WATER QUALITY OF LAKE WARAMAUG AND SURROUNDING WATERSHED, LITCHFIELD COUNTY, CONNECTICUT

by Kenneth P. Kulp and David Grason

U.S. GEOLOGICAL SURVEY
Water-Resources Investigations Report 85-4267

Prepared in cooperation with the

LAKE WARAMAUG TASK FORCE NORTHWEST CONNECTICUT REGIONAL PLANNING AGENCY



Hartford, Connecticut 1992

U.S. DEPARTMENT OF THE INTERIOR

MANUEL LUJAN, JR., Secretary

U.S. GEOLOGICAL SURVEY

Dallas L. Peck, Director

For additional information write to:

Chief, Connecticut District U.S. Geological Survey, WRD 450 Main Street, Room 525 Hartford, CT 06103 Copies of this report can be purchased from:

U.S. Geological Survey Books and Open-File Services Section Federal Center, Box 25425 Denver, CO 80225

CONTENTS

Page	-
Introduction	24457831
ILLUSTRATIONS	
Page	_
·	-
2. Surface-water inflow and atmospheric-deposition sampling sites.	3
TABLES	
Page	9
Table 1. Selected physical characteristics of Lake Waramaug	2
descriptions	5 7
descriptions)
descriptions	
7. Chemical, physical, and biological characteristics of surface-water inflows	
8. Chemical and physical characteristics of atmospheric deposition	6
9. Chemical and physical characteristics of ground water and seepage	7
10. Chemical and physical characteristics of lake bottom material. 18 11. Chemical, physical, and biological characteristics of lake water	

CONVERSION FACTORS AND ABBREVIATIONS

Multiply inch-pound unit	Ву	To obtain metric unit
inch (in.)	25.4	millimeter
foot (ft)	0.3048	meter
pound`(lb)	0.4536	kilogram
acre	0.4047	hectare
mile (mi)	1.609	kilometer
square mile (mi²)	2.590	square kilometer
cubic foot per second (ft ³ /s)	0.02832	cubic meter per second
ton per day (ton/d)	907.2	kilogram per day
gallon per minute (gal/min)	0.0631	liter per second
pound per acre (lb/acre)	1.1218	kilograms per hectare
degree Fahrenheit (°F)	5/9 (°F-32)	degree Celsius

WATER QUALITY OF LAKE WARAMAUG AND SURROUNDING WATERSHED, LITCHFIELD COUNTY. CONNECTICUT

by Kenneth P. Kulp and David Grason

ABSTRACT

Lake Waramaug and its watershed in western Connecticut were sampled from March 1977 to March 1978 to develop information for a lake-management plan. Nutrient enrichment has degraded the lake water quality, resulting in an increased algal population in recent years. Chemical analyses of surface-water inflow, ground-water inflow, and atmospheric deposition in the watershed indicate that surface-water inflow at the northeastern corner of the lake is the major source of nutrients discharged to the lake. Atmospheric deposition contains 0.01 to 0.47 milligrams per liter total phosphorus and 0.52 to 3.2 milligrams per liter total nitrogen. During the 7.3-month period of investigation, atmospheric deposition contributed 61,400 pounds of nitrogen and 3,150 pounds of phosphorus to the lake's watershed. Nutrient concentrations in ground water were relatively low, with total phosphorus ranging from 0.008 to 0.14 milligrams per liter.

Lake bottom materials contain high concentrations of nitrogen and phosphorus which seem to be released into the hypolimnion when anaerobic conditions develop. These anaerobic conditions exist in the hypolimnion during most of the summer when the lake develops strong thermal stratification. The ultimate fate of these nutrients was not determined, although the nitrogen and phosphorus seem to be confined to the hypolimnion during the summer stratification.

During the summer, the dominant phytoplankton in the lake are blue-green algae. During the spring and fall, golden-brown algae are the dominant phytoplankton. No severe algal blooms occurred in the lake during this investigation.

INTRODUCTION

Lake Waramaug, in western Connecticut, has had increased algal populations in recent years, indicative of nutrient enrichment. Norvell and Frink (1975) identified the lake as one where accelerated eutrophication has occurred during the past 35 to 40 years. In a subsequent report by the King's Mark Resource Conservation and Development Area's Environmental Review Team (1976), it was recommended that a more extensive investigation of the lake be made to define specific causes and sources of the accelerated eutrophication.

In March 1977, the U.S. Geological Survey, in cooperation with the Lake Waramaug Task Force of the Northwestern Connecticut Regional Planning Agency, initiated an investigation of selected water-quality characteristics in Lake Waramaug and its watershed. The purpose of this report is to present data from this investigation that will be used in conjunction with the information obtained from other studies in the development of a lakemanagement program.

The main objectives of the investigation were to identify the sources of the nutrients which are causing the lake's accelerated eutrophication, to obtain a water-quality baseline of the lake throughout the growing season, and to determine areal and temporal variations in nutrient concentrations within the lake and its watershed. To achieve these objectives, samples were collected from (1) the lake, (2) surface-water inflow and outflow, (3) selected ground-water sources, and (4) atmospheric deposition within the watershed, and analyzed for chemical and physical characteristics.

Acknowledgment is made to Carol Youell and William H. Vogel, Jr., of the Lake Waramaug Task Force, who assisted in the collection of data during this investigation.

DESCRIPTION OF THE STUDY AREA

Lake Waramaug is located in Litchfield County in western Connecticut in the townships of Kent, Warren, and Washington (fig. 1). The lake has a watershed of 14.07 mi² (square miles), of which about 66 percent is forested, 10 percent is agricultural, 5 percent is wetland, 5 percent is residential or commercial, and the remaining 14 percent is composed of fields, recreational areas, or open-water (King's Mark, 1976). The land immediately adjacent to the lake is primarily residential, with the exception of that area along the northwestern end which is encompassed by Lake Waramaug State Park and the Arrow Point area. This area consists mainly of forests and fields.


Lake Waramaug is of glacial origin, and is the second largest naturally formed lake in the State. It is used primarily for recreational purposes, including boating, fishing, and swimming. The primary inflow is Lake Waramaug Brook, locally known as Sucker Brook, which discharges into the northeastern corner of the lake. Several small streams enter at various points around the lake, some of which are intermittent. The outflow is located at the southeastern corner and forms the East Aspetuck River, which is a part of the Housatonic River drainage system.

The Lake Waramaug watershed is underlain by the Waramaug Formation, which consists of mica-quartz gneisses and schists, feldspathic mica quartzites, and mica quartzites (Gates and Bradley, 1952). The surficial geology of the area is mainly glacial till, with numerous rock outcrops and some small alluvial fan deposits near the the lakeshore (Colton, 1969).

Selected physical characteristics of the lake are presented in table 1. The bathymetry and aquatic vegetation of the lake are described by Kulp (1981).

Table 1.--Selected physical characteristics of Lake Waramaug [King's Mark Environmental Review Team, 1976]

Surface area	672 acres
Maximum depth	40 feet
Mean depth	22.1 feet
Volume Average retention time	647 × 10 ⁶ cubic feet 302 days






Figure 1.--Location of study area.

DATA COLLECTION

Because data collection consisted of sampling various systems of the lake and its watershed, separate programs were required. Unless otherwise indicated, the techniques used for the collection and analysis of chemical samples were in accordance with those described by Brown and others (1970). The algal growth potential determinations and bacteriological analysis were done by the techniques described by Greeson and others (1977).

Surface-Water Inflow Sampling

Twenty-seven surface-water sites were sampled in the lake's drainage. The location of these sampling sites is shown in figure 2; descriptions are given in table 2.

Discharge measurements were made at each site at the time of the sampling, using techniques described by Buchanan and Somers (1969). At some sites where flow was very low, measurements were made by timing discharge into a graduated container. Field determinations of water temperature, specific conductance, pH, dissolved-oxygen concentration, fecal coliform, fecal streptococcus, and total coliform bacteria were made at each site. Water samples were analyzed by the U.S. Geological Survey Central Laboratory for constituents that included dissolved solids, organic carbon, suspended sediment, nitrogen, phosphorus, and algal growth potential.

Specific amounts and dates of precipitation during the sampling period are given in table 3. The initial sampling of surface-water inflow took place on March 14, 1977. This coincided with a rain storm which began on March 13, and continued through March 14. Melting snow, in conjunction with the storm, caused streamflow to be high. The second sampling was conducted on June 1 and 2, coinciding with a light rain following a long, dry period. Streamflow was low at the time of sampling, and some smaller streams were dry. The third sampling was conducted on June 7, when weather and streamflow were similar to that of the June 1 and 2 sampling.

On September 20, the fourth sampling was conducted during a light rain which had begun on the previous day. Streamflow and runoff were again low. Based on an evaluation of previously collected data, laboratory analyses were modified at this time. Dissolved-solids concentration was not determined because a satisfactory correlation with specific conductance had been established, and analysis of orthophosphate was replaced by analysis of dissolved phosphorus, which was felt to be more directly related to algal growth in the lake. Analysis of temperature, pH, dissolved-oxygen concentration, bacteria, algal growth potential, and nitrogen were reduced to once per site.

The fifth sampling was conducted on October 20, following several days of rain when streamflow was above average. The sixth sampling was conducted during heavy rain on November 8, when streamflows were again high.




Figure 2.--Surface-water inflow and atmospheric deposition sampling sites.

Table 2.--Surface-water inflow sampling site identification and descriptions

Site	USGS station	ما مهمه ا	ما مسمع المسلم	Doggistics on semants
number	number	Latitude	Longitude	Description or remarks
I1	414151073204000	41°41'51"	73°20'40 "	Small unnamed brook just south of private road leading to Boulder's Inn
12	01201030	41°42'02"	73°20'50"	Lake Waramaug Brook near New Preston at bridge on private drive
13	414207073210400	41°42'07"	73°21'04"	·
14	414210073211800	41°42'10"	73°21'18"	
15	414210073213000	41°42'10"	73°21'30"	
16	414206073214500	41°42'06 "	73°21'45"	
17	01201010	41°43'23"	73°20'30"	
18	414210073221200	41°42'10"	73°22'12"	Largest of several culverts under Lake Street just east of Arrow Point that carries water to the lake during heavy runoff
19	414229073224500	41°42'29 "	73°22'45"	Unnamed brook just west of Arrow Point
I10	414234073230300	41°42'34 "	73°23'03"	Small unnamed brook flowing into the northwestern corner of the lake
I11	414223073225900	41°42'23"	73°22'59"	Small unnamed brook flowing from the middle of Lake Waramaug State Park
I12	414157073224700	41°41'57"	73°22'47 "	Culvert flowing under road at the Washington-Kent town line. Culvert flows only during heavy runoff
I13	414146073224300	41°41'46"	73°22'43"	Culvert from Golf Course Hill under West Shore Road. Flows only during heavy runoff periods
I14	414145073224000	41°41'45"	73°22'40"	•
I15	414120073224700	41°41'20"	73°22'47 "	·
I16	414136073228900	41°41'36 "	73°22'29 "	Small culvert under West Shore Road. Flows only during heavy runoff
I 17	414133073222300	41°41'33"	73°22'23"	Small unnamed brook entering the southwestern corner of the lake
I18	414144073215900	41°41'44"	73°21'59 "	Small culvert under West Shore Road. Flows only during heavy runoff
I 19	414141073215000	41°41'41"	73°21'50"	Small culvert under West Shore Road. Flows only during heavy runoff
120	414143073213100	41°41'43"	73°21'31"	Small culvert from Tinker Hill under West Shore Road. Flows only during heavy runoff
121	414119073212200	41°41'19"	73°21'22"	Small culvert flowing under West Shore Road at boat landing
122	414102073211400	41°41'02"	73°21'14"	Flow from rock cliffs above West Shore Road
123	01201050	41°40'57 "	73°21'15"	East Aspetuck River at New Preston at lake outlet
124	414121073205800	41°41'21"	73°20'58"	Flow from rock cliffs above State Route 45
125	414133073205000	41°41'33"	73°20'05"	Flow from rock cliffs above State Route 45
126	01201020	41°43'10"	73°19'56"	Lake Waramaug Brook near Warren at State Route 341
127	414237073204600	41°42'37"	73°20'46"	Lake Waramaug Brook at State Route 45

Table 3.--Precipitation data, U.S. Geological Survey station 414232073210900 [Data provided by William H. Vogel, Jr.]

	· · · · · · · · · · · · · · · · · · ·			Precip	itation	in incl	nes				
Day	Mar. 1977	Apr. 1977	May 1977	June 1977	July 1977	Aug. 1977	Sept. 1977	Oct. 1977	Nov. 1977	Mar. 1978	Apr. 1978
1 2				0.42				0.70 0.10			
3	0.25	0.60				0.25				0.60 ^a	
5 6 7	0.60	0.70	0 .6 5	0.50		0.65 1.00		0.10			0.15 0.10
3 4 5 6 7 8 9		0.10	1.25	1.10	0.15	1.00		1.10 0.10	1.15 0.25		
11 12 13 14	0.85				0.35	0.10 0.30	0.55	0.50	0.25	0.60	0.20
14 15 16	0.70 0.10					0.20		0.50 0.15	0.30	0.60 0.10 6.00 ^a	
17	6 as 8				0.20	0.30	0.80	0.25	0.25		- 10
18 19	0.60 ^a			0.80			0.25	0.25			0.10 0.90
20	0.20 ^a			0.80			0.25 0.70	0.15 0.65			0.10
21 22				0.50			0.70	0.65	0.15	0.10	0.10
23 24 25	1.60 ^a	1.40			1.20	0.15	0.65 0.50		0.10		
26 27							1.40	0.20 0.10	0.70	0.40 ^a 0.30	
28 29	0.40			0.20							
30 31	0.10										

Approximate value (precipitation was snow which was converted to approximate rainfall equivalent by dividing by 10).

In the spring of 1978, after all the collected data were evaluated, a final sampling was planned. Because phosphorus was thought to be the most important constituent, the sampling was designed to measure its concentrations extensively at several sites during a spring freshet. Sites I2, I7, I9, and I26 (table 2) were selected for extensive sampling. Staff gages were installed at these sites to measure water stage at the time of sampling and to develop a stage-discharge relation. Sampling was conducted at these sites from March 20 to 31, 1978, coinciding with a period of light rains and some snow. On March 27, the small amount of snow still on the ground was melted by rain, creating high streamflow. During this day, most sites were sampled, and multiple samples were collected at the four extensive sampling sites. The samples were analyzed for phosphorus concentration, algal growth potential, and suspended-sediment concentration. Temperature, pH, and specific conductance were measured on site at the time of sampling.

Atmospheric-Deposition Sampling

One site, near the town of Warren (fig. 2), was established for the collection of wet and dry atmospheric deposition. The samples were collected using a bulk-type sampler, consisting of a polyethylene funnel fitted to a glass bottle. Data collection began April 1, 1977, and continued through November 9, 1977. The first sample was a composite of atmospheric deposition for the entire month of April. Subsequently, samples were collected for analysis after every major rainstorm. For samples collected prior to August 24, 1977, a plastic wool material was placed in the collector to filter out large objects such as leaves and twigs; thereafter, a copper screen was used for this purpose. Laboratory analyses verified that neither of these filter materials contributed significant phosphorus to the sample. The plastic wool may have contributed nitrogen in the range of 0.01 to 0.10 mg/L (milligrams per liter) to the sample. The copper screen had no effect on nitrogen concentration.

All samples were taken from the collector and immediately refrigerated. Specific conductance and pH were determined upon receipt of the samples by the U.S. Geological Survey which in some instances was several days after collection. The samples were subsequently sent to the laboratory for nitrogen and phosphorus analysis. The quantities and dates of precipitation over the collection period were recorded by a local resident, William H. Vogel, Jr. These data are shown in table 3.

Ground-Water Sampling

Fourteen wells and springs in the vicinity of Lake Waramaug and four seepage sources on the lake bottom near shore were sampled to determine the contribution of nutrients by ground water to the lake. The location of these sampling sites is shown in figure 3; descriptions are given in table 4. Samples of ground-water seepage within the lake were collected using the samplers and techniques described by Lee (1977), and were analyzed for nitrogen and phosphorus. Ground-water samples from the wells and springs were also analyzed for these nutrients, and measurements of specific conductance, pH, and temperature were made at the time of sampling.




Figure 3.--Ground-water and seepage sampling sites.

Table 4.--Ground-water and seepage sampling site identification and descriptions [m, meter; μ m/s, micrometer per second]

Site	USGS station			
number	number	Latitude		Description or remarks
			Ground	-water sites
G1	414202073205901	41°42'02 "	73°20'59 "	Dug well with concrete sleeves, approximately 2.5 m deep and 1.0 m in diameter.
G2	414157073205801	41°41'57 "	73°20'58"	Dug well, approximately 10 years old. Approximatel 2.5 m deep and 1 m in diameter.
G3	414159073215201	41°41'59"	73°21'52"	Dug well with gravel bottom, approximately 2.4 m deep and 1 m in diameter.
G4	414230073212201	41°42'30"	73°21'22 "	Spring in pasture. Stone holding tank approximatel 1.5 m deep.
G5	414207073210701	41°42'07"	73°21'07 "	Dug well with concrete sleeves, approximately 40 years old. Approximately 3.4 m deep and 1 m in diameter. Sand and gravel bottom
G6	414232073231001	41°42'32"	73°23'10"	Drilled well, 66 m deep. In field near showers in Lake Waramaug State Park
G 7	414219073230201	41°42'19	73°23'02"	Drilled well, 34 m deep. Near campsites in Lake Waramaug State Park
G8	414214073230001	41°42'14"	73°23'00"	Drilled well, 36 m deep. In back of campsites in Lake Waramaug State Park
G9	414271073225801	41°42'17"	73°22'58 "	Drilled well, 53 m deep. On lakeside of road in Lake Waramaug State Park
G10	414236073225301	41°42'36 "	73°22'53"	Artesian well
G11	414122073212001	41°41'22 "	73°21'20"	Drilled well approximately 3 years old and 47 m deep. Yield of approximately 10 gal/min. Below 4 m well was drilled through Cambrian rock.
G12	414138073213501	41°41'38"	73°21'35 "	Spring with concrete holding tank.
G13	414139073213701	41°41'39 "	73°21'37"	Spring approximately 0.6 m deep. Concrete holding tank
G14	414140073224801	41°41'40"	73°22'48"	Dug well, approximately 25 years old, approximately 5 m deep.
	•		Seep	page sites
S 1	414155073204300	41°41'55"	73°20'43"	Seepage rate of 3.4 μ m/s, approximately 8 m from shore.
S 2	414142073214200	41°41'42"	73°21'42"	Seepage rate of 0.2 μ m/s, approximately 6 m from shore.
\$3	414146073220000	41°41'46"	73°22'00"	Seepage rate of 0.1 $\mu \mathrm{m/s}$, approximately 12 m from shore.
S 4	414156073205000	41°41'56"	73°20'50"	Seepage rate of 0.05 $\mu m/s$, approximately 20 m from shore.

Lake-Bottom-Material Sampling

On May 4 and 5, 1977, lake-bottom-material samples were collected from 13 points in the lake using a ponar dredge. The locations of these sampling sites are shown on figure 4; descriptions are given in table 5. The samples were analyzed for concentrations of nitrogen, phosphorus, and organic carbon, and particle-size distribution.




Figure 4.--Lake-bottom-material sampling sites.

Table 5.--Lake-bottom-material sampling site identification and descriptions [m. meter]

Site	USGS station		,	
number	number	Latitude	Longitude	Description or remarks
B1	414157073205400	41°41'57"	73°20'54"	At mouth of Lake Waramaug Brook
B2	414205073210900	41°42'05"	73°21'09"	At mouth of small unnamed brook on northeast side of
				lake
В3	414209073212500	41°42'09 "	73°21'25" lake	Approximately 20 m from shore on northeast side of
B4	414225073224500	41°42'25"	73°22'45"	At mouth of unnamed brook at northwest end of lake
B5	414225073225000	41°42'25"	73°22'50 "	At site L2, approximately 100 m from shore at
	northwest end of	lake		
B6	414225073225700	41°42'25"	73°22'57"	At mouth of brook entering from Lake Waramaug State Park
В7	414145073224000	41°41'45	73°22'40"	At site I14, at end of pipe connecting to Ash Swamp
B8	414147073215600	41°41'47"	73°21'56"	Approximately 30 m from shore on south side of Lake
В9	414152073214300	41°41'52"	73°21'43"	At site L8, approximately 300 m from Randalls Point
B10	414145073213100	41°41'45"	73°21'31"	Approximately 30 m from shore on south side of Lake
B11	414119073212200	41°41'19"	73°21'22"	At site L21, mouth of culvert entering southeast end of lake
B12	414121073211000	41°41'21"	73°21'10"	At site L14, approximately 200 m from shore at southeast end of lake
B13	414104073211000	41°41'404	73°21'10"	At site L16, near outflow at southeast end of Lake

Lake-Water Sampling

Initially, lake-water samples were collected at 16 sites. The locations of these sites are shown in figure 5; descriptions are given in table 6. At each site, field measurements of secchi disk transparency, water temperature, pH, specific conductance, and dissolved-oxygen concentration were made at various depths through the water column. Samples were collected at various depths at six mid-lake sites (L2, L5, L8, L11, L14, and L16), using a point sampler. These samples were analyzed for alkalinity, nitrogen, and phosphorus. Samples collected in April and August 1977 were also analyzed for total organic carbon and algal growth potential. Phytoplankton samples were collected at each mid-lake site, and the dominant types were identified.

Lake sampling began on April 4, 1977, and continued until April 13, 1978. Analysis of results obtained through June 1977 indicated that the water quality of nearshore sites was essentially the same as that found in the upper levels of the mid-lake sites. For this reason, the nearshore sites (L1, L3, L4, L6, L7, L9, L10, L12, L13, and L15) were eliminated from the sampling program beginning with the July 1977 sampling. The analysis of orthophosphorus was also eliminated at this time, as it was felt to be of doubtful utility. During the two rounds of sampling in October 1977, analysis of iron concentration was added to the laboratory determinations. The November 16, 1977 sampling was limited to field determinations, because the growing season had ended. The April 13, 1978 sampling was limited to sites L5, L8, and L14, and only the field determinations and analysis for phosphorus were performed. This final sampling took place just after the ice had melted from the lake.




Figure 5.--Lake sampling sites.

Table 6.--Lake sampling site identification and descriptions

Site	USGS station			
number	number	Latitude	Longitude	Description or remarks
L1	414227073224800	41°42'48"	73°22'48"	Field values only. Discontinued in June 1977
L2	414225073225000	41°42'25"	73°22'50"	Field and laboratory values
L3	414224073225500	41°42'40"	73°22'55"	Field values only. Discontinued in June 1977
L4	414149073222400	41°41'49"	73°22'24 "	Field values only. Discontinued in June 1977
L5	414144073222400	41°41'44"	73°22'24"	Field and laboratory values
L6	414136073222400	41°41'36"	73°22'24"	Field values only. Discontinued in June 1977
L7	414200073214900	41°42'00"	73°21'49 "	Field values only. Discontinued in June 1977
L8	414152073214300	41°41'52"	73°21'43 "	Field and laboratory values
L9	414145073213700	41°41'45 "	73°21'37 "	Field values only. Discontinued in June 1977
L10	414155073205000	41°41'55 "	73°20'50"	Field values only. Discontinued in June 1977
L11	414150073205400	41°41'50"	73°20'54 "	Field and laboratory values
L12	414144073210100	41°41'44 "	73°21'01"	Field values only. Discontinued in June 1977
L13	414121073210100	41°41'21 "	73°21'01"	Field values only. Discontinued in June 1977
L14	414121073211000	41°41'21"	73°21'10"	Field and laboratory values
L15	414121073211700	41°41'21"	73°21'17 "	Field values only. Discontinued in June 1977
L16	414104073211000	41°41'04"	73°21'10"	Field and laboratory values

RESULTS AND CONCLUSIONS

Surface-Water Inflows

Results of surface-water-inflow analyses are listed in table 7 (at back of report). Fecal coliform and fecal streptococcus bacteria populations were relatively high at sites I2, I3, I4, I6, I11, I14, I19, I26, and I27. These bacteria are typically found in the fecal wastes of warm-blooded animals, and their presence in large numbers in water is indicative of fecal pollution. In general, the ratio of the number of fecal coliform bacteria to that of fecal streptococcus bacteria at these sites was less than 0.7, indicating nonhuman sources, such as dogs, cats, rodents, livestock, and poultry (Geldreich and Kenner, 1969). In some instances, however, the ratios were higher, indicating possible human fecal contamination. Ratios exceeding 2, which indicate a human source of pollution, were found at sites I4, I14, and I19.

In general, the streams along the northeastern shore of the lake have the highest nutrient concentrations and have the highest algal growth potential values. Lake Waramaug Brook, the largest stream entering the lake, flows through this area, and is the largest stream source of nutrients discharging to the lake. Sites I3 and I4, located in the same general area, had consistently high phosphorus concentrations. A large fraction of the phosphorus was in a dissolved form, which is readily available for use by algae. This is reflected in the high algal growth potential recorded at these sites. Although these streams are both small, their contribution of nutrients to the lake is significant.

Sites located along the steeper slopes of the lake's eastern shoreline (I22, I24, and I25) had low pH and specific conductance values, characteristic of the values recorded from the atmospheric deposition samples. Nutrient concentrations were low at these sites, and it seems that the nutrient contribution from these inflows is small and results directly from precipitation.

Nutrient contributions from streams discharging to the western-most finger of the lake generally are low, although site III occasionally had moderate algal growth potential values. Because this area of the lake is relatively shallow (approximately 15 feet) and is isolated by Arrow Point, the nutrient loading from these streams, particularly III, may have significant local impact. In recent years, this area has experienced more algal blooms than other areas of the lake, although none were observed during this investigation.

Results of the March 1978 sampling indicate that the greatest quantity of phosphorus enters the lake during periods of high surface-water flow. Data indicate that sharp increases in phosphorus concentrations occur during the initial rise of the stream. Concentrations of phosphorus begin to decrease gradually, as peak flow is reached and recession begins. For example, data from site I2 (table 7) indicate that on March 26, 1977, when the streamflow was 30 ft 3 /s (cubic feet per second), the total phosphorus concentration was 0.01 mg/L. Rain increased the streamflow to 245 ft 3 /s on March 27, and the phosphorus increased to 0.44 mg/L. As streamflow increased to 325 ft 3 /s, the phosphorus concentration increased to 0.50 mg/L. During the following 2 hours, streamflow remained constant, but the phosphorus concentration decreased to 0.25 mg/L. On the following day, when the streamflow had decreased to 116 ft 3 /s, the phosphorus concentration was 0.04 mg/L. Similar streamflow-phosphorus relations were observed at the other three intensively sampled sites.

Atmospheric Deposition

The results of chemical analyses of atmospheric deposition are given in table 8. Specific conductance values ranged from 12 to 72 μ S/cm (microsiemens per centimeter at 25 degrees Celsius). Concentrations of the nutrients nitrogen and phosphorus varied, and occasionally were relatively high. Total phosphorus concentrations ranged from 0.01 to 0.47 mg/L and total nitrogen concentrations ranged from 0.50 to 3.2 mg/L. In several instances, the nutrient concentrations found in some of the inflows were similar to those of the atmospheric deposition samples for the same time period. It is possible that a substantial portion of the nutrients found in these streams originated from atmospheric deposition. During the data-collection period, April 1 to November 9, 1977, atmospheric deposition contributed approximately 6.28 lb/acre (pounds per acre) of nitrogen and 0.35 lb/acre of phosphorus. These amounts are equivalent to 61,400 lb (pounds) of nitrogen and 3,150 lb of phosphorus from the atmosphere during this 7.3-month period, which is a significant amount.

Table 8.--Chemical and physical characteristics of atmospheric deposition [Results are in milligrams per liter (mg/L) except as indicated; μ S/cm, microsiemens per centimeter]

Sample	Precipi- tation	Specific	Total nitrogen	Total Kjeldahl nitrogen	Total nitrate + nitrate	•		Total nitrogen as N deposited on	Total phosphorus as P deposited on
collection	quantity	conductance	as N	as N	as N	as NO ₃	phosphorus	lake surface a/	lake surface <u>a</u> /
period	(inches)	(µS/cm)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	as P	(pounds)	(pounds)
04-01-77/									
05-01-77 05-01-77/	2.80	35	0.94	0.44	0.50	4.2	0.010	401	4.26
05-05-77	.65	22	1.0	.66	.36	4.5	.010	99	0.99
05-05-77/ 05-11-77	1.25	25	1.0	.56	.45	4.5	.110	190	20.9
05-11-77/									
06-02-77 06-02-77/	.42	41	3.2	2.4	.78	14	.470	205	30.1
06-02-77	1.60	28	.70	.30	.40	3.1	.030	171	7.31
06-11-77/ 06-21-77	1.30	65	2.2	1.1	1.1	9.7	.030	436	5.94
06-21-77	1.50	03	2.2	1.1	1.1	3.7	.030	430	3.34
07-25-77 07-25-77/	2.10	63	1.6	.75	.84	7.0	.024	512	7.68
08-10-77	2.90	72	1.7	.68	1.0	7.4	.020	751	8.83
08-10-77/ 08-24-77	1.05	53	1.2	.53	.64	5.2	.09 0	192	14.4
08-24-77	1.05	33	1.2	.55	.04	5.2	.090	192	14.4
09-13-77 09-13-77/	0.55	61	2.2	1.3	.90	9.7	.211	184	17.7
09-13-77	1.75	42	1.7	1.1	.63	7.7	.169	453	45.0
09-21-77/	0 55	10	00	E1	21	2.6	100	210	20.6
09-26-91 09-26-91/	2.55	19	.82	.51	.31	3.6	.102	318	39.6
10-01-77	.70	35	1.5	.94	.53	6.5	.056	160	5.97
10-01-77/ 10-10-77	1.40	13	.64	.52	.12	2.8	.023	136	4.90
10-10-77/	1 05	24	63	22	0.4	0.5	0.47	100	0.05
10-18-77 10-18-77/	1.25	34	.57	.33	.24	2.5	.047	109	8.95
10-20-77	.80	15	1.1	.46	.65	4.9	.089	134	10.8
10-20-77/ 11-09-77	1.70	12	.52	.34	.18	2.3	.020	135	5.18
04-01-77/								4,586	238.51
11-09-77									

 $[\]underline{\underline{a}}/$ Calculation based on lake surface area of 672 acres.

Ground Water and Seepage

Results of ground-water and lake-seepage sample analyses are given in table 9. Phosphorus concentrations in ground water were greatest at sites G1 and G10, with concentrations of 0.136 and 0.102 mg/L, respectively. Concentrations of phosphorus at other ground-water sites ranged from 0.008 to 0.074 mg/L. Without a more thorough study of ground-water flow to the lake, it is not possible to determine the nutrient contribution of ground water to the lake, although it appears to be minimal, compared to other sources.

With the exception of station S1, lake seepage data showed values similar to those found in bottom waters during the same time period. Based on the limited seepage and other ground-water data collected, seepage does not appear to be contributing a significant quantity of nutrients to the lake.

Table 9.--Chemical and physical characteristics of ground water and seepage [Results are in milligrams per liter (mg/L) except as indicated; μ S/cm, microsiemens per centimeter; °C, degrees Celsius]

Site		Specific conductance	pH (units)	Temper- ature	Total nitrite +nitrate as N	Dissolved ammonia nitrogen as N	Total organic nitrogen as N	Total nitrogen as N	Total nitrogen as NO ₃	Total phosphorus as P
number	Date	(µS/cm)	(units)	(°C)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
G 1	08-25-77	360	5.9	15.0	21	0.00	0.15	21	94	0.136
G2	08-25-77	360	7.6	17.0	.07	.01	.65	.73	3.2	.030
G3	08-25-77	285	6.7	16.0	.06	.01	.14	.21	.93	.014
G4	08-25-77		5.8	10.0	1.0	.08	.51	1.6	7.0	.045
G5	08-25-77		6.8	14.0	.35	.00	.04	.39	1.7	.043
G6	08-25-77	265	7.7	11.0	.03	.01	.03	.07	.31	.010
G7	08-25-77	250	7.0	13.5	1.1	.00	.00	1.1	4.9	.036
G8	08-25-77	185	6.8	13.0	.07	.00	.02	.09	.40	.014
G9	08-25-77	250	7.6	15.0	.08	.00	.05	.13	.58	.042
G10	08-25-77	87	7.1	13.0	.06	.00	.21	.27	1.2	.102
G11	08-25-77		6.6	20.0	.13	.00	.00	.13	.58	.008
G12	08-25-77		6.6	14.0	.05	.00	.21	.26	1.2	.046
G13	08-25-77	87	6.9	14.0	.08	.01	.08	.17	.75	.053
G14	08-25-77	295	7.2	15.0	1.4	.00	.16	1.6	6.9	.074
S 1	08-02-77	_	_	-	.18	.03	.24	.45	2.0	.032
S2	08-02-77	-	_	-	.02	.21	.74	.97	4.3	.049
S2	08-04-77	-	-	_	.01	.21	.99	1.2	5.4	.061
\$3	08-11-77	_	-	-	.01	.20	1.0	1.2	5.4	.047
S4	08-16-77	_	_	_	.08	.34	.76	1.2	5.2	.060

Lake-Bottom Materials

The data from lake bottom-material sampling are given in table 10. The highest concentration of nutrients was found at the deepest site (B12). In general, nutrient concentrations are higher at deeper sites than at the shallower sites. This may be related to the characteristics of the bottom materials, which are finer and contain higher concentrations of organic carbon at the deeper sites than at shallower sites.

The data indicate that under anaerobic conditions, nutrients are released from the bottom material into the water of the lake. The extent to which this occurs and the ultimate fate of these nutrients are unknown.


Table 10.--Chemical and physical characteristics of lake-bottom material

[ft, feet; mg/kg, milligrams per kilogram; g/kg, grams per kilogram; mm, millimeter]

Site number	Date	Sampling depth (ft)	Total nitrogen in bottom material as N (mg/kg)	Total phosphorus in bottom material as P (mg/kg)	Organic carbon in bottom material as C (g/kg)	Bed material fall diameter (percent finer than 0.004mm)	Bed material fall diameter (percent finer than 0.062mm)	Bed material sieve diameter (percent finer than 2.00 mm)
B1	05-05-77	2.0	3,000	850	10	8	21	100
B2	05-05-77	2.0	12,000	790	28	12	29	100
B3	05-04-77	20	23,000	930	56	26	81	100
B4	05-04-77	2.0	2,200	430	7.6	5	11	100
B5	05-04-77	15	34,000	1,000	94	37	91	100
B6	05-05-77	2.0	4,500	980	18	5	13	100
B7	05-04-77	2.0	1,700	100	65.0	3	8	100
B8	05-04-77	25	22,000	990	51	27	85	100
В9	05-04-77	27	33.000	1,100	75	39	95	100
B10	05-04-77	3.0	730	700	2.2	1	8	100
B11	05-04-77	2.0	1,200	95	2.8	0	3	100
B12	05-04-77	40	42,000	5,000	103	56	86	100
B13	05-04-77	10	4,300	990	15	7	22	100

Lake Water

Results of lake water analyses are given in table 11 (at back of report). The data indicate that the lake begins to stratify thermally in late April. By mid-June, the lake is well stratified, and dissolved oxygen levels are near zero in the hypolimnion. Stratification continues through early September, with anaerobic conditions existing in regions of the lake below 20 to 24 ft. Stratification begins to break down in mid-September and the lake is essentially homothermous by early October. Figure 6 shows temperature and dissolved-oxygen profiles during the period of study at site L8.



DISSOLVED OXYGEN, IN MILLIGRAMS PER LITER AND TEMPERATURE, IN DEGREES CELSIUS

Figure 6.--Temperature and dissolved-oxygen profiles at lake station L8.

Nutrient data (table 11) show phosphorus, and often nitrogen concentrations, are usually highest near the lake bottom. This is most evident during the summer when the lake is stratified. It is not observed during turnover in April or October. The nutrient concentrations begin to increase near the bottom of the lake in June. They continue to increase through early September, then they begin to decrease rapidly. The June increase is probably due to reducing reactions that take place in the bottom sediments under anaerobic conditions, releasing nutrients to the surrounding water. Because the lake is stratified at this time, little mixing of water takes place, and nutrients are not made available to support algal growth near the lake surface. This is reflected in the nutrient concentrations in water near the surface of the lake, which appear to react independently of the concentrations at the bottom water during this period. Surface nutrient concentrations during the summer growing period are probably controlled to a greater extent by precipitation and inflows than by mixing with other levels of the lake. As stratification begins to break down in September, mixing takes place and nutrient-rich waters near the bottom reach the surface. This produces a marked decrease in nutrient concentrations at the bottom. and an increase in concentrations at the surface.

Algal populations in the lake varied during the 1977 growing season. Golden-brown algae (Chrysophyta) predominated in April, but blue-green algae (Cyanophyta) predominated by mid-May and continued throughout the summer. They were replaced by golden-brown algae and euglenoids (Euglenophyta) in October. It is notable that the change in predominant algal types is concurrent with the spring and fall turnover of the lake. This may be related to the release of silica from bottom sediments into the surface of the lake during turnover. Silica is essential to the growth of certain golden-brown algae, in particular, the diatoms. The change in dominance of algal types may also be controlled by nitrogen concentrations. The change in May and June occurred in conjunction with a sharp decrease in surface nitrogen concentrations. Blue-green algae are capable of using atmospheric nitrogen in place of high nitrogen concentrations in the water, essential to most other types of algae. The decrease in nitrogen concentrations would thus impede the production of other algal types, while having no effect on blue-green algae. In late September and October, nitrogen concentrations increased in surface waters and dominance reverted back to golden-brown algae, along with euglenoids. No serious algal blooms were observed during the 1977 sampling period, and the lake was said to have had a "good year" by local residents.

SUMMARY

Data collected from March 1977 to March 1978 indicate that the major sources of nutrients to the lake are surface-water inflows at the northeastern corner of the lake, with Lake Waramaug Brook being the largest single contributor. The highest nutrient concentrations occur in streams during periods of high flow. Bacteriological data from surface-water inflows frequently indicate high levels of fecal contamination, primarily from nonhuman sources.

Atmospheric deposition in the lake watershed contains 0.01 to 0.47 mg/L total phosphorus and 0.52 to 3.2 mg/L total nitrogen, and may be a significant source of nutrients to the lake. During the period April 1 to November 9, 1977, atmospheric deposition contributed about 61,400 lb of nitrogen and 3,150 lb of phosphorus to the lake watershed.

Ground-water and in-lake seepage do not appear to contribute a significant quantity of nutrients to the lake, although a more thorough study of ground-water flows and quality is necessary to confirm this.

Bottom materials from deep sections of the lake contain higher concentrations of nutrients and organic carbon, and are of finer size than those from shallower sites in the lake. In general, lake bottom materials contained high concentrations of nutrients.

Lake Waramaug was found to develop strong thermal stratification during the summer months, with anaerobic conditions existing in the hypolimnion during most of this period. Nutrients in the lake were also found to concentrate at specific levels during stratification. The highest nutrient concentrations were located near the lake bottom, possibly because of the release of nutrients from bottom materials during anaerobic conditions. These nutrients do not appear to mix with water in other levels of the lake until the fall turnover. Blue-green algae predominate in the lake during the summer, with golden-brown algae predominating in the spring and fall.

Lake morphology probably contributes to the lake's history of local algal blooms in the northwestern finger, although no significant blooms were observed during the period of this study.

CITED REFERENCES

- Brown, Eugene, Skougstad, M.W., and Fishman, M.J., 1970, Methods for collection and analysis of water samples for dissolved minerals and gases: U.S. Geological Survey Techniques of Water-Resources Investigations, book 5, chap. Al, 160 p.
- Buchanan, T.J., and Somers, W.P., 1969, Discharge measurements at gaging stations: U.S. Geological Survey Techniques of Water-Resources Investigations, book 3, chap. A8, 65 p.
- Colton, R.B., 1969, Surficial geologic map of the New Preston quadrangle, Litchfield County, Connecticut: U.S. Geological Survey Quadrangle Map GQ-782, scale 1:24,000.
- Gates, R.M., and Bradley, W.C., 1952, The geology of the New Preston quadrangle: Connecticut Geological and Natural History Survey Miscellaneous Series no. 5, 46 p.
- Geldreich, E.E., and Kenner, B.A., 1969, Concepts of fecal streptococci in stream pollution: Journal Water Pollution Control Federation, v. 41, p. R336-R352.
- Greeson, P.E., Ehlke, T.A., Irwin, G.A., Lium, B.W., and Slack, K.V., eds., 1977, Methods for collection and analysis of aquatic biological and microbiological samples: U.S. Geological Survey Techniques of Water-Resources Investigations, book 5, chap. A4, 332 p.
- King's Mark Environmental Review Team, 1976, Eutrophication of Lake Waramaug, Kent, Warren, and Washington, Connecticut: King's Mark Resource Conservation and Development Area, Warren, Connecticut, 34 p.
- Kulp, K.P., 1981, Bathymetry and aquatic plants of Lake Waramaug, Connecticut: U.S. Geological Survey Open-File Report 81-477, 1 sheet.
- Lee, D.R., 1977, A device for measuring seepage flux in lakes and estuaries: Limnology and Oceanography, v. 22, no. 1, p. 140-177.
- Norvell, W.A., and Frink, C.R., 1975, Water chemistry and fertility of twenty-three Connecticut lakes: Connecticut Agricultural Experiment Station Bulletin 759, 45 p.

SUPPLEMENTAL TABLES

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows

[ft 3 /s, cubic feet per second, μ S/cm, microsiemens per centimeter; $^\circ$ C, degrees Celsius; mg/L, milligrams per liter; immed., immediate; μ M, micrometer; MF, membrane filter; cols./100 mL, colonies per 100 milliliters; t/day, tons per day; <, less than; --, no data; e (following a number), estimated; B (following a number), non-ideal colony count]

						INFLOW	SITE II					
		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	Coli- form, fecal,	Strep- tococci	Solids, residue	
		flow,	con-	pН		0xygen,	(per-	total,	0.7	fecal	at 180°C	Solid
		instan- taneous	duct- ance	(stand- ard	ature	dis- solved	cent satur-	immed. (cols./	µM-MF (cols./	KF agar (cols./	dis- solved	dis- solved
Date	Time	(ft ³ /s)	(µS/cm)	units)	(°C)	(mg/L)	ation)	100 mL)	100 mL)	100 mL)	(mg/L)	(t/day)
March 1977												
14	1125	9.8	39	5.0	5.0	12.4	97	240	2	16	27	0.71
	1620	8.5	33	5.2	5.0	11.6	91	<8		35		.53
14 June 1977								<8	<1		23	
01 September 1	1545 977	.46	48	6.2	12.0	9.5	88		4,800	640	53	.07
20	1430	.60	62	6.4	15.0			12,000	900	2,800		
October 197												
20 November 19	1430 177	3.0	35	5.7	10.5				6	29		
08 March 1978	1510	4.1	35	5.5	10.0				220	540		
27	1305	16	33	6.8	2.0							
Date	Nitro- gen, NO2+NO3 total (mg/L as N)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)	Phos- phorus, dis- solved (mg/L as P)	Phos- phorus, ortho, total (mg/L as P)	Carbon, organic total (mg/L as C)	Algal growth poten- tial, bottle test (mg/L)	Sedi- ment, sus- pended (mg/L)	Sedi- ment, dis- charge, sus- pended (t/day)
March 1977												
14	0.09	0.030	0.52	0.64	2.8	0.020		0.010	3.5	0.2	5	0.13
14	.08	.020	.53	.63	2.8	.020		<.010	3.5	.1	4	.09
June 1977		.020		.55	5	.020			5,5	•••	•	.03
01	.16	.010	.22	.39	1.7	.030		<.010	5.8	3.1	18	.02
September 1		.010		.55					0.0		-0	.02
20	.08	.050	.10	.23	1.0	.010	0.010		7.6	1.8	8	.01
October 197		- 010	67	0.7	21	010	010		6.0		•	.02
20	<.10	<.010	.07	.07	. 31	.010	.010		6.0		2	.02
November 19		. 010	24	2.4	, ,	020	0.00					^.
08 March 1978	<.10	<.010	.24	.24	1.1	.030	.020		11	4.1	4	.04
27						.048	.006			.3	57	2.5

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

						INFLOW	SITE 12				
		Stream- flow, instan-	Spe- cific con- duct-		Temper-	0×ygen, dis-	Oxygen, dis- solved (per- cent	Coli- form total, immed.	Coli- form, fecal, 0.7	Strep- tococci fecal KF agar	
D-+-	T.2	taneous (ft³/s)	ance	ard	ature (°C)	solved	satur-	(cols./	(cols./	(cols./	
Date	Time	(ft°/s)	(µS/cm)	units)	()	(mg/L)	at ion)	100 mL)	100 mL)	100 mL)	
larch 1977											
14 14	1155 1555	196 171	75 72	6.6 6.B	4.0 4.0	13.2 13.2	100 100	2,100 7,400	350 4,700	18,000B 15,000B	
June 1977	1333	1/1	<i>,</i> -	0,0	4.0	13.2	100	7,400	4,700	15,0000	
01	1405	2.8	130	6.9	15.0	9.9	97	4,500	4,000	2,800	
02	1015	7.6	120	7.1	15.0	10.4	102	9,500	1,400	5,200	
07	1030	5.1	127	7.4	12.5	10.3	96	3,800	1,900	5,000	
September 197		6.0	106		16.0	0.6	0.6	16 000	100 0000	1 400 0000	
20	1115	6.9 5.2	195 195	7.7	16.0	9.5	96	16,000	120,0008	1,400,000B	
20 October 1977	1325	3.2	130		16.0						
20	1345	53	95	7.2	9.5	11.7	102		3,900	5,200	
lovember 1977			,,	/	7.5	,	102	- -	5,500	5,200	
08	1050	102	102	7.2	8.5	12.6	108		55,000	200,000	
08	1330	90	100		8.5						
08	1455	80	85		9.0						
larch 1978											
20	0915	16									
21	0918	16 36									
22 23	1015 09 06	36 32									
24	1345	45									
25	1400	35									
26	1220	30									
27	1005	245									
27	1100	325									
27	1200	325	7 7	6.9	1.0						
27	1230	325									
27	1300	325				••					
27	1330	325									
27 27	1400 1430	325 290									
27	1500	268									
27	1530	268									
27	1600	245									
27	1630	215									
28	1245	116	58	6.6	3.5						
29	1205	117									
30	1330	78									
31	1130	64	52	6.8	3.5						
		Solids, residue		Nitro- gen,	Nitro- gen, ammonia	Nitro- gen,					
			C-114	-		•					
		at 180°C		N02+N0		organio	-				
		dis- solved	dis- solved	total (mg/L	solved (mg/L	total (mg/L					
Date	Time	(mg/L)	(t/day)	as N)	as N)	as N)					
March 1977		<u> </u>									
14	1155	43	23	0.41	0.130	0.62					
14 June 1977	1555	41	19	.40	.140	.64					
01	1405	93	.70	.59	.030	.35					
02	1015	78	1.6	.42	.040	.41					
07 Cantember 10	1030	81	1.1	.51	.040	.41					
September 193 20	1115			.61	.070	.84					
20	1325			.61	.380	1.0					
October 1977	1525			.01	,500	1.0					
	1345			.20	< .010	.22					
20											
20 November 1971	7										
	7 1050			.38	.130	1.7					
lovember 1971				.38 .29 .29	.130 .010 .010	1.7 .79 .57					

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

						INFLUM	SITE 13					
		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	Coli- form, fecal,		Solids, residue	
		flow,	con-	pН		Oxygen,	(per-	total,	0.7	fecal	at 180°C	Solids
		instan- taneous	duct - ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-MF (cols./	KF agar (cols./	dis- solved	dis- solved
Date	Time	(ft ³ /s)	(µS/cm)	units)	(°C)	(mg/L)	at ion)	100 mL)	100 mL)	100 mL)	(mg/L)	(t/day
March 1977												
14	1235	7.4	85	6.9	6.0	12.3	98	5,200	3,300	24,000	60	1.2
14	1640	8.0	97	6.8	5.0	12.3	95	12,000	4,700	40,000	46	
June 1977	1040	0.0	<i>J</i> ,	0.0	3.0	12.2	33	12,000	7,700	70,000	40	
01	1455	.28	117	6.7	14.0	8.6	83	16,000	2,400	9,000	94	.0
02	1100	.20	113	6.8	15.0	9.2	90	13,000	10,000	4,800	74	.0
07	1105	.23	117	6.9	13.5	9.8	93	5,600	4,200	2,600	76	
eptember 1		.23	11/	0.5	15.5	7.0	,,,	3,000	7,200	2,000	/0	•••
20	1210	.05e	240		15.0	8.9	87	160,000	180,000	850,000	в	_
20	1410	.05	240		15.0			100,000				_
ctober 197		.03	2.40		15.0							-
20	1410	2.0	115	7.0	10.5	10.4	93		2,600	2,700		
lovember 19		2.0	113	,.0	10.5	10.7	33		2,000	2,700		
08	1215	6.2	111	6.8	9.5	10.6	92		150,000	590,000		
08	1445	4.0	112		11.0	10.0			130,000			
March 1978	1775	4.0	112		11.0							
27	1345	16	77	6.7	3.0							_
		Nitro-								Algal		Sedi-
	Nitro-	gen,	Nitro-				Phos-	Phos-		growth		ment,
	gen,	ammonia	gen,	Nitro-	Nitro-	Phos-	phorus,	phorus,	Carbon,	poten-	Sedi-	dis-
	NO2+NO3	dis-	organic	gen,	gen,	phorus,	dis-	ortho.	organic	tial,	ment,	charge,
	total	solved	total	total	total	total	solved	total	total	bottle	sus-	sus-
	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	test	pended	pended
Date	as N)	as N)	as N)	as N)	as NO3)	as P)	as P)	as P)	as C)	(mg/L)	(mq/L)	(t/day)
	as 117	43 11)	<u>as 117</u>	as 11/	as nos)	as r)	as rj	as rj	<u>us c)</u>	(mg/L)	(mg/ L)	(c/uuy/
March 1977		0.050	0.50		3.0					0.4	50	
14	0.81	0.250	0.58	1.6	7.3	0.130		0.050	6.5	24	59	1.2
14	. 85	. 340	. 52	1.7	7.6	. 200		.090	6.2	48	84	1.8
June 1977		•••	٠.			• • •				2.6		٠.
01	1.3	.130	. 84	2.3	10	.110		.040	4.7	3.8	54	.04
02	.92	.040	. 26	1.2	5.4	.030		.020	5.9	12	17	.00
07	.82	.040	. 21	1.1	4.7	.050		.010	7.0	12		
September 1		202		• •		05-				30	_	
20	2.4	.020	1.4	3.8	17	. 350	.250		9.2	79	8	
20	2.2	.070	. 77	3.0	13	. 240	.200		8.7		5	.00
ktober 197												
20	.92	.010	.22	1.2	5.1	. 130	.120		7.3	33	3	.02
lovember 19			<u>.</u> .									
	. 96	. 100	. 84	1.9	8.4	. 280	.160		9.8	63	47	.79
08												
08	. 85	.080	. 35	1.3	5.7	. 140	.160		12		25	.27
	. 85	.080	. 35	1.3	5.7	. 140	.160		12	40	25 576	.27 25

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

						INFLOW S	ITE 14				
		Stream-	Spe- cific			Coli- form,	Coli- form, fecal,		o- Solids		Nitro- gen,
		flow,	con-	рH		total		fec a1		_	ds, NO2+NO3
		instan-		(stand-	- Temper				-		
		taneous		ard	ature	(cols.		_			
Date	Time	(ft ³ /s)	(µS/cm) units)	(°C)	100 mL) 100 ml	.) 100	mL) (mg	/L) (t	/day) as N)
March 1977											
14	1300	0.13e	210	7.2	6.0	5,000	3,600	950	111	0.04	2.0
ctober 1977 20	1425	.05	310	7.6	11.0		380	310			1.1
lovember 197 08	7 1310	5.1	320		10.5		1,800	5,400			1.0
March 1978							1,000	3,400			1.0
27	1435	.09	182	6.8	4.0						
	Nitro-								Algal		Sed i -
	gen, ammonia	Nitro- gen,	Nitro-	Nitro-	Phos-	Phos- phorus,	Phos- phorus, (Carbon,	growth poten-	Sedi-	ment, dis-
	dis-	organic	gen,	gen, p	phorus,	dis-	ortho, d	organic	tial,	ment,	charge,
	solved		total	total	total	so lved	total	total	bottle	sus-	Sus-
Date	(mg/L as N)	(mg/L as N)	(mg/L as N) a	(mg/L as NO3)	(mg/L as P)	(mg/L as P)	(mg/L as P) a	(mg/L as C)	test (mg/L)	pended (mg/L)	pended (t/day)
March 1977											
14	0.150	1.1	3.2	14	0.070		0.040	5.8	42	40	
October 1977 20	.660	.44	2.2	9.7	.600	.570		7.1		7	.00
November 197	7								110		
08 March 1978	1.60	1.1	3.7	16	.980	.980		12	118	24	. 33
27					.204	.167			52	35	.00
						INFLOW S	1TE 15				
							Coli-				
		Stream-	Spe- cific			Coli- form,	form, fecal		p- Solids ci residu		Nitro- gen,
		flow,	con-	рH		total	, 0.7	feca	1, at 180	°C Soli	ds, NO2+NO3
		instan- taneous		(stand- ard	ature	(cols.	•				
Date	Time	(ft ³ /s)	(µS/cn	n) units)	(°C)	100 mL) 100 mi	L) 100	mL) (mg	<u>/L) (t</u>	/day) as N)
March 1977	1215	0.00	• • •		0.0	445		170			
14 October 1977	1315	0.26e	104	7.1	8.0	44B	1	170	59	0.04	1.1
20	1420	.05	150	7.3	10.5		280	330			1.8
November 197 08	1400	2.7	1 40		11.0		780	1,500			.94
March 1978 27	1500	.18	83	6.8	4.0						**
						***			Alosi		Sodi
	Nitro- gen,	Nitro-				Phos-	Phos-		Algal growth		Sedi- ment,
	ammonia		Nitro-	Nitro-		p horus ,		Carbon,	pot en-	Sedi-	dis-
	dis- solved	organic total	gen, total	gen, p	phorus, total	dis- solved	ortho, o	organic total	tial, bottle	ment, sus-	charge, sus-
	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	test	pended	pended
Date	as N)			as NO3)		as P)		as C)	(mg/L)	(mg/L)	(t/day)
March 1977											
14 October 1977	0.050	0.70	1.9	8.2	0.020		0.010	7.4	7.3	35	
20	.010	.41	2.2	9.8	.020	.010		7.6	1.8	30	.00
November 197 08	7 <.010	.48	1.4	6.3	.030	.020		8.9	14	4	.03
March 1978											
27					.044	.016			12	70	.03

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

							0		Coli-			
		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	form, fecal,		Solids, residue	
		flow,	con-	pН		Oxygen,	(per-	total,	0.7	fecal	at 180°C	Solids
		instan- taneous	duct - ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-mF (cols./	KF agar (cols./	dis- solved	dis- solved
Date	Time	(ft ³ /s)	(US/cm)	units)	(°C)	(mg/L)	at ion)	100 mL)	100 mL)	100 mL)	(mg/L)	(t/day)
March 1977												
14	1325	11	62	6.5	6.0	12.2	98	220	20	3,300	39	1.2
14 June 1977	1705	8.6	57	6.7	5.0	12.6	98	67B	<1	15,000E	41	.95
01 September 1	1425 977	.55	97	6.8	15.0	9.4	92	130,000B	12,000B	35,000	74	.11
20 October 197	1425	.08	140		17.0			4,800	610	900E	3	
20 November 19	1325	.75	85	6.5	11.0				200	800		
08	1030	.68	76	6.8	9.0				9,200	27,000		
March 1978 27	1520	15	47	6.8	3.0							
	Nitro-	Nitro- gen,	Nitro-				Phos-	Phos-		Algal growth		Sedi- ment,
	qen,	ammonia	gen,	Nitro-	Nitro-	Phos-	phorus.	phorus,	Carbon.	pot en-	Sedi-	dis-
	N02+N03	dis-	organic	gen,	gen,	phorus,	dis-	ortho,	organic	tial,	ment,	charge,
	total	solved	total	total	total	total	solved	total	total	bottle	sus-	sus-
	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	test	pended	pended
Date	às N)	as N)	as N)	as N)	as NO3)	as P)	as P)	às P)	as C)	(mg/L)	(mg/L)	(t/day)
March 1977												
14	0.34	0.050	0.55	0.94	4.2	0.D30		<0.010	9.1	1.4	53	1.6
14	. 35	.050	. 58	.98	4.3	.030		<.010	6.6	1.0	54	1.3
June 1977												
01	. 21	.040	1.4	1.6	7.1	.120		<.010	6.1	1.0	184	.27
September :		01.0				050	010		0.5		•	00
20	.23	.010	. 47	.71	3.1	.050	.010		9.5	1.2	8	.00
October 19: 20	.25	<.010	. 29	.54	2.4	.010	.010		9.2	.6	5	.01
20 November 19		·.UIU	. 29	.54	2.4	.010	.010		9.2	.0	9	.01
08	.22	.010	.99	1.2	5.4	.170	.030		14	1.0	115	.21
March 1978												

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

						INFLOW	\$1TE 17	
		Stream- flow, instan- taneous	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Oxygen, dis- solved	Oxygen, dis- solved (per- cent satur-	Coli- form total, immed. (cols./
Date	Time	(ft ³ /s)	(µS/cm)	units)	(°C)	(mg/L)	ation)	100 mL)
					•			
March 1977 14 June 1977	1405	80e	63	6.4	3.0	13.5	100	1,500
01	1710	6.7	130	7.1	15.0	10.0	98	3,700
02	1205	2.5	118	7.1	17.0	9.9	102	4,700
07	1155	3.0	125	7.1	13.0	10.0	94	5 40
September 19		_						
20	1530	.82	245		14.5			1,800B
October 197		20	7.5					
20	1525	20	75	6.6	9.5			
November 19:	1555	24	75		0.0			
08 March 1978	1 22 2	34	/5		9.0			
20	0855	6.7						
21	0900	6.1						
22	0955	26						
23	0850	25						
24	1415	28						
25	0900	22						
26	1045	17						
27	0940	52						
27	1100	64						
27	1230	64	94	6.7	.0			
27	1320	65						
27	1350	65						
27	1430	6 9						
27	1500	71						
27	1600	74						
27	1620	74						
28	1100	52	50	6.9	2.0			
29	1120	42						
30	1400	32						
31	1225	25	50	6.4	3.5			

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

					<u>1N</u>	FLOW SITE I	7Continu
	Coli-					Nitro-	
	form, fecal,	Strep- tococci,	Solids, residue		Nitro- gen,	gen, ammonia	Nitro- gen,
	0.7	fecal,	at 180°C	Solids,	N02+N03	dis-	organic
Date	μM-MF (cols./ 100 mL)	KF agar (cols./ 1DO mL)	dis- solved (mg/L)	dis- solved (t/day)	total (mg/L as N)	solved (mg/L as N)	total (mg/L as N)
March 1977							
14 June 1977	45	420	44	9.5	0.32	0.070	0.63
01	1,900B	4,300	92	1.7	. 45	. 040	1.2
02	160	540	80	.54	. 22	.020	. 43
07	210	520	91	.74	. 25	.010	. 31
September							
20	1,000	2,600			. 21	.010	. 21
October 19							
20	65	140			. 03	<.010	. 16
November 1							
08	3,100	2,400			. 09	<.010	. 38
March 1978							
20							
21							
22 23							
24							
25							
26							
27							
27							
27							
27							
27							
27							
27							
27							
27							
28							
29					-		
30							
31							

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

						INFLOW	SITE 18					
		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	Coli- form, fecal,		Solids, residue	**************************************
		flow,	con-	pН		Oxygen,	(per-	total,	0.7	fecal	at 180°C	Solids
		instan- taneous	duct- ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-MF (cols./	KF agar (cols./	dis- solved	dis- solved
Date	Time	(ft ³ /s)	(µS/cm)	units)	(°C)	(mg/L)	ation)	100 mL)	100 mL)	100 mL)	(mg/L)	(t/day
larch 1977												
14 lune 1977	1135	2.2	62	8.0	5.5	12.0	95	1108	1	74	43	0.26
01 October 197	1555 7	.20e	116	6.6	15.0	8.6	84	4,700	6308	4,9008	85	.05
20 November 19		.10	108	6.3	11.0				84	270		
08 March 1978	1145	.26	80	6.3	9.5				3,200	11,000		
27	1600	.35	47	7.1	3.0							
		Nitro-								Algal		Sedi-
	Nitro-	gen,	Nitro-			ъ.	Phos-	Phos-		growth		ment,
	gen, NO2+N O 3	ammonia dis-	gen, organic	Nitro- gen.	Nitro-	Phos- phorus,	phorus, dis-	phorus, ortho.	Carbon, organic	poten- tial,	Sedi- ment,	dis- charge.
	total	solved	total	total	gen, total	total	solved	total	tota1	bottle	sus-	sus-
	(mg/L	(mq/L	(mq/L	(mq/L	(mg/L	(mq/L	(mg/L	(mq/L	(mg/L	test	pended	pended
Date	às N)	às N)	as N)	as N)	as NO3)	as P)	as P)	as P)	as C)	(mg/L)	(mg/L)	(t/day)
March 1977	0.12	0.030	0.50	0.65	2.9	0.030		0.010	4.8	5.5	65	0.39
June 1977	U. 12	0.030	0.30	0.05	2.9	0.030		0.010	4.0	3,5	υɔ	0.39
01 October 197	.30	.020	. 50	.82	3.6	.110		.040	5.4	13	66	
20 lovember 19	.60	.010	. 20	.81	3.6	.020	.020		7.1	7.0	6	.00
	.13	<.010	. 17	.30	1.3	.080	.050		10	14	44	.03
08 March 1978	.13	1.010	• • •							- '		

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

						INFLOW	SITE 19					
		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	Coli- form, fecal,		Solids, residue	
		flow,	con-	pН		0×ygen,	(per-	total,	0.7	fecal	at 180°C	Solids
		instan- taneous	duct- ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-MF (cols./	KF agar (cols./	dis- solved	dis- solved
Date	Time	(ft ³ /s)	(µS/cm)	units)	(°C)	(mg/L)	at ion)	1DO mL)	10C mL)	100 mL)	(mg/L)	(t/day)
March 1977												
14	1210	46	36	6.7	4.0	13.1	99	100B	<1	35	30	3.7
14	1745	40e	34	6.5	3.0	13.5	102	17B	<1	37	30	3.2
June 1977	2/43	406	54	0.5	5.0	15.5	102	• • • •	-1	3,	30	5.2
01	1640	.64	48	6.4	14.0	9.7	93	4,200	2,100E	2,4008	3 44	.08
02	1030	.84	43	6.7	16.0	10.0	101	4,200	<1	300	43	.10
07	1030	.66	45	6.5	13.0	10.4	98	1,500	100	720	42	.07
September 19		.00	-43	0.5	-5.0	10.7	70	2,500	100	, 20	-16	.07
20	1340	.05	50		15.0			820	150	1,500	R	
October 1977	1040	.05	30		13.0			OLU	130	1,5001		
20	1250	9.3	52	6.5	10.0				42	100		
November 197		3.3	32	0.5	10.0				42	100		
08	1230	4.7	57		10.0				640	2,200		
March 1978	1230	7./	3/		10.0				040	2,200		
20	0926	2.9										
21	0920	3.2										
22	1025	3.2 7.8										
23	0915	9.2										
	1350	14										
24 25	1410	12										
25 26	1215	7.4										
	1015	45										
27		45 38	28	6.3	4.5							
27	1115											
27	1145	37 38	30 30	6.4	4.5 4.5							
27	1215	38 38	30 28	6.3	4.5							
27	1245 1315	38 37	28 27	6.4 6.4	4.0							
27 27	1315	3/ 36	2/ 28	6.4	4.0							
27	1415	36 34	28 27	6.5	4.0							
			27 26	6.5								
27	1445	34			4.0							
27	1515	34 34	28 29	6.4	4.0							
27	1545			6.4	4.0 3.5							
27	1615	34	27	6.4								
27	1645	32	30	6.5	3.5							
28	1215	26	29	6.7	3.5							
29	1215	21										
30	1345	18										
31	1200	15	24	6.7	4.0							

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

					<u>I N</u>	FLOW SITE	19Cont	inued				
	Nitro- gen, NO2+NO3 total (mg/L	Nitro- gen, ammonia dis- solved (mg/L	Nitro- gen, organic total (mg/L	Nitro- gen, total (mg/L	Nitro- gen, total (mg/L	Phos- phorus, total (mg/L	Phos- phorus, dis- solved (mg/L	Phos- phorus, ortho, total (mg/L	Carbon, organic total (mg/L	Algal growth poten- tial, bottle test	Sedi- ment, sus- pended	Sedi- ment, dis- charge, sus- pended
Date	as N)	as N)	as N)	as N)	as NO3)	as P)	as P)	as P)	as C)	(mg/L)	(mg/L)	(t/day)
March 1977	,											
14	0.12	0.060	0.47	0.65	2.9	0.030		0.010	5.9	5.1	31	3.9
14	.14	.050	.65	.84	3.7	.030		.010	7.9	2.3	8	0.5
June 1977	. 14	.030	. 00	.54	3.7	.030		.010	,.,	2.5	·	
01	. 50	.030	.59	1.1	5.0	.060		.020	3,2	16	33	.0€
02	.26	.010	.34	.61	2.7	.060		.020	5.2	11	28	.06
07	.36	<.010	. 49	.85	3.8	.050		.020	6.3	10	11	.02
		\.U10	. 49	.03	3.0	.030		.020	0.3	10	11	.02
September		01.0	.17	A.C	2.0	.030	.020		8.5	9.6	8	.00
20	. 28	.010	.17	.46	2.0	.030	.020		0.5	9.0	0	.00
October 19		000	٠.		4 2	000	000		10		,,	0.0
20	.02	.020	. 91	.95	4.2	. 020	.020		10	1.7	11	.28
November 1		***			• •							^-
08	.06	.010	. 39	.46	2.0	.060	.020		9.1	2.2	16	.20
March 1978	3											
20			~-			.012						
21						.004						
22						.011						
23						.016						
24						.017						
25						.013						
26						.010						
27			~-			.170	<.010					
27						.110	.005					
27						.066	.004					
27						.035	.007			.9	80	8.3
27						.051	.004				51	5.2
27						.042	.004				52	5.2
27						.034	.004				40	3.9
27						.028	.004				38	3.5
27						.028	,007				31	2.8
27						.048	.006				33	3.0
27						.028	.009				25	2.3
27						.020	.009				23	2.1
27						.047	.005				26	2.2
						.048	.005				20 8	.50
28											3	.1
29						.015						
30						. 024					11	.53
31				~-		.017	.008	~~			2	.08

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

01 07	Time 1250 1730 1720	Stream- flow, instan- taneous (ft ³ /s) 0.44 .44e	Spe- cific con- duct- ance (µS/cm)	pH (stand- ard units) 7.0	Temper- ature (°C)	Oxygen, dis- solved (mg/L)	Oxygen, dis- solved (per- cent satur- ation)	Coli- form total, immed. (cols./	Coli- form, fecal, 0.7 µM-MF (cols./	Strep- tococci fecal KF agar (cols./	Solids, residue at 180°C dis- solved (mg/L)	Solids dis- solved
March 1977 14 14 June 1977 01 07	1250 1730 1720	instan- taneous (ft ³ /s)	duct- ance (µS/cm)	(stand- ard units)	ature	dis- solved	cent satur-	immed. (cols./	μM-MF (cols./	KF agar (cols./	dis- solved	dis- solved
March 1977 14 14 June 1977 01 07	1250 1730 1720	taneous (ft^3/s)	ance (µS/cm) 72	ard units)	ature	solved	satur-	(cols./	(cols./	(cols./	so I ved	solved
March 1977 14 14 June 1977 01 07	1250 1730 1720	0.44	72		(°C)	(mg/L)	at ion)	100 ml \	100 -11	100 -11	(mo (1)	
14 14 June 1977 01 07	1730 1720			7.0				AUU IIIL)	TOO MIL)	TOO IDT)	(mg/ r)	(t/day
14 14 June 1977 01 07	1730 1720			7 A								
14 June 1977 01 07	1730 1720				5.0	10.8	84	42B	1	110	50	0.06
June 1977 01	1720	.446		7.0	5.0	12.6	98	<8	<1	100	47	.06
01 07			07	7.0	5.0	12.0	70	`0	~1	100	47	.00
07		1.5e	63	6.7	14.0	9.4	90	5,200	4,200B	1,900	53	.21
			56									
	1130	.14	20	6.9	14.0	10.2	98	1,400	600	1,900	46	.02
		1.0	0.0		10.0				100			
20	1230	.10	96	6.4	10.0				100	62		
November 1							_					
08	1315	.28	80		11.0	10.4	94		520	1,500		
March 1978												
27	1245	1.3	49	6.5	6.0							
Date	Nitro- gen, NO2+NO3 total (mg/L as N)	Nitro- gen, ammonia dis solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)	Phos- phorus, dis- solved (mg/L as P)	Phos- phorus, ortho, total (mg/L as P)	Carbon, organic total (mg/L as C)	Algal growth poten- tial, bottle test (mg/L)	Sedi- ment, sus- pended (mg/L)	Sedi- ment, dis- charge, sus- pended (t/day)
March 1977												
14	0.18	0.050	0.83	1.1	4.7	0.020		< 0.010	5.6	0.7	14	0.02
14	.16	.050	. 53	.74	3.3	.020		<.010	4.4	.6	7	
June 1977			. 30	•••					••		•	
01	. 19	.010	1.8	2.0	8.8	.120		.010	7.2	. 4	119	
07	.04	.010	.05	.10	. 44	.010		<.010	5.7	1.1	0	.00
October 19											•	
20	. 12	<.010	, 27	.39	1.7	.010	.010		9.3	3.2	12	.00
November 1			• /			.010			٠.٠	٧. ٤		
08	.01	<.010	. 31	.32	1.4	.020	.010		9.2	3.2	10	.00
March 1978		0		.52	•••	.0.0	.010		٥	٠.٢	10	.00
27						.013	<.010			.7	21	.07

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

								-				
		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	Coli- form, fecal,		Solids, residue	
		flow,	con-	pН		Oxygen,	(per-	total,	0.7	fecal	at 180°C	Solids
		instan- taneous	duct- ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	μM-MF (cols./	KF agar (cols./	dis- solved	dis- solved
Date	Time	(ft ³ /s)	(µS/cm)	units)	(°C)	(mg/L)	at ion)	100 mL)	100 mL)	100 mL)	(mg/L)	(t/day
March 1977												
14	1310	1.9	88	7.1	5.0	10.2	80	520B	10	500	63	0.32
14	1720	1.5e	90	6.8	3.5	11.0	82	360B	24		59	. 24
June 1977	1.20	1.50	,,,	0.0	3.3	11.0	υZ	3300	£ 71	. 500	33	7
01	1810	1.0e	104	6.9	14.0	9.6	92	25,000B	24,000	3 20,000E	75	. 20
02	1130	.10	125	7.3	16.0	10.2	103	2,700	<0	2,100	83	.02
07	1205	.12	120	7.4	13.0	10.4	98	4,800B	1,300	2,500	82	.03
September :		•	120			20.1	,,	.,	-,000	-,000		
20	1230	.04e	190		15.5	10.0	99	9,500B	780	5,500		
October 19		.540	130		-0.0	-0.0	,,,	-,000	.00	-,		
20	1205	.30e	146	6.8	10.0				65	270		
November 19		.500	170	0.0	10.0	J.			03	-70	-	
08	1350	.99	120		11.0	10.2	92		6,100	5,200		
March 1978	1330		120		11.0	10.2	32		0,100	3,200		
27	1310	3.5	51	6.4	5.0							
		Nitro-								Algal		Sedi-
	Nitro-	gen,	Nitro-				Phos-	Phos-		growth		ment,
	gen,	ammonia	gen,	Nitro-	Nitro-	Phos-	phorus,	phorus,	Carbon,	pot en -	Sedi-	dis-
	N02+N03	dis	organic	gen,	gen,	phorus,	dis-	ortho,	organic	tial,	ment,	charge,
	total	solved	total	total	total	total	solved	total	total	bottle	sus-	sus-
	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	test	pended	pended
Date	as N)	as N)	as N)	as N)	as NO3)	as P)	as P)	as P)	as C)	(mg/L)	(mg/L)	(t/day)
March 1977				•					•			
14	0.61	0.150	0.95	1.7	7.6	0.080		0.020	12	17	40	D.21
14	.58	.140	.96	1.7	7.4	.080		.020	15	21	32	0,21
June 1977	. 50	, 140	. 70	1.7	7.4	.000		.020	13	£ 1	32	
01	. 37	.020	2.4	2.8	12	. 300		.030	9.6	13	218	
02	.11	.020	.43	.56	2.5	.060		.060	4.2	6.7	17	.00
	.09	.010	. 43	.34	1.5	.060		.040	5.3	7.2	6	.00
07 Sept em ber 1		.010	. 24	.34	1,5	.000		.040	5,3	1.2	0	.00
	<.10	010	Λ0	10		020	010		8.5	23	12	
20 October 19		.010	. 09	.10	. 44	.020	.010		8.5	23	12	
		- 010	40	0.0	2.0	020	0.20		0.5	16	2	
20	.37	<.010	. 49	.86	3.8	.030	.030		8.5	16	3	
November 1	-	010	۲۵		٠.	070	0.00		10		10	
08	. 48	.010	. 68	1.2	5.2	.070	.060		12	9.7	18	.05
08 March 1978 27	. 40	.010	.08	1.2	5,2	.165	.010		12	3.5	91	.86

Table 7--Chemical, physical, and biological characteristics of surface-water inflows--Continued

						INFLO	SITE 112	<u>!</u>			
		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	Coli- form, fecal,	Strep- tococci	Solids, residue
		flow,	con-	pН		Oxygen,	(per-	total,	0.7	fecal	at 180°C
		inst a n- taneous	duct - ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-mf (cols./	KF agar (cols./	dis- solved
<u>Date</u>	Time	(ft ³ /s)	(µS/cm)	units)	(°C)	(mg/L)	ation)	100 mL)	100 mL)	100 mL)	(mg/L)
March 1977 14	1340	0.30e	114	7.4	5.0	10.0	78	<8	3	260	71
(olids, dis- olved	Nitro- gen, NO2+NO3 total (mg/L	Nitro- gen, ammonia dis- solved (mg/L	Nitro- gen, organic total (mg/L	Nitro- gen, total (mg/L	Nitro- gen, total (mg/L	Phos- phorus, total (mg/L	Phos- phorus, ortho, total (mg/L	Carbon, organic total (mg/L	Algal growth poten- tial, bottle test	Sedi- ment, sus- pended
Date (as N)	as N)	as N)	as N)	as NO3)	as P)	as P)	as C)	(mg/L)	(mg/L)
March 1977 14		0.30	0.050	0.73	1.1	4.8	0.020	<0.010	6.9	0.7	3
						INFLO	W SITE 113	<u>1</u>			
		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	Coli- form, fecal,	Strep- tococci	Solids, residue
		flow,	con-	pН		0xygen,	(per-	total,	0.7	fecal	at 180°C
		instan- taneous	duct - ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-mf (cols./	KF agar (cols./	dis- solved
Date	Time	(ft ³ /s)	(µS/cm)	units)	(°C)	(mg/L)	at ion)	100 mL)	100 mL)	100 mL)	(mg/L)
March 1977 14 March 1978	1400		144	6.7	6.0	11.8	94	120B	1	41	86
27	1335	.13	96	6.3	6.0						
	Nitro- gen,	Nitro- gen, ammonia	Nitro- gen,	Nitro-	Nitro-	Phos-	Phos-	Phos-	Algal growth poten-	Sed i-	Sedi- ment, dis-
0	NO2+NO3 total (mg/L	dis- solved (mg/L	organic total (mg/L	gen, total (mg/L	gen, total (mg/L	phorus, total (mg/L	dis- solved (mg/L	ortho, total (mg/L	tial, bottle test	ment, sus- pended	charge, sus- pended
Date	as N)	as N)	as N)	as N)	as NO3)	as P)	as P)	as P)	(mg/L)	(mg/L)	(t/day)
M arch 1977	1.0	0.040	0.86	1.9	8.4	0.050		0.010	2,8	3	
14 March 1978		0.040	0.00								

 $\label{thm:continued} \textbf{Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued}$

							Ovvoer		Coli-			
		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	form, fecal,	Strep- tococci	Solids, residue	Nitro- gen,
		flow,	con-	рH		Oxygen,	(per-	total,	0.7	fecal	at 180°C	NO2+NO3
		instan- taneous	duct - ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-MF (cols./	KF agar (cols./	dis- solved	total (mg/L
Date	Time	(ft ³ /s)	(µS/cm)	units)	(°C)	(mg/L)	at ion)	100 mL)	100 mL)	100 mL)	(mg/L)	(as N)
March 1977	1620		100	6.6	4.5	7.8	58	12000	4000	280	56	0.18
June 1977 01 March 1978	1800		106	6.7	19.0	6.2	67	260	4	69	60	.02
27	1420	6.5	58	6.3	1.0							
	Nitro-				· · · · · · · · · · · · · · · · · · ·				Algal		Sedi-	
	gen, ammonia	Nitro-	Nitoo	Nitro-	Phos-	Phos-	Phos-	Cambon	growth	Sedi-	ment, dis-	
	dis-	gen, organic	Nitro- gen,	gen,	phorus,	phorus, dis-	phorus, ortho,	Carbon, organic	poten- tial,	ment,	charge,	
	so lved	total	total	total	total	solved	total	total	bott le	sus-	sus-	
Date	(mg/L as N)	(mg/L as N)	(mg/L as N)	(mg/L as NO3)	(mg/L as P)	(mg/L as P)	(mg/L as P)	(mg/L as C)	test (mg/L)	pended (mg/L)	pended (t/day)	
March 1977												
14	0.080	0.67	0.93	4.1	0.030		0.010	5.8	3.1	3		
June 1977 01	.120	.55	. 69	3.1	.060		.040	4.8	8.1	33		
March 1978						020			0.0	7	10	
27					.050	.038			8.9	,	.12	
						INFLOW	SITE ILE	<u> </u>				
						Oxygen,		Coli-				
		Spe- cific				dis- solved	Coli- form	form, fecal,	Strep- tococci	Solids, residue		
		con-	рΗ		Oxygen,	(per-	total,	0.7	fecal	at 180°(:	
		duct- ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-mF (cols./	KF agar (cols./	dis- solved		
Date	Time	(µS/cm)	units)	(°C)	(mg/L)	ation)	100 mL)	100 mL)	100 mL)	(mg/L)		
March 1977 14	1455	262	7.1	2,0	6.8	49	42B	2	7500	139		
	3	-31	, 	2.0	2.0	,,		-	. 530	.33		
		Nitro-							Algal	_		
	Nitro-	gen,	Nitro-	Nitma	Nitas	Dhee	Phos -	Cambon	growth	Sod:		
	gen, NO2+NO3	ammonia dis-	gen, organic	Nitro- gen,	Nitro- gen,	Phos- phorus,	phorus, ortho,	Carbon, organic	poten- tial,	Sedi- ment.		
	total	solved	total	total	total	total	total	total	bott le	sus-		
Date	(mg/L as N)	(mg/L as N)	(mg/L as N)	(mg/L as N)	(mg/L as NO3)	(mg/L as P)	(mg/L as P)	(mg/L as C)	test (mg/L)	pended (mg/L)		
March 1977												
14	0.43	0.080	0.85	1.4	6.0	0.090	0.020	5.5	0.8	26		

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	Coli- form, fecal,	Strep- tococci	Solids, residue	
		flow,	con-	рH		Oxygen,	(per-	total,	0.7	fecal	at 180° C	Solids
		instan- taneous	duct- ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-MF (cols./	KF agar (cols./	dis- solved	dis- solved
Date	Time	(ft ³ /s)	(US/cm)	units)	(°C)	(mg/L)	at ion)	100 mL)	100 mL)	100 mL)	(mg/L)	(t/day)
March 1977 14 March 1978	1700	0.10e	460	6.8	4.0	9.4	71	25B	2	31	227	0.06
27	1500	.01	99	6.3	7.0							
Date	Nitro- gen, NO2+NO3 total (mg/L as N)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)	Phos- phorus, dis- solved (mg/L as P)	Phos- phorus, ortho, total (mg/L as P)	Carbon, organic total (mg/L as C)	Algal growth poten- tial, bottle test (mg/L)	Sedi- ment, sus- pended (mg/L)	Sedi- ment, dis- charge, sus- pended (t/day)
March 1977 14 March 1978	0.12	0.060	0.94	1.1	5.0	0.050		0.010	8.6	1.4	48	
27						.039	.021			3.4	7	.00

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

							Oxygen,	-	Coli-			
		Stream-	Spe- cific				dis- solved	Coli- form	form, fecal,	Strep- tococci	Solids, residue	
		flow,	con-	pН		Oxygen,	(per-	total,	0.7	fecal	at 180°C	Solids
		instan- taneous	duct - ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	μM-MF (cols./	KF agar (cols./	dis- solved	dis- solved
Date	Time	(ft ³ /s)	(µS/cm)	units)	(°C)	(mg/L)	at ion)	100 mL)	100 mL)	100 mL)	(mg/L)	(t/day)
March 1977												
14	1130	4.1	53	6.8	3.5	13.0	96	<8	<1	57	28	0.31
14 June 1977	1600	4.0	57	6.8	3.0	13.4	99	<8	<1	67	33	.36
01	1430	.13	90	6.5	12.0	9.0	83	4,000	1,000	7,700B	58	.02
September 1	1977 121 5	0.6	101	6.8	15.0			12,000	000	1,800		
20 October 197		.06	101	0.8	13.0			12,000	880	1,000		
20	1500	3.0	54	6.6	11.0				17	40		
November 19 08	1600	.87	60	6.6	10.5				500	380		
March 1978									-	-3-		
27	1555	5.1	43	7.0	1.0	••				,		
		Nitro-	····							Algal		Sedi-
	Nitro-	gen,	Nitro-				Phos-	Phos-		growth		ment,
	gen, NO2+NO3	ammonia dis-	gen, organic	Nitro- gen,	Nitro- gen,	Phos- phorus,	phorus, dis-	phorus, ortho,	Carbon, organic	poten- tial,	Sedi- ment,	dis- charge,
	total	solved	total	total	total	total	solved	total	total	bottle	sus-	sus-
	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(m g/L	(mg/L	(mg/L	test	pended	pended
Date	as N)	as N)	as N)	as N)	as NO3)	as P)	as P)	as P)	as C)	(mg/L)	(mg/L)	(t/day)
March 1977												
14	0.28	0.050	0.55	0.88	3.9	0.020		<0.010	3.7	0.7	16	0.18
14	. 24	.020	. 24	.50	2.2	.030		<.010	4.4	4.4	13	.14
June 1977 01	. 15	.040	. 62	.81	3.6	.040		.010	4.1	1.0	37	.01
September 1		.040	. 02	.01	3.0	.040		.010	7.1	1.0	3/	.01
20	. 32	.010	.37	.70	3.1	.090	.080		7.3	2.6	5	.00
	77											
October 197		01.0	0.3	ΛE	22	01.0	010		E 3	A	2	0.2
October 197 20	.02	.010	.02	.05	. 22	.010	.010		5.3	. 4	3	.02
October 197	.02	.010 <.010	.02	.05 .21	.22	.010	.010 .020		5.3 11	.4 11	3 6	.02

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

						INFLOW	SITE I18					
		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	Coli- form, fecal,	Strep- tococci	Solids, residue	
		flow,	con-	рH		Oxygen,	(per-	total,	0.7	fecal	at 180°C	Solids
		instan- taneous	duct- ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	μM-MF (cols./	KF agar (cols./	dis- solved	dis- solved
Oate	Time	(ft ³ /s)	(US/cm)	units)	(°C)	(mg/L)	at ion)	100 mL)	100 mL)	100 mL)	(mg/L)	(t/day)
March 1977												
14 November 19	1215 977	0.25	57	6.9	3.0	13.6	101	50B	<4	340	34	0.02
08	1515	.05	68	6.1	10.5				220	2,600		
	Nitro-	Nitro- gen,	Nitro-				Phos-	Phos-		Algal growth		Sedi- ment,
	gen,	ammonia	gen,	Nitro-	Nitro-	Phos-	phorus,	phorus,	Carbon,	pot en-	Sedi-	dis-
	NO2+NO3	dis-	organic	gen,	gen,	phorus,	dis-	ortho,	organic	tial,	ment,	charge,
	total	solved	total	total	total	total	solved	total	total	bottle	sus- pended	SUS-
Date	(mg/L as N)	(mg/L as N)	(mg/L as N)	(mag/L as N)	(mg/L as NO3)	(mg/L as P)	(mg/L as P)	(mg/L as P)	(mg/L as C)	test (mg/L)	(mg/L)	pended (t/day)
						1	==					
March 1977 14 November 19	0.97	0.080	0.82	1.9	8.3	0.180		0.010	8.0	2.1	75	0.05
08	.95	<.010	. 49	1.4	6.4	.070	.070		9.4	48	2	.00
		Stream-	Spe- cific				dis- solved	Coli- form	form, fecal,	Strep- tococci	Solids, residue	
		flow,	con-	pН		Oxygen,	(per-	total,	0.7	fecal	at 180°C	Solids
		instan- taneous	duct - ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	μM-MF (cols./	KF agar (cols./	dis- solved	dis- solved
Date	Time	(ft ³ /s)	(µS/cm)	units)	(°C)	(mg/L)	ation)	100 mL)	100 mL)	100 mL)	(mg/L)	(t/day)
March 1977												
14 October 19	1240	0.45	55	7.0	3.0	13.6	101	25B	<1	120	30	0.04
20	1520	.10	81	6.6	11.0				11,000	1,400		
November 19 08	97 <i>7</i> 1 5 00	.15	70	6.7	11.0				21,000	9,800		
March 1978 27	1530	.18	47	7.0	2.0							
	Nitro	Nitro-	Nitro				Phos -	Phos-		Algal arowth	,	Sedi-
	Nitro-	gen,	Nitro-	Nitro-	Nitro-	Phos-	Phos-	Phos-	Carbon.	growth	Sedi-	ment,
	Nitro- gen, NO2+NO3	gen, ammonia	Nitro- gen, organic	Nitro- gen,	Nitro- gen,	Phos- phorus,	Phos- phorus, dis-		Carbon, organic		Sedi- ment,	ment, dis-
	gen, NO2+NO3 total	gen, ammonia dis- solved	gen, organic total	gen, total	gen, total	phorus, total	phorus, dis- solved	phorus, ortho, total	organic total	growth poten- tial, bottle	ment, sus-	ment, dis- charge, sus-
	gen, NO2+NO3 total (mg/L	gen, ammonia dis- solved (mg/L	gen, organic total (mg/L	gen, total (m g/L	gen, total (mg/L	phorus, total (mg/L	phorus, dis- solved (mg/L	phorus, ortho, total (mg/L	organic total (mg/L	growth poten- tial, bottle test	ment, sus- pended	ment, dis- charge, sus- pended
Date	gen, NO2+NO3 total	gen, ammonia dis- solved	gen, organic total	gen, total	gen, total	phorus, total	phorus, dis- solved	phorus, ortho, total	organic total	growth poten- tial, bottle	ment, sus-	ment, dis- charge, sus-
Date March 1977 14	gen, NO2+NO3 total (mg/L as N)	gen, ammonia dis- solved (mg/L	gen, organic total (mg/L	gen, total (m g/L	gen, total (mg/L	phorus, total (mg/L	phorus, dis- solved (mg/L	phorus, ortho, total (mg/L	organic total (mg/L	growth poten- tial, bottle test	ment, sus- pended	ment, dis- charge, sus- pended
Date March 1977 14 October 19 20	gen, NO2+NO3 total (mg/L as N) 0.63	gen, ammonia dis- solved (mg/L as N)	gen, organic total (mg/L as N)	gen, total (mg/L as N)	gen, total (mg/L as NO3)	phorus, total (mg/L as P)	phorus, dis- solved (mg/L as P)	phorus, ortho, total (mg/L as P)	organic total (mg/L as C)	growth poten- tial, bottle test (mg/L)	ment, sus- pended (mg/L)	ment, dis- charge, sus- pended (t/day)
<u>Date</u> March 1977 14 October 19	gen, NO2+NO3 total (mg/L as N) 0.63	gen, ammonia dis- solved (mg/L as N)	gen, organic total (mg/L as N)	gen, total (mg/L as N)	gen, total (mg/L as NO3)	phorus, total (mg/L as P)	phorus, dis- solved (mg/L as P)	phorus, ortho, total (mg/L as P)	organic total (mg/L as C)	growth poten- tial, bottle test (mg/L)	ment, sus- pended (mg/L)	ment, dis- charge, sus- pended (t/day)

Table 7.--Chemical, physical and biological characteristics of surface-inflows--Continued

						INFLOW	SITE 120	!				
		Stream-	Spe- cific		·		Oxygen, dis- solved	Coli- form	Coli- form, fecal,	Strep- tococci	Solids, residue	
		flow,	con-	рH		Oxygen,	(per-	total,	0.7	fecal	at 180°C	Solids
		instan- taneous	duct- ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-MF (cols./	KF agar (cols./	dis- solved	dis- solved
Date	Time	(ft ³ /s)	(µS/cm)	units)	(°C)	(mg/L)	at ion)	100 mL)	100 mL)	100 mL)	(mg/L)	(t/day)
March 1977 14 October 19	1255	0.78	185	7.0	3.0	13.2	98	1,600	<1	1,600	107	0.23
20 lovember 1	1525	.05	210	6.6	11.0				10	17		
08 March 1978		.08	170	6.8	10.5				230	480		
27	1510	.18	88	7.0	1.0							
	Nitro-	Nitro- gen,	Nitro-				Phos-	Phos-		Algal growth		Sedi- ment,
	gen, NO2+NO3 total	ammonia dis- solved	gen, organic total	Nitro- gen, total	Nitro- gen, total	Phos- phorus, total	phorus, dis- solved	phorus, ortho, total	Carbon, organic total	poten- tial, bottle	Sedi- ment, sus-	dis- charge, sus-
0at e	(mg/L as N)	(mg/L as N)	(mg/L _as N)	(mg/L as N)	(mg/L as NO3)	(mg/L as P)	(mg/L as P)	(mg/L as P)	(mg/L as C)	test (mg/L)	pended (mg/L)	pended (t/day)
larch 1977 14	0.52	0.090	0.59	1.2	5.3	0.310		0.020	4.9	17	125	0.26
ctober 19 20 ovember 1	. 31	.070	.14	.52	2.3	.040	.030		5.9	14	3	.00
08 arch 1978	. 46	.010	. 45	.92	4.1	.070	.050		9.0		10	.00
27						. 117	.016			7.5	110	.05

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

						INFLOW	SITE I21					
		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	Coli- form, fecal,	Strep- tococci	Solids, residue	0.1
		flow,	con-	pН	_	Oxygen,	(per-	total,	0.7	fecal	at 180°C	
		instan- taneous	duct- ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-mF (cols./	KF agar (cols./	dis- solved	dis- solved
Date	Time	(ft ³ /s)	(US/cm)	units)	(°C)	(mg/L)	at ion)	100 mL)	100 mL)	100 mL)	(mg/L)	(t/day)
March 1977 14 June 1977	1315	4.5	51	6.9	3.0	13.2	98	480B	25	200	22	0.27
02 October 197	1030	.25	94	6.9	14.0	9.6	92	6,200	5	1,100	51	.03
20 November 19	1550	.50	48	6.8	11.0				280	2,800		
08 March 1978	1300	1.8	58	6.8	10.0				6,300	8,000		
27	1440	.03	45	7.0	1.0							
Date March 1977	Nitro- gen, NO2+NO3 tota1 (mg/L as N)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)	Phos- phorus, dis- solved (mg/L as P)	Phos- phorus, ortho, total (mg/L as P)	Carbon, organic total (mg/L as C)	Algal growth poten- tial, bottle test (mg/L)	Sedi- ment, sus- pended (mg/L)	Sedi- ment, dis- charge, sus- pended (t/day)
14 June 1977	0.16	0.050	0.34	0.55	2.4	0.050		<0.010	4.2	0.8	92	1.1
02 October 197	.06	.050	. 19	.30	1.3	.020		<.010	7.0	.3	27	.02
20 November 19	. 10	.010	.12	.23	1.0	.010	.010			1.6	2	.00
08 March 1978	. 20	.030	. 18	.41	1.8	.050	.030		10	16	16	.08
27						.106	.016			4.4	180	.01
						INFLOW	SITE 122	<u> </u>				
		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	Coli- form, fecal,	Strep- tococci	Solids, residue	
		flow,	con-	pН		Oxygen,	(per-	total,	0.7	fecal	at 180°C	
		inst a n- taneous	duct- ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-MF (cols./	KF agar (cols./	dis- solved	
Date	Time	(ft ³ /s)	(µS/cm)	units)	(°C)	(mg/L)	at ion)	100 mL)	100 mL)	100 mL)	(mg/L)	
March 1977 14	1345	0.23	37	4.5	3.0	13.5	100	<8	3	46	19	
	olids, dis- solved	Nitro- gen, NO2+NO3 total (mg/L	Nitro- gen, ammonia dis- solved (mg/L	Nitro- gen, organic total (mg/L	Nitro- gen, total (mg/L	Nitro- gen, total (mg/L	Phos- phorus, total (mg/L	Phos- phorus, ortho, total (mg/L	Algal growth poten- tial, bottle test	Sedi- ment, sus- pended	Sedi- ment, dis- charge, sus- pended	
Date March 1977	(t/day)	as N)	as N)	as N)	as N)	as NO3)	as P)	as P)	(mg/L)	(mg/L)	(t/day)	
14	0.01	0.04	0.010	0.19	0.24	1.1	0.010	<0.010	0.0	23	0.01	

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

						INFLOW ST	TE 123	
								
		Stream- flow, instan- taneous	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature)×ygen, (dis-	kygen, dis- solved (per- cent satur-	Coli- form total, immed. (cols./
Date	Time	(ft ³ /s)	(µS/cm)	units)	(°C)	(mg/L) a	at ion)	100 mL)
March 1977								
14	1415		86	7.2	4.0	12.0	91	130B
14	1645	197	88	7.2	4.0	13.0	98	120B
June 1977								
01	1615	6.1	96	8.5	19.0	9.4	101	180
02	1115	7.8	97	8.7	19.5	10.2	111	3,400
07	1025	7.9	95	7.9	19.0	9.0	97	85
September 1								
20	1040	2.1	81	7.1	21.0	8.2	91	1,200
October 197								
20	1600	64	89	6.8	12.0			
November 19						•••	100	
08	1200	25	92	7.2	12.0	10.8	100	
08	1630	33	89	7.0	12.0			
March 1978 27	1410	160	94	7.2	2.0			
27		100		,,,				
	Coli-					Nitro-		
	form.	Strep-	Solids,		Nitro-	gen,	Nitro-	
	fecal,	tococci	residue		gen,	ammonia	gen,	
	0.7	fecal	at 180°C	Solide	NO2+NO3		_	^
		recal		201105	NUZ+NU3	dis-	organi	L
	μ M-mf	KF agar	dis-	dis-	total	solved	4.4.1	
	(cols./	(cols./					total	
Date				solved	(mg/L	(mg/L	(mg/L	
	100 mL)	100 mL)	solved (mg/L)	solved (t/day)	(mg/L as N)			
March 1977	100 mL)					(mg/L	(mg/L	
		100 mL)	(mg/L)		as N)	(mg/L as N)	(mg/L as N)	
14	35	100 mL) 200e	(mg/L) 43	(t/day)	as N) 0.23	(mg/L as N) 0.170	(mg/L as N) 0.66	
14 14		100 mL)	(mg/L)		as N)	(mg/L as N)	(mg/L as N)	
14 June 1977	35 <1	200e 420	(mg/L) 43 46	(t/day) 24	0.23 .23	(mg/L as N) 0.170 .150	(mg/L as N) 0.66 .34	
14 14 June 1977 01	35 <1 58	200e 420 220	(mg/L) 43 46 52	(t/day) 24 .86	0.23 .23	(mg/L as N) 0.170 .150	(mg/L as N) 0.66 .34 .73	
14 14 June 1977 01 02	35 <1	200e 420 220 220	(mg/L) 43 46	(t/day) 24 .86 1.2	0.23 .23 .01	(mg/L as N) 0.170 .150 .070 .050	(mg/L as N) 0.66 .34 .73 .95	
14 14 June 1977 01 02 07	35 <1 58 6 9	200e 420 220	(mg/L) 43 46 52 55	(t/day) 24 .86	0.23 .23	(mg/L as N) 0.170 .150	(mg/L as N) 0.66 .34 .73	
14 14 June 1977 01 02 07	35 <1 58 6 9	200e 420 220 220	(mg/L) 43 46 52 55	(t/day) 24 .86 1.2	0.23 .23 .01	(mg/L as N) 0.170 .150 .070 .050	(mg/L as N) 0.66 .34 .73 .95	
14 14 June 1977 01 02 07 September 1 20	35 <1 58 6 9 977	200e 420 220 220 160	43 46 52 55 64	.86 1.2 1.4	0.23 .23 .01 .01	(mg/L as N) 0.170 .150 .070 .050 .120	(mg/L as N) 0.66 .34 .73 .95 .02	
14 14 June 1977 01 02 07 September 1 20 October 197 20	35 <1 58 6 9 977 150 7	200e 420 220 220 160	43 46 52 55 64	.86 1.2 1.4	0.23 .23 .01 .01	(mg/L as N) 0.170 .150 .070 .050 .120	(mg/L as N) 0.66 .34 .73 .95 .02	
14 14 June 1977 01 02 07 September 1 20 October 197 20 November 19	35 <1 58 6 9 977 150 7 54	200e 420 220 220 160 360B	(mg/L) 43 46 52 55 64	.86 1.2 1.4	0.23 .23 .01 .01 .01 .50	(mg/L as N) 0.170 .150 .070 .050 .120 .170	(mg/L as N) 0.66 .34 .73 .95 .02 .60	
14 14 June 1977 01 02 07 September 1 20 October 197 20 November 19	35 <1 58 6 9 977 150 7 54	200e 420 220 220 160 360B	(mg/L) 43 46 52 55 64	.86 1.2 1.4	0.23 .23 .01 .01 .01 .50 .06	(mg/L as N) 0.170 .150 .070 .050 .120 .170 .180	(mg/L as N) 0.66 .34 .73 .95 .02 .60 .28	
14 14 June 1977 01 02 07 September 1 20 October 197 20 November 19 08	35 <1 58 6 9 977 150 7 54	200e 420 220 220 160 360B	(mg/L) 43 46 52 55 64	.86 1.2 1.4	0.23 .23 .01 .01 .01 .50	(mg/L as N) 0.170 .150 .070 .050 .120 .170	(mg/L as N) 0.66 .34 .73 .95 .02 .60	
14 14 June 1977 01 02 07 September 1 20 October 197 20 November 19	35 <1 58 6 9 977 150 7 54	200e 420 220 220 160 360B 190	(mg/L) 43 46 52 55 64	.86 1.2 1.4	0.23 .23 .01 .01 .01 .50 .06	(mg/L as N) 0.170 .150 .070 .050 .120 .170 .180	(mg/L as N) 0.66 .34 .73 .95 .02 .60 .28	

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

						INFLOW	SITE 124					
		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	Coli- form, fecal,	Strep- tococci	Solids, residue	
		flow,	con-	pН		Oxygen,	(per-	total,	0.7	feca1	at 180°C	Solids
		instan- taneous	duct- ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-MF (cols./	KF agar (cols./	dis- solved	dis- solved
Date	Time	(ft ³ /s)	(US/cm)	units)	(°C)	(mg/L)	at ion)	100 mL)	100 mL)	100 mL)	(mg/L)	(t/day
March 1977 14 June 1977	1435	1.2	46	4.8	4.0	13.6	103	∢8	<1	<1	23	0.07
01 October 197	1655	.22	61	6.4	11.0	10.6	95	23	<1	140	38	.02
20 November 19	1615	.25	39	5.7	10.0				2	8		
08 March 1978	1515	.30	43	5.4	10.0				5	28		
27	1340	2.6	45	6.9	2.0							
Date	Nitro- gen, NO2+NO3 total (mg/L as N)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)	Phos- phorus, dis- solved (mg/L as P)	Phos- phorus, ortho, total (mg/L as P)	Carbon, organic total (mg/L as C)	Algal growth poten- tial, bottle test (mg/L)	Sedi- ment, sus- pended (mg/L)	Sedi- ment, dis- charge, sus- pended (t/day)
March 1977 14 June 1977	0.03	0.010	0.30	0.34	1.5	0.020		<0.010	9.2	0.0	200	0.65
01 October 197	. 19	.010	. 66	.86	3.8	.010		<.010	4.9	.6	26	.02
20 November 19	<.10	<.010	.00	.01	.04	<.010	.010		5.4	40	2	.00
08 March 1978	.04	.010	.00	.04	. 18	.010	.010		11	.0	0	.00
27						.009	.017			.0	28	.20

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	Coli- form, fecal,	Strep- tococci	Solids, residue	
		flow,	con-	рH		Oxygen,	(per-	total,	0.7	fecal	at 180°C	Solids
		instan- taneous	duct- ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-MF (cols./	KF agar (cols./	dis- solved	dis- solved
Date	Time	(ft ³ /s)	(US/cm)	units)	(°C)	(mg/L)	at ion)	100 mL)	100 mL)	100 mL)	(mg/L)	(t/day)
March 1977 14	1455	0.68	40	4.5	4.0	13.8	105	∢8	<1	1	16	0.03
October 197 20 November 19	1630	.10	38	5.6	10.0				3	7		
08	1545	.10e	40	6.3	10.0				11	53		
		Nitro-								Algal	· · · · · · · · · · · · · · · · · · ·	Sed i -
	Nitro-	gen,	Nitro-				Phos-	Phos-		growth		ment.
	gen,	ammonia	gen,	Nitro-	Nitro-	Phos-	phorus,	phorus,	Carbon,	pot en -	Sedi-	dis-
	N02+N03	dis-	organic	gen,	gen,	phorus,	dis-	ortho,	organic	tial,	ment,	charge,
	total	solved	total	total	total	total	so lved	total	total	bottle	SUS-	sus-
Date	(mg/L as N)	(mg/L as N)	(mg/L as N)	(mg/L as N)	(mg/L as NO3)	(mg/L as P)	(mg/L as P)	(neg/L ars P)	(mg/L as C)	test (mg/L)	pended (mg/L)	pended (t/day)
4												
larch 1977 14	0.02	0.010	0.17	0.20	0.89	0.010		<0.010	1.6	0.0	4	0.00
14 October 197		0.010	0.17	0.20	0.09	0.010		~0.010	1.0	0.0	4	0.00
20	<.10	<.010	.06	.06	. 27	.010	.010		6.8	.1	2	.00
November 19												
08	<.10	<.010	.00	.01	.04	.020	.010		9.5	.2	8	

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

						INFLOW S	ITE 126				
		Stream- flow, instan- taneous	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Oxygen, dis-	xygen, dis- solved (per- cent satur-	Coli- form total, immed. (cols./			
Date	Time	(ft ³ /s)	(US/cm)	units)	(°C)	(mg/L)	at ion)	100 mL)	 	 	
June 1977	1620	4 -	140	2 1	14.0	10.0	06	36 000			
01 02	1630 1130	4.5 4.5	140 135	7.1 7.1	14.0 16.0	10.0 10.0		36,000 12,000			
October 197		1.0	133		10.0	10.0	101	11,000			
20	1515	30	82	6.7	10.0						
March 1978											
20	0905	14 14									
21 22	0906 1000	46									
23	0855	38			••						
24	1430	58									
25	0845	38									
26	1040	27									
27	0946	250									
27	1115	355									
27 27	1300 1420	340 310									
27	1450	310									
27	1530	280									
27	1615	260									
28	1145	89	61	6.8	3.0						
29	1135	64									
30	1310	55									
31	1300	38	42		3.5						
31							Nitro gen,			 	
31	Coli- form, fecal,	Strep- tococci	42 Solids, residue		3.5 Nitro- gen,	Nitro- gen, ammonia	Nitro gen,		 	 	
31	Coli- form,	38 Strep-	42 Solids,		3.5	Nitro- gen, ammonia	Nitro	i-		 	
	Coli- form, fecal, 0.7 µM-MF (cols./	Strep- tococci fecal KF agar (cols./	Solids, residue at 180°C dis-solved	Solids dis- solved	Nitro- gen, NO2+NO3 total (mg/L	Nitro- gen, ammonia dis- solved (mg/L	Nitro gen, organi tota (mg/L	ic			
Date	Coli- form, fecal, 0.7	Strep- tococci fecal KF agar	Solids, residue at 180°C dis-	Solids dis-	Nitro- gen, NO2+NO3 total	Nitro- gen, ammonia dis- solved	Nitro gen, organi tota	ic			
Date June 1977	Coli- form, fecal, 0.7 µM-MF (cols./ 100 mL)	Strep- tococci fecal KF agar (cols./ 100 mL)	Solids, residue at 180°C dissolved (mg/L)	Solids dis- solved (t/day)	Nitro- gen, NO2+NO3 total (mg/L as N)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro gen, organi tota (mg/L	ic			
Date June 1977 01	1300 Coli- form, fecal, 0.7 µM-MF (cols./ 100 mL)	Strep- tococci fecal KF agar (cols./ 100 mL)	Solids, residue at 180°C dissolved (mg/L)	Solids dis- solved (t/day)	3.5 Nitro- gen, NO2+NO3 total (mg/L as N)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro gen, organi tota (mg/L as N)	ic		 	
Date June 1977	1300 Coliform, fecal, 0.7 µM-MF (cols./100 mL) 14,000 1,100	Strep- tococci fecal KF agar (cols./ 100 mL)	Solids, residue at 180°C dissolved (mg/L)	Solids dis- solved (t/day)	Nitro- gen, NO2+NO3 total (mg/L as N)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro gen, organi tota (mg/L	ic			
Date June 1977 01 02 0ctober 197 20	1300 Coliform, fecal, 0.7 µM-MF (cols./100 mL) 14,000 1,100	Strep- tococci fecal KF agar (cols./ 100 mL)	Solids, residue at 180°C dissolved (mg/L)	Solids dis- solved (t/day)	3.5 Nitro- gen, NO2+NO3 total (mg/L as N)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro gen, organi tota (mg/L as N)	ic			
Date June 1977 01 02 0ctober 197 20	Coli- form, fecal, 0.7 µM-MF (cols./ 100 mL)	Strep- tococci fecal KF agar (cols./ 100 mL) 28,000 3,300	Solids, residue at 180°C dissolved (mg/L)	Solids dis- solved (t/day) 1.0 .90	Nitro- gen, NO2+NO3 total (mg/L as N) 0.62	Nitro- gen, ammonia dis- solved (mg/L as N) 0.190	Nitro gen, organi tota (mg/L as N) 0.72	ic			
Date June 1977 01 02 0ctober 197 20 March 1978 20 21	Coli- form, fecal, 0.7 µM-MF (cols./ 100 mL) 14,000 1,100	Strep- tococci fecal KF agar (cols./ 100 mL) 28,000 3,300 6,200	Solids, residue at 180°C dissolved (mg/L)	Solids dis- solved (t/day) 1.0 .90	3.5 Nitrogen, NO2+NO3 total (mg/L as N) 0.62 .35	Nitro- gen, ammonia dis- solved (mg/L as N) 0.190 .040	Nitro gen, organi tota (mg/L as N) 0.72 .19	ic			
Date June 1977 01 02 0ctober 197 20 March 1978 20 21 22	Coli- form, fecal, 0.7 µM-MF (cols./) 100 mL) 14,000 1,100	Strep- tococci fecal KF agar (cols./ 100 mL) 28,000 3,300 6,200	Solids, residue at 180°C dis-solved (mg/L)	Solids dis- solved (t/day) 1.0 .90	3.5 Nitro- gen, NO2+NO3 total (mg/L as N) 0.62 .35	Nitro- gen. ammonia dis- solved (mg/L as N) 0.190 .040	Nitro gen, organi tota (mg/L as N) 0.72 .19	ic			
Date June 1977 01 02 0ctober 197 20 March 1978 20 21 22 23	Coli- form, fecal, 0.7 µM-MF (cols./ 100 mL) 14,000 1,100	Strep- tococci fecal KF agar (cols./ 100 mL) 28,000 3,300 6,200	Solids, residue at 180°C dissolved (mg/L)	Solids dis- solved (t/day) 1.0 .90	Nitro- gen, N02+N03 total (mg/L as N) 0.62 .35	Nitro- gen, ammonia dis- solved (mg/L as N) 0.190 .040	Nitro gen, organi tota (mg/L as N) 0.72 .19	ic			
Date June 1977 01 02 0ctober 197 20 March 1978 20 21 22 23 24	Coli- form, fecal, 0.7 µM-MF (cols./ 100 mL) 14,000 1,100	Strep- tococci fecal KF agar (cols./ 100 mL) 28,000 3,300 6,200	Solids, residue at 180°C dis-solved (mg/L)	Solids dis- solved (t/day) 1.0 .90	Nitro- gen, N02+N03 total (mg/L as N) 0.62 .35 .17	Nitro- gen, ammonia dis- solved (mg/L as N) 0.190 .040	Nitro gen, organi tota (mg/L as N) 0.72 .19	ic			
Date June 1977 01 02 0ctober 197 20 March 1978 20 21 22 23 24 25	Coli- form, fecal, 0.7 µM-MF (cols./) 100 mL) 14,000 1,100	Strep- tococci fecal KF agar (cols./ 100 mL) 28,000 3,300 6,200	Solids, residue at 180°C dis-solved (mg/L) 85 74	Solids dis- solved (t/day) 1.0 .90	3.5 Nitro- gen, NO2+NO3 total (mg/L as N) 0.62 .35	Nitro- gen. ammonia dis- solved (mg/L as N) 0.190 .040	Nitro gen, organi tota (mg/L as N) 0.72 .19	ic			
Date June 1977 01 02 0ctober 197 20 March 1978 20 21 22 23 24 25	Coli- form, fecal, 0.7 µM-MF (cols./ 100 mL) 14,000 1,100	Strep- tococci fecal KF agar (cols./ 100 mL) 28,000 3,300 6,200	Solids, residue at 180°C dis-solved (mg/L) 85 74	Solids dis- solved (t/day) 1.0 .90	Nitro- gen, N02+N03 total (mg/L as N) 0.62 .35 .17	Nitro- gen, ammonia dis- solved (mg/L as N) 0.190 .040	Nitro gen, organi tota (mg/l as N) 0.72 .19	ic			
Date June 1977 01 02 0ctober 197 20 21 22 23 24 25 26 27	Coli- form, fecal, 0.7 µM-MF (cols./ 100 mL) 14,000 1,100	Strep- tococci fecal KF agar (cols./ 100 mL) 28,000 3,300 6,200	Solids, residue at 180°C dis-solved (mg/L)	Solids dis- solved (t/day) 1.0 .90	3.5 Nitrogen, NO2+NO3 total (mg/L as N) 0.62 .35 .17	Nitro- gen, ammonia dis- solved (mg/L as N) 0.190 .040 .010	Nitro gen, organi tota (mg/L as N) 0.72 .19	ic			
Date June 1977 01 0ctober 197 20 March 1978 20 21 22 23 24 25 26 27 27	Coli- form, fecal, 0.7 µM-MF (cols./ 100 mL) 14,000 1,100 520	Strep- tococci fecal KF agar (cols./ 100 mL) 28,000 3,300 6,200	Solids, residue at 180°C dissolved (mg/L) 85 74	Solids dis- solved (t/day) 1.0 .90	3.5 Nitrogen, N02+N03 total (mg/L as N) 0.62 .35 .17	Nitro- gen, ammonia dis- solved (mg/L as N) 0.190 .040 .010	Nitro gen, organi tota (mg/L as N) 0.72 .19 .12	ic			
Date June 1977 01 02 0ctober 197 20 March 1978 20 21 22 23 24 25 27 27 27	Coli- form, fecal, 0.7 µM-MF (cols./ 100 mL) 14,000 1,100 520	38 Strep-tococci fecal KF agar (cols./) 100 mL) 28,000 3,300 6,200	Solids, residue at 180°C dis-solved (mg/L) 85 74	Solids dis- solved (t/day) 1.0 .90	3.5 Nitrogen, N02+N03 total (mg/L as N) 0.62 .35	Nitro- gen, ammonia dis- solved (mg/L as N) 0.190 .040 .010	Nitro gen, organi tota (mg/L as N) 0.72 .19 .12	ic			
Date June 1977 01 02 0ctober 197 20 March 1978 20 21 24 25 26 27 27 27	1300 Coliform, fecal, 0.7 µM-MF (cols./100 mL) 14,000	38 Streptococci fecal KF agar (cols./ 100 ml) 28,000 3,300 6,200	\$50 ids, residue at 180°C dis- solved (mg/L) 85 74	Solids dis- solved (t/day) 1.0 .90	3.5 Nitrogen, N02+N03 total (mg/L as N) 0.62 .35 .17	Nitro- gen, ammonia dis- solved (mg/L as N) 0.190 .040	Nitro gen, organi tota (mg/L as N) 0.72 .19 .12	ic			
Date June 1977 01 02 0ctober 197 20 21 22 23 24 25 26 27 27 27 27 27	Coli- form, fecal, 0.7 µM-MF (cols./ 100 mL) 14,000 1,100	38 Strep-tococci fecal KF agar (cols./ 100 mL) 28,000 3,300 6,200	Solids, residue at 180°C dis-solved (mg/L) 85 74	Solids dis- solved (t/day) 1.0 .90	3.5 Nitrogen, N02+N03 total (mg/L as N) 0.62 .35 .17	Nitro- gen, ammonia dis- solved (mg/L as N) 0.190 .040 .010	Nitro gen, organi tota (mg/L as N) 0.72 .19 .12	ic			
Date June 1977 01 02 02 0ctober 197 20 21 22 23 24 25 26 27 27 27 27 27 27	1300 Coliform, fecal, 0.7 µM-MF (cols./100 mL) 14,000	38 Streptococci fecal KF agar (cols./ 100 ml) 28,000 3,300 6,200	\$50 ids, residue at 180°C dis- solved (mg/L) 85 74	Solids dis- solved (t/day) 1.0 .90	3.5 Nitrogen, N02+N03 total (mg/L as N) 0.62 .35 .17	Nitro- gen, ammonia dis- solved (mg/L as N) 0.190 .040	Nitro gen, organi tota (mg/L as N) 0.72 .19 .12	ic			
Date June 1977 01 02 02 0ctober 197 20 21 22 23 24 25 27 27 27 27 27 28	Coli- form, fecal, 0.7 µM-MF (cols./ 100 mL) 14,000 1,100 520	38 Strep-tococci fecal KF agar (cols./ 100 mL) 28,000 3,300 6,200	Solids, residue at 180°C dis-solved (mg/L) 85 74	Solids dis- solved (t/day) 1.0 .90	3.5 Nitrogen, N02+N03 total (mg/L as N) 0.62 .35 .17	Nitro- gen, ammonia dis- solved (mg/L as N) 0.190 .040 .010	Nitro gen, organi tota (mg/L as N) 0.72 .19 .12	ic			
Date June 1977 01 02 02 0ctober 197 20 21 22 23 24 25 26 27 27 27 27 27	Coli- form, fecal, 0.7 µM-MF (cols./ 100 mL) 14,000 1,100 520	38 Strep-tococci fecal KF agar (cols./) 100 mL) 28,000 3,300 6,200	Solids, residue at 180°C dis-solved (mg/L) 85 74	Solids dis- solved (t/day) 1.0 .90	3.5 Nitrogen, N02+N03 total (mg/L as N) 0.62 .35 .17	Nitro- gen, ammonia dis- solved (mg/L as N) 0.190 .040 .010	Nitro gen, organi tota (mg/L as N) 0.72 .19 .12	ic			

Table 7.--Chemical, physical, and biological characteristics of surface-water inflows--Continued

							SITE 127					
		Stream-	Spe- cific				Oxygen, dis- solved	Coli- form	Coli- form, fecal,		Solids, residue	
		flow,	con-	pН		Oxygen,	(per-	total,	0.7	fecal	at 180°C	Solids
		instan- taneous	duct - ance	(stand- ard	Temper- ature	dis- solved	cent satur-	immed. (cols./	µM-MF (cols./	KF agar (cols./	dis- solved	dis- solved
Date	Time	(ft^3/s)	(µS/cm)	units)	(°C)	(mg/L)	ation)	100 mL)	100 mL)	100 mL)	(mg/L)	(t/day
June 1977												
02 September	1225 1977	4.5	126	6.9	17.0	9.6	99	5,000	1,200B	1,300	76	0.92
20 March 1978	1350	6.5	165	7.1	16.0			46,000B	29,000B	48,000E		
27	1730	239	55	6.6	3.0							
		Nitro-								Algal		Sed i-
	Nitro-	gen,	Nitro-	M 2 A	W	Dhira	Phos-	Phos-	Carrhan	growth	6. 4:	ment,
	gen, NO2+NO3	ammonia dis-	gen, organic	Nitro- gen,	Nitro- gen,	Phos- phorus,	phorus, dis-	phorus, ortho.	Carbon, organic	poten- tial,	Sedi- ment.	dis- charge,
	total	solved	total	total	total	total	solved	total	total	bottle	sus-	sus-
	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	(mg/L	test	pended	pended
Date	as N)	às N)	às N)	as N)	as NO3)	as P)	às P)	às P)	às C)	(mg/L)	(mg/L)	(t/day)
June 1977												
02	0.37	0.070	0.26	0.70	3.1	0.040		0.010	5.0	19	16	0.19
September												
20	<.10	. 050	.30	.35	1.6	.030	.020		7.0	41	13	.23
March 1978												

Table 11.--Chemical, physical, and biological characteristics of lake water $[\text{ft, feet; } \mu\text{S/cm, microsiemens per centimeter; }^{\circ}\text{C, degrees Celsius; m, meters; } \\ \text{mg/L, milligrams per liter; FET-fld, fixed end-point titration in the field<, less than; --, no data}$

			Lake	site L1			
	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Oxygen, dis- solved	Oxygen, dis- solved (per- cent satur-
Date		(ft)	(µS/cm)	units)	(°C)	(mg/L)	ation)
Ann:1 10	77						
April 19 04	1315	3.30	69	6.7	6.0	14.0	112
04	1316	6.60	70	6.7	6.0	14.0	112
04	1317	9.90	70	6.5	6.0	13.9	111
04	1318	13.2	70	6.9	6.0	13.9	111
26	1155	1.60	88	8.4	12.0	11.9	110
26	1156	3.30	88	8.4	11.5	11.9	109
26	1157	6.60	88	8.4	11.5	11.9	109
May 1977		0.00	~	0.4	11.5	,	103
16	1145	1.60	80	7.5	14.0	10.6	102
16	1146	3.30	80	8.0	13.0	11.2	106
16	1147	5.00	80	8.0	13.0	11.2	106
June 197			-	0.0	10.0		
13	1135	1.60	84	6.8	17.5	9.1	95
13	1137	3.30	84	5.5	17.0	8.8	91
13	1139	6.60	85	5.5	16.5	8.6	88
13	1141	8.20	85	5.5	16.5	8.3	85

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

						Lake site	. L2					
-	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Trans- par- ency (secchi disk)	0xygen, dis- solved	Oxygen, dis- solved (per- cent satur-	Bicar- bonate FET-fld (mg/L as	Alka- linity field (mg/L as	Carbon dioxide dis- solved (mg/L	Nitro- gen, NO2+NO3 total (mg/L
Date		(ft)	(µS/cm)	units)	(°C)	(m)	(mg/L)	ation)	HC03)	CACO3)	as CO2)	as N)
	_											
April 197												
04	1400	1.60				1.40			1B	15		.08
04	1401	3.30	73	6.3	6.0		13.8	110				
04	1402	6.60	73	6.3	6.0		13.8	110				
04	1405	8.20							18	15		.08
04	1406	9.90	73	6.2	6.0		13.8	110				
04	1407	13.2	74	6.1	6.0		13.8	110				
04	1410	15.0							17	14		.10
04	1411	16.5	74	6.5	6.0		13.8	110				
26	1210	1.60	80	8.3	12.0	1.70	11.9	110	23	19	.2	.01
26	1211	3.30	81	8.0	11.5		11.6	106				
26	1214	6.60	79	7.3	11.5		11.1	102				
26	1215	8.20							18	15		.01
26	1217	9.80	76	7.4	11.0		11.6	105				
26	1219	13.2	B1	7.4	10.0		11.7	104				
26	1220	15.0	88	7.1	10.0		11.2	99	22	18	2.8	.06
May 1977	1110	10.0		,	1010							
04	1125	15.0										
16	1220	1.60	83	7.7	14.0	1.00	10.9	105	25	21	.8	< .10
16	1223	3.30	82	7.8	13.0	1.00	11.0	104				
16	1224	6.60	82	7.0	13.0		11.1	105				
	1225								19	16		< .10
16		8.20	81		12.5		11.0	102	_			
16	1226	9.80		7.7				103				
16	1227	13.2	81	7.4	12.5		10.8	101				
16	1230	15.0	81	7.0	12.5		10.1	94	19	16	3.0	.01
June 1977												
13	1200	1.60	85	6.7	17.0	2.20	9.0	93		16	6.2	<.10
13	1202	3.30	85	5.5	17.0		8.8	91				
13	1204	6.60	86	5.4	16.0		8.0	81				
13	1205	8.20								18		< .10
13	1207	9.80	85	5.3	16.0		8.0	81	~-			
13	1209	13.2	67	5.3	16.0		6.6	67				
13	1210	14.8	85	5.3	16.0		6.4	65		18	175	< .10
July 1977	7											
11	1105	1.60	92	7.7	24.5	2.90	9.3	110		19	.7	.02
11	1107	3.30	92	7.7	24.5		9.3	110	~-			
11	1108	6.60	92	7.7	24.0		9.2	109				
11	1110	7.40						103		19		<.10
11	1112	9.80	92	7.7	24.0		9.2	108				1.10
11	1115	13.0	92	7.6	23.0		7.8	90		20	1.0	.01
August 19		13.0	76	7.0	23.0		7.0	30		20	1.0	.01
		1 60	0.4	7 6	24 5	2 60	0.0	107		10	1 2	- 10
04	1155	1.60	94	7.5	24.5	2.60	9.0	107		19	1.2	<.10
04	1157	3.30	94	7.4	24.5		9.0	107				
04	1159	6.60	94	7.4	24.0		8.9	105				

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

								0xygen,			
	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Trans- par- ency (secchi disk)	0×ygen, dis- solved	dis- solved (per- cent satur-	Alka- linity field (mg/L as	Carbon dioxide dis- solved (mg/L	Nitro- gen, NO2+NO3 total (mg/L
Date		(ft)	(µS/cm)	units)	(°C)	(m)	(mg/L)	at ion)	CACO3) as	C02)	as N)
August	1977										
04	1200	7.40							19		<.10
04	1202	9.80	94	7.3	24.0		8.2	96			
04	1205	13.0	115	7.3	23.0		3.8	44	20	1.9	.01
Septemb				.,.	20.0		0.0				
06	1440	1.60			24.0	2.70	8.4	99	20		<.10
06	1441	3.30			24.0		8.3	98			
06	1442	6.60			24.0		8.2	96	20		<.10
06	1443	9.80			23.5		6.4	74			
06	1444	13.2			23.0		4,2	48	21		<.10
07	1230	1.60	87		23.5		7.8	91			
07	1231	3.30	87		23.5		7.8	91			
07	1232	6.60	87		23.0		7.6	87			
07	1236	9.80	87		23.0		7.3	84			
07	1240	13.2	87		23.0		7.3 5.9	68			
October		13.2	u,		23.0		3.5	00			
04	1255	1.60	97		15.0	1.80	8.5	83	22		.03
04	1256	3.30	97		15.0	1.00	8.5	83			
04	1257	6.60	99		15.0		8.6	84			
04	1300	7.40							22		
04	1300	9.80	99		15.0		8.4	82			.03
04	1301	13.2	99		15.0		8.4	82	23		
21	0945	1.60		7.4	11.5	1.70			20	1.5	. 04 . 02
21	0945	3.30	80 80	7.4	11.5	1.70	8.1	74 74	20	1.5	
21				7.4							
21	0947 0950	6.60 8.20	80	7.3	11.5		8.0	73	20		
21	0951	9.80		7.3	11.5		8.0		-		.04
			80	7.3				73			
21	0952	13.2	80		11.5		8.0	73	20	1	
21	0955	14.9	80	7.4	11.5		8.1	74	20	1.5	.03
Novembe		1.60	20	7.4	0.0						
16	1315	1.60	80	7.4	9.0		5.0	43			
16	1316	3.30	80	7.3	9.0		5.0	43			
16	1317	6.60	80	7.4	9.0		5.0	43			
16	1318	9.80	80	7.4	9.0		5.0	43			
16	1319	13.2	80	7.4	9.0		4.9	42			
16	1320	14.9	80	7.4	9.0		4.9	42			

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

			Nitro-							Algal
			gen, ammonia	Nitro- gen,	Nitro-	Nitro-	Phos-	Phos- phorus,	Carbon,	growth poten-
		Sam- pling	dis-	organic	gen,	gen,	phorus,	ortho,	organic	tial,
0-1-	Time	depth	solved (mg/L	total (mg/L	total (mg/L	total (mg/L	total (mg/L	total (mg/L	total (mg/L	bottle test
Date		(ft)	as N)	as N)	as N)	as NO3)	as P)	as P)	as C)	(mg/L)
April 19	77									
04	1400	1.60	.080	. 52	.68	3.0	.010	<.010	3.1	.8
04	1401	3 .3 0								
04	1402	6.60								
04	1405	8.20	.080	. 42	.58	2.6	.010	<.010	5.8	
04	1406	9.90								
04	1407	13.2								
04	1410	15.0	.060	.74	.90	4.0	.020	<.010	6.7	
04	1411	16.5								
26	1210	1.60	. 020	. 40	.43	1.9	.020	<.010		
26	1211	3.30								
26	1214	6.60								
26	1215	8.20	.070	. 49	.57	2.5	.020	<.010		
26	1217	9.80								
26	1219	13.2								
26	1220	15.0	.060	. 81	.93	4.1	.070	<.010		
May 1977										
04	1125	15.0								
16	1220	1.60	.030	. 42	.45	2.0	.030	<.010		
16	1223	3.30								
16	1224	6.60								
16	1225	8.20	.050	. 35	.40	1.8	.030	<.010		
16	1226	9.80								
16	1227	13.2								
16	1230	15.0	.030	. 61	.65	2.9	.070	<.010		
June 197	7									
13	1200	1.60	. 050	. 28	.33	1.5	.020	<.010		
13	1202	3.30						••		
13	1204	6.60								
13	1205	8.20	.090	.27	.36	1.6	.030	<.010		
13	1207	9.80								
13	1209	13.2								
13	1210	14.8	. 120	. 35	.47	2.1	.040	<.010		
July 197										
11	1105	1.60	.050	. 61	.68	3.0	.020			
11	1107	3.30								
11	1108	6.60								
11	1110	7.40	.050	. 39	.44	1.9	.020			
11	1112	9.80								
11	1115	13.0	.070	. 41	.49	2.2	.020			
August 1										
04	1155	1.60	.010	. 45	.46	2.0	.020		4.3	.8
04	1157	3.30					.020			

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

Nitrogen, Nitrogen, Nitrogen, Sampling Sampli				Lake	site L2	Continued	<u>l</u>		
04 1200 7.40 .020 .53 .55 2.4 .030 8.0 04 1205 13.0 .030 .39 .43 1.9 .050 5.9 September 1977 06 1440 1.60 .020 .31 .33 1.5 .030 06 1441 3.30	Date	Time	pling depth	gen, ammonia dis- solved (mg/L	gen, organic total (mg/L	gen, total (mg/L	gen, total (mg/L	phorus, total (mg/L	organic total (mg/L
04 1200 7.40 .020 .53 .55 2.4 .030 8.0 04 1205 13.0 .030 .39 .43 1.9 .050 5.9 September 1977 06 1440 1.60 .020 .31 .33 1.5 .030 06 1441 3.30	August 1	977							
04 1202			7.40	.020	.53	. 55	2.4	.030	8.0
04 1205 13.0 .030 .39 .43 1.9 .050 5.9 September 1977 06 1440 1.60 .020 .31 .33 1.5 .030 06 1441 3.30 <									
September 1977 06 1440									5 9
06 1440			10.0		.05			.030	0.5
06 1441			1.60	.020	.31	. 33	1.5	.030	
06 1442 6.60 .020 .33 .35 1.6 .030 06 1443 9.80 06 1444 13.2 .110 .37 .48 2.1 .050 06 1445020 .33 06 1445020 .33 06 1450110 .37 07 1230 1.60 07 1231 3.30 07 1232 6.60 07 1236 9.80 07 1240 13.2 07 1255 1.60 .050 .38 .46 2.0 .030 04 1255 3.30 04 1257 6.60 04 1300 7.40 .040 .38 .45 2.0 .100 04 1301 9.80 04 1301 9.80 04 1305 13.2 .040 .42 .50 2.2 .100 04 1305 13.2 .040 .42 .50 2.2 .100 21 0945 1.60 .010 .89 .92 4.1 .020 21 0947 6.60 21 0947 6.60 21 0951 9.80 21 0952 13.2 21 0951 9.80 21 0952 13.2 21 0951 9.80 16 1316 3.30									
06 1443									
06 1444 13.2 .110 .37 .48 2.1 .050 06 1445020 .33 06 1450110 .37 07 1230 1.60 07 1231 3.30 07 1236 9.80 07 1240 13.2 07 1256 9.80 07 1256 3.30 04 1256 3.30 04 1257 6.60 04 1300 7.40 .040 .38 .46 2.0 .030 04 1301 9.80 04 1305 13.2 .040 .42 .50 2.2 .100 04 1305 13.2 .040 .42 .50 2.2 .100 21 0945 1.60 .010 .89 .92 4.1 .020 21 0946 3.30 21 0947 6.60 21 0950 8.20 .030 .55 .62 2.7 .030 21 0951 9.80 21 0952 13.2 21 0955 14.9 .030 .80 .86 3.8 .020 November 1977 16 1315 1.60 16 1317 6.60 16 1318 9.80									
06 1445									
06 1450						-			
07 1230 1.60									
07 1231 3.30					-				
07 1232 6.60									
07 1236 9.80									
07 1240 13.2									
October 1977 04 1255 1.60 .050 .38 .46 2.0 .030 04 1256 3.30									
04 1255			10.1						
04 1256			1.60	.050	. 38	. 46	2.0	.030	
04 1257 6.60									
04 1300 7.40 .040 .38 .45 2.0 .100 04 1301 9.80 04 1305 13.2 .040 .42 .50 2.2 .100 21 0945 1.60 .010 .89 .92 4.1 .020 21 0946 3.30 21 0950 8.20 .030 .55 .62 2.7 .030 21 0951 9.80 21 0952 13.2 21 0955 14.9 .030 .80 .86 3.8 .020 November 1977 16 1315 1.60 16 1316 3.30 16 1318 9.80 16 1318 9.80 16 1318 9.80 16 1319 13.2									
04 1301 9.80									
04 1305 13.2 .040 .42 .50 2.2 .100 21 0945 1.60 .010 .89 .92 4.1 .020 21 0947 6.60 21 0950 8.20 .030 .55 .62 2.7 .030 21 0951 9.80 21 0952 13.2 21 0955 14.9 .030 .80 .86 3.8 .020 November 1977 16 1315 1.60 16 1317 6.60 16 1318 9.80 16 1318 9.80 16 1319 13.2									
21 0945									
21 0946 3.30									
21 0947 6.60									
21 0950 8.20 .030 .55 .62 2.7 .030 21 0951 9.80 21 0952 13.2 21 0955 14.9 .030 .80 .86 3.8 .020 November 1977 16 1315 1.60 16 1316 3.30 16 1317 6.60 16 1318 9.80 16 1318 9.80 16 1319 13.2									
21 0951 9.80 21 0952 13.2 21 0955 14.9 .030 .80 .86 3.8 .020 November 1977 16 1315 1.60 16 1316 3.30 16 1317 6.60									
21 0952 13.2									
21 0955 14.9 .030 .80 .86 3.8 .020 November 1977 16 1315 1.60 16 1316 3.30 16 1317 6.60 16 1318 9.80 16 1319 13.2									
November 1977 16 1315									
16 1315									
16 1316 3.30 16 1317 6.60 16 1318 9.80			1.60						
16 1317 6.60 16 1318 9.80 16 1319 13.2									
16 1318 9.80 16 1319 13.2									
16 1319 13.2									
AV.,, AVEO 17.J	16	1320	14.9						

Phytoplankton analyses

			Þ	hytoplar	nkton anal	yses						
DATE TIME		4,77 1400		26•77 200		26•77 210		16•77 220		13•77 200		11•77 105
DIVEPSITY: DIVISION .CLASS .ORDERFAMILYGENUS		0.0 0.0 1.0 1.0		0.9 0.9 1.6 1.6		1.0 1.0 1.0 1.0		0.9 0.9 0.9 1.6 1.6		0.9 0.9 1.6 1.6		0.9 0.9 1.6 1.6
ORGANISM	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT
CHLOROPHYTA (GREEN ALGAE) •CHLOROPHYCEAE												
CHLOROCOCCALESCHARACIACEAE												
SCHROEDERIAMICRACTINIACEAE		-	•	0		-		-		-		-
MICPACTINIUMOOCYSTACEAE		-	*	0		-	*	0		-		-
ANKISTRODESMUS		0		0		-	*	0 0		-		-
CHODATELLACLOSTERIOPSIS		0 -		-		-		-		-		-
DICTYOSPHAERIUMKIRCHNERIELLA		-		-		-	*	0	#	33	2500# 	26
NEPHROCYTIUMOOCYSTIS		-		-		-		-		-	59	1
QUADRICOCCUS		-		-		-		0		-		-
TETRAEDRONSCENEDESMACEAE	*	0		-		-	*	0		-		-
••••SCENEDESMUS ••TETRASPORALES	*	0		-		-	*	0		-		-
PALMELLACEAE									_			_
SPHAEROCYSTISVOLVOCALES		-		-		-		-		-		•
CHLAMYDOMDNADACEAECHLAMYDOMONAS		0		-		-		-		_		_
ZYGNEMATALES		·										
DESMIDIACEAECLOSTERIUM		_		-		-		0		-		-
····COSMARIUM ····STAURASTRUM		-		-		-		-		-	69 #	1
CHRYSOPHYTA												
.BACILLARIOPHYCEAECENTRALES												
COSCINODISCACEAE				•		_		•		_		_
CYCLOTELLA MELOSIRA		\$ 50 0	*	0		-		0		-		-
STEPHANODISCUSPENNALES		-	*	0		-	*	0		-		-
ACHNANTHACE AE	_	•				_		_		_		_
ACHNANTHESCYMBELLACEAE	*	0		-		-		-		_		
CYMBELLAFRAGILARIACEAE		-		0		-		-		-		-
ASTERIONELLA	. 1		#	33	•	5 0		# 33		-		-
····FRAGILARIA ····SYNEDRA		0	*	0		-	*	ō		-		-
MERIDIONACEAEMERIDION		_		0		_		-		-		_
NITZSCHIACEAE		0				_		_		_		_
NITZSCHIA TABELLARIACEAE	٠	U				-		_		_	_	
*TABELLARIA .CHRYSOPHYCEAE		-		0		-		-		-		•
CHRYSOMONADALES OCHROMONADACEAE												
DINOBRYON		0		-		-		-		-		-
CYANOPHYTA (BLUE-GREEN ALGAE) •CYANOPHYCEAE												
CHROCCOCCALES												
CHROCCOCCAEAEANACYSTIS		_	*	33		-		-		-		-
HORMOGONALES NOSTOCACEAE												
ANABAENA		-		-		- + 50		33		-	3900#	41
APHANIZOMENONOSCILLATORIACEAE	•	0	*	0	•	50						-
OSCILLATORIA CHROCCOCCALES		0	#	3 3		-	*	3 3	#	33		-
CHROCCOCCAEAE				0		_		_		. 23	3000#	: 31
••••GOMPHOSPHAERIA EUGLENOPHYTA (EUGLENOIDS)		-	*	0		-		-	#	33	3000#	. 31
.EUGLENOPHYCEAEEUGLENALES												
EUGLENACEAE EUGLENA		_		_		-		-		_	-	_
····TRACHELOMONAS		-		-		-		-		-	*	0
PYRRHOPHYTA (FIRE ALGAE)												
.DINOPHYCEAE PERIDINIALES												
PERIDINIACEAE				•					•	_		_
PERIDINIUM		-	*	0		-		-		•		-

^{* =} Present in sample but not a dominant type.
= A dominant type in sample.

Phytoplankton analyses

			Phy	toblankt	on analys	es				
DATE TIME		4•77 1155		17,77 1115		6•77 .440		4 , 77 255		21•77 945
DIVERSITY: PIVISION		1.5		1.5		0.0		1.0		1.5
•CLASS		1.5		1.5		0.0		1.5		1.5
••ORDER		1.5		1.5		1.0		2.3		2.0
FAMILY		1.5		1.5		1.0		5.6		2.0
· · · · GENUS		2.0		2.0		2.0		2.6		2.0
ORGANISM	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PFR- CENT	CELLS /ML	PER- CENT
CHLOROPHYTA (GREEN ALGAE)										
· CHLOROPHYCFAE										
CHLOROCOCCALES										
CHARACIACEAESCHROEDERIA		_		_		_		_		_
MICRACTINIACEAE										
MICRACTINIUM		-		-		-		-		-
OOCYSTACEAE										
ANKISTRODESMUS		0		-		-		-		_
CHOOATELLACLOSTERIOPSIS	*	0		-		-		_		_
DICTYOSPHAERIUM		Ö		_		_		-		_
KIRCHNERIELLA		_		-		-		-		-
NEPHROCYTIUM		0		-		-		-		-
OOCYSTIS		0		-		-		-		-
QUADRICOCCUS		-		-		-		-		-
TETRAEDRONSCENEDESMACEAE		-		-		-		-		-
SCENEDESMUS		0		_		-		_		-
TETRASPORALES		•								
PALMELLACEAE										
SPHAEROCYSTIS	*	25	•	ŧ 2S		-		-		-
VOLVOCALESCHLAMYDOMONADACEAE							,			
· · · · CHLAMYDOMONAS		_		_		-		-		_
ZYGNEMATALES										
DESMIDIACEAE										
CLOSTERIUM		-		-		-		-		-
COSMARIUM		-		-		-		-		-
STAURASTRUM		0		-		-		-		_
CHRYSOPHYTA										
.BACILLARIOPHYCEAE										
CENTRALES										
COSCINODISCACEAE										
CYCLDTELLA		-		-		-				-
MELOSIRASTEPHANODISCUS		-		-		-		17		25
PENNALES		_		_		_		_		_
ACHNANTHACEAE										
ACHNANTHES		-		-		-		-		-
CYMBELLACEAE										
CYMBELLA		-		-		-		-		-
FRAGILARIACEAEASTERIONELLA		_		_		_	#	17		-
FRAGILARIA		-		-		_		-		-
SYNEDRA		-		-		-		-		-
MERIDIONACEAE										
MERIDION		-		-		-		-		-
NITZSCHIACEAENITZSCHIA		0		_		_		_		_
TABELLARIACEAE		•								
TABELLARIA		-		-		-		-		-
.CHRYSOPHYCEAE										
CHRYSOMONADALES										
•••OCHROMONADACEAE ••••DINOBRYON		_		_		_		17		_
• • • • DINOBRION							"	.,		
CYANOPHYTA (BLUE-GREEN ALGAE)										
.CYANOPHYCEAE										
CHROCCOCCALES										
CHROCCOCCAEAE		0		_	1 #	25		_		_
ANACYSTISHORMOGONALES	-	U		_	1"	23				
NOSTOCACEAE										
ANABAENA	#	25		25		25		-		-
APHANIZOMENON	*	25	#	25	1 #	25	#	17	#	25
OSCILLATORIACEAE										
OSCILLATORIACHROCCOCCALES		-		-		-	*	17		-
CHROCCOCCAEAE										
GOMPHOSPHAERIA		0		_	1#	25	#	17	#	25
EUGLENOPHYTA (EUGLENOIDS)										
• EUGLENOPHYCEAE										
EUGLENALES										
EUGLENACEAE		0		_		_		_		_
TRACHELOMONAS		25		25		-		-	#	25
		-		-						-
PYRRHOPHYTA (FIRE ALGAE)										
•PINOPHYCEAE										
PERIDINIALESPERIDINIACEAE										
PERIDINIACEAE		-		-		-		_		_
* - Procent in sample but not a domina	ant type									

^{* =} Present in sample but not a dominant type.
= A dominant type in sample.

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

		Lake	site L3				
Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Oxygen, dis- solved	Oxygen, dis- solved (per- cent satur-	
 Date	(ft)	(µS/cm)	units)	(°C)	(mg/L)	ation)	
April 1977							
04 1500	3.30	70	6.9	6.0	14.0	112	
04 1501	6.60	70	6.9	6.0	14.0	112	
04 1502	8.20	70	6.9	6.0	14.0	112	
26 1245	1.60	83	8.1	12.0	11.6	107	
26 1246	3.30	83	8.2	11.5	11.7	107	
26 1247	6.60	83	7.6	11.5	11.5	106	
26 1248	9.80	89	7.2	11.0	10.8	98	
May 1977							
16 1235	1.60	84	7.8	14.5	11.0	107	
16 1236	3.30	84	7.8	14.0	11.0	106	
16 1237	6.60	83	7.6	13.0	10.9	103	
16 1238	9.80	83	7.4	13.0	10.8	102	
16 1239	11.0	83	7.4	13.0	10.7	101	
June 1977							
13 1235	1.60	8 6	6.1	17.0	9.0	93	
13 1237	3.30	85	5.6	16.5	8.0	82	
13 1239	6.60	85	5.4	16.5	7.5	77	
13 1241	8.20	85	5.4	16.5	7.0	72	
 		Spe- cific				0xygen, dis-	
	Sam-	con-	pН		Oxygen,	solved (per-	
Time	pling	con- duct-	(stand-	Temper- ature	Oxygen, dis- solved	(per- cent	
Time Nato	pling depth	con- duct- ance	(stand- ard	ature	dis- solved	(per- cent satur-	
Time Date	pling	con- duct-	(stand-		dis- solved	(per- cent	
Date April 1977	pling depth (ft)	con- duct- ance (µS/cm)	(stand- ard units)	ature (°C)	dis- solved (mg/L)	(per- cent satur- ation)	
 Date April 1977 04 1515	pling depth (ft)	con- duct- ance (µS/cm)	(stand- ard units) 6.9	ature (°C) 6.0	dis- solved (mg/L)	(per- cent satur- ation)	
Date April 1977 04 1515 04 1516	pling depth (ft) 1.60 3.30	con- duct- ance (µS/cm) 75 76	(stand- ard units) 6.9 7.0	ature (°C) 6.0 6.0	dis- solved (mg/L) 14.1 14.1	(per- cent satur- ation) 113 113	
Date April 1977 04 1515 04 1516 04 1517	pling depth (ft) 1.60 3.30 6.60	con- duct - ance (µS/cm) 75 76 76	(stand- ard units) 6.9 7.0 7.0	ature (°C) 6.0 6.0 6.0	dis- solved (mg/L) 14.1 14.1 14.1	(per- cent satur- ation) 113 113	
Date April 1977 04 1515 04 1516 04 1517 04 1518	pling depth (ft) 1.60 3.30 6.60 9.80	con- duct - ance (µS/cm) 75 76 76 76	(stand- ard units) 6.9 7.0 7.0 7.0	ature (°C) 6.0 6.0 6.0 6.0	dis- solved (mg/L) 14.1 14.1 14.1 14.1	(per- cent satur- ation) 113 113 113	
Date April 1977 04 1515 04 1516 04 1517 04 1518 04 1519	pling depth (ft) 1.60 3.30 6.60 9.80 11.6	con- duct - ance (//S/cm) 75 76 76 76 76 77	(stand- ard units) 6.9 7.0 7.0 7.0 7.1	6.0 6.0 6.0 6.0 6.0	dis- solved (mg/L) 14.1 14.1 14.1 14.1 14.0	(per- cent satur- ation) 113 113 113 113 113	
Date April 1977 04 1515 04 1516 04 1517 04 1518 04 1519 26 1355	pling depth (ft) 1.60 3.30 6.60 9.80 11.6 1.60	con- duct - ance (µS/cm) 75 76 76 76 77 81	(stand- ard units) 6.9 7.0 7.0 7.0 7.1 9.0	6.0 6.0 6.0 6.0 6.0 12.0	dis- solved (mg/L) 14.1 14.1 14.1 14.0 13.0	(per- cent satur- ation) 113 113 113 113 112 120	
Date April 1977 04 1515 04 1516 04 1517 04 1518 04 1519 26 1355 26 1356	1.60 3.30 6.60 9.80 11.6 1.60 3.30	con- duct - ance (µS/cm) 75 76 76 76 77 81 82	(stand- ard units) 6.9 7.0 7.0 7.0 7.1 9.0 9.0	6.0 6.0 6.0 6.0 6.0 12.0	dis- solved (mg/L) 14.1 14.1 14.1 14.0 13.0 13.0	(per- cent satur- ation) 113 113 113 113 112 120	
Date April 1977 04 1515 04 1516 04 1517 04 1518 04 1519 26 1355 26 1355 26 1355	pling depth (ft) 1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60	con- duct - ance (µS/cm) 75 76 76 76 77 81 82 82	(stand- ard units) 6.9 7.0 7.0 7.1 9.0 9.0	6.0 6.0 6.0 6.0 6.0 12.0 12.0	dis- solved (mg/L) 14.1 14.1 14.1 14.0 13.0 13.0	(per- cent satur- ation) 113 113 113 113 112 120 120 121	
Date April 1977 04 1515 04 1516 04 1518 04 1519 26 1355 26 1356 26 1357	1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80	con- duct - ance (µS/cm) 75 76 76 76 76 77 81 82 82 82 81	(stand- ard units) 6.9 7.0 7.0 7.1 9.0 9.0 9.2 8.7	6.0 6.0 6.0 6.0 6.0 12.0 12.0 11.5	dis- solved (mg/L) 14.1 14.1 14.1 14.0 13.0 13.0 13.2 12.4	(percent saturation) 113 113 113 113 112 120 120 120 121 113	
Date April 1977 04 1515 04 1516 04 1517 04 1518 04 1519 26 1355 26 1356 26 1357 26 1358	pling depth (ft) 1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60	con- duct - ance (µS/cm) 75 76 76 76 77 81 82 82	(stand- ard units) 6.9 7.0 7.0 7.1 9.0 9.0	6.0 6.0 6.0 6.0 6.0 12.0 12.0	dis- solved (mg/L) 14.1 14.1 14.1 14.0 13.0 13.0	(per- cent satur- ation) 113 113 113 113 112 120 120 121	
Date April 1977 04 1515 04 1516 04 1518 04 1518 04 1519 26 1355 26 1357 26 1358 26 1359 May 1977	1.60 3.30 6.60 9.80 11.6 3.30 6.60 9.80	conduct - ance (µS/cm) 75 76 76 76 77 81 82 82 81 81	(stand-ard units) 6.9 7.0 7.0 7.0 9.0 9.0 9.2 8.7 8.3	6.0 6.0 6.0 6.0 6.0 12.0 11.5 11.0	dis- solved (mg/L) 14.1 14.1 14.1 13.0 13.0 13.2 12.4 12.1	(percent saturation) 113 113 113 113 112 120 120 121 113 110	
Date April 1977 04 1515 04 1516 04 1517 04 1518 04 1519 26 1355 26 1356 26 1357 26 1358 26 1359 May 1977 16 1255	1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 11.6	conduct - ance (µS/cm) 75 76 76 76 77 81 82 82 81 81	(stand-ard units) 6.9 7.0 7.0 7.1 9.0 9.0 9.2 8.7 8.3	ature (°C) 6.0 6.0 6.0 6.0 12.0 11.5 11.0	dis- solved (mg/L) 14.1 14.1 14.1 14.0 13.0 13.0 13.2 12.4 12.1	(percent saturation) 113 113 113 113 112 120 120 120 121 113 110	
Date April 1977 04 1515 04 1516 04 1518 04 1519 26 1355 26 1356 26 1357 26 1358 26 1359 May 1977 16 1255	1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 11.6	conduct - ance (µS/cm) 75 76 76 76 76 77 81 82 82 81 81 86	(stand-ard units) 6.9 7.0 7.0 7.0 7.1 9.0 9.0 9.2 8.7 8.3 8.0 8.2	ature (°C) 6.0 6.0 6.0 6.0 12.0 12.0 11.5 11.0 15.0 13.5	dis- solved (mg/L) 14.1 14.1 14.1 14.0 13.0 13.0 13.2 12.4 12.1	(percent saturation) 113 113 113 113 112 120 120 120 120 120 120 120 100 108 107	
Date April 1977 04 1515 04 1516 04 1518 04 1519 26 1355 26 1356 26 1357 26 1359 May 1977 16 1255 16 1256	1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 11.6	conduct - ance (µS/cm) 75 76 76 76 77 81 82 82 81 81 86 85	(stand-ard units) 6.9 7.0 7.0 7.0 7.1 9.0 9.0 9.2 8.7 8.3 8.0 8.2 7.9	ature (°C) 6.0 6.0 6.0 6.0 12.0 11.5 11.0 15.0 13.5 13.0	dis- solved (mg/L) 14.1 14.1 14.1 14.0 13.0 13.0 13.2 12.4 12.1	(percent saturation) 113 113 113 113 112 120 120 121 113 110 108 107 104	
Date April 1977 04 1515 04 1516 04 1518 04 1518 04 1519 26 1355 26 1356 26 1356 26 1359 May 1977 16 1255 16 1256	1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 11.6	conduct - ance (µS/cm) 75 76 76 76 77 81 82 82 81 81 86 85 85	(stand-ard units) 6.9 7.0 7.0 7.1 9.0 9.0 9.2 8.7 8.3 8.0 8.2 7.9 9.7,7	ature (°C) 6.0 6.0 6.0 6.0 12.0 11.5 11.0 15.0 13.5 13.0 13.0	dis- solved (mg/L) 14.1 14.1 14.1 14.0 13.0 13.0 13.2 12.4 12.1 11.0 11.2 11.0	(percent saturation) 113 113 113 113 112 120 120 121 113 110 108 107 104 103	
Date April 1977 04 1515 04 1516 04 1518 04 1519 26 1355 26 1356 26 1357 26 1358 26 1359 May 1977 16 1255 16 1256 16 1256	1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 11.5	conduct - ance (µS/cm) 75 76 76 76 77 81 82 82 81 81 86 85 85	(stand-ard units) 6.9 7.0 7.0 7.0 7.1 9.0 9.0 9.2 8.7 8.3 8.0 8.2 7.9 7.7 7.4	ature (°C) 6.0 6.0 6.0 6.0 12.0 11.5 11.0 11.5 13.0 13.0 13.0	dis- solved (mg/L) 14.1 14.1 14.1 14.0 13.0 13.0 13.2 12.4 12.1 11.0 11.2 11.0 10.9	(percent saturation) 113 113 113 113 112 120 120 121 113 110 108 107 104 103 101	
Date April 1977 04 1515 04 1516 04 1517 04 1518 04 1519 26 1355 26 1356 26 1359 May 1977 16 1256 16 1257 16 1257 16 1257	1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 11.6	conduct - ance (µS/cm) 75 76 76 76 77 81 82 82 81 81 86 85 85	(stand-ard units) 6.9 7.0 7.0 7.1 9.0 9.0 9.2 8.7 8.3 8.0 8.2 7.9 9.7,7	ature (°C) 6.0 6.0 6.0 6.0 12.0 11.5 11.0 15.0 13.5 13.0 13.0	dis- solved (mg/L) 14.1 14.1 14.1 14.0 13.0 13.0 13.2 12.4 12.1 11.0 11.2 11.0	(percent saturation) 113 113 113 113 112 120 120 121 113 110 108 107 104 103	
Date April 1977 04 1515 04 1516 04 1518 04 1519 26 1355 26 1355 26 1356 26 1359 May 1977 16 1255 16 1257 16 1258 16 1258 16 1258	1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 13.2	conduct - ance (\(\mu S / cm\)) 75 76 76 76 76 781 82 82 81 81 81 86 85 85 85 85 85	(stand-ard units) 6.9 7.0 7.0 7.0 7.0 9.0 9.0 9.2 8.7 8.3 8.0 8.2 7.9 7.7 7.4 7.0	ature (°C) 6.0 6.0 6.0 6.0 12.0 11.5 11.0 15.0 13.5 13.0 13.0 13.0	dis- solved (mg/L) 14.1 14.1 14.1 14.1 14.0 13.0 13.0 13.2 12.4 12.1 11.0 10.9 10.7 9.8	(percent saturation) 113 113 113 113 112 120 120 121 113 110 108 107 104 103 101 93	
Date April 1977 04 1515 04 1516 04 1518 04 1519 26 1355 26 1356 26 1357 26 1359 May 1977 16 1256 16 1256 16 1257 16 1258 16 1259	1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 11.6	conduct - ance (µS/cm) 75 76 76 76 77 81 82 82 81 81 86 85 85 85	(stand-ard units) 6.9 7.0 7.0 7.1 9.0 9.0 9.2 8.7 8.3 8.0 8.2 7.9 7.4 7.0 7.2	ature (°C) 6.0 6.0 6.0 6.0 12.0 11.5 11.0 15.0 13.0 13.0 18.0	dis- solved (mg/L) 14.1 14.1 14.1 14.0 13.0 13.2 12.4 12.1 11.0 11.2 11.0 9.8	(percent saturation) 113 113 113 113 112 120 120 121 113 110 108 107 104 103 101 93	
Date April 1977 04 1515 04 1516 04 1518 04 1518 04 1519 26 1355 26 1356 26 1357 26 1358 26 1359 May 1977 16 1255 16 1256 16 1257 16 1259 16 1259 16 1259 16 1300 June 1977 13 1250	1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 11.5	conduct - ance (µS/cm) 75 76 76 76 77 81 82 82 81 81 86 85 85 85 85	(stand-ard units) 6.9 7.0 7.0 7.0 7.1 9.0 9.0 9.2 8.7 8.3 8.0 8.2 7.9 7.7 4 7.0 7.2 5.9	ature (°C) 6.0 6.0 6.0 12.0 12.0 11.0 11.0 13.5 13.0 13.0 13.0 18.0	dis- solved (mg/L) 14.1 14.1 14.1 14.0 13.0 13.0 13.2 12.4 12.1 11.0 10.9 10.7 9.8 9.8	(percent saturation) 113 113 113 113 112 120 120 121 113 110 108 107 104 103 103 103 103	
Date April 1977 04 1515 04 1516 04 1518 04 1519 26 1355 26 1356 26 1357 26 1359 May 1977 16 1256 16 1256 16 1257 16 1258 16 1259	1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 11.6 1.60 3.30 6.60 9.80 11.6	conduct - ance (µS/cm) 75 76 76 76 77 81 82 82 81 81 86 85 85 85	(stand-ard units) 6.9 7.0 7.0 7.1 9.0 9.0 9.2 8.7 8.3 8.0 8.2 7.9 7.4 7.0 7.2	ature (°C) 6.0 6.0 6.0 6.0 12.0 11.5 11.0 15.0 13.0 13.0 18.0	dis- solved (mg/L) 14.1 14.1 14.1 14.0 13.0 13.2 12.4 12.1 11.0 11.2 11.0 9.8	(percent saturation) 113 113 113 113 112 120 120 121 113 110 108 107 104 103 101 93	

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

						Lake sit	e <u>L5</u>					
	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Trans- par- ency (secchi disk)	0×ygen, dis- solved	Oxygen, dis- solved (per- cent satur-	Bicar- bonate FET-fld (mg/L as	Alka- linity field (mg/L as	Carbon dioxide dis- solved (mg/L	Nitro- gen, NO2+NO3 total (mg/L
Date		(ft)	(µS/cm)	units)	(°C)	(m)_	(mg/L)	at ion)	HC03)	CACO3) a	s CO2)	as N)
April 19	17.7											
04	1525	1.60	78	6.8	6.0	1.40	14.0	112	19	16	4.8	.10
04	1526	3.30	78	6.8	6.0			112				
04	1527	6.60	78	6.8	6.0			112				
04	1528	9.80	78	6.8	6.0			112				
04	1530	13.2	78	6.8	6.0			111	18	15	4.5	.10
04	1531	16.5	77	6.8	6.0			110			7.5	
04	1532	20.0	77	6.8	6.0			109				
04	1533	23.1	77	6.7	6.0			108				
04	1535	25.0	76	6.6	6.0			106	19	16	7.6	.14
26	1415	1.60	80	9.0	12.5	1.50		120	20	16	.0	<.10
26	1416	3.30	80	9.0	12.0			119				
26	1420	4.90							19	16		.01
26	1421	6,60	80	8.7	11.5			113				
26	1425	9.80	79	8.0	11.0			108	20	16	.3	.01
26	1426	13.2	79	7.5	10.0		11.8	104				
26	1427	16.5	79	6.9	9.0			91				
26	1428	20.0	80	6.9	9.0			88				
26	1430	21.5	80	6.9	9.0			87	20	16	4.0	.08
May 1977		20.0	-		3.0			٠.			****	
16	1310	1.60	86	7.9	15.5	1.30	11.0	109	19	16	.4	<.10
16	1312	3.30	85	8.2	14.0			107				
16	1315	6,60	85	8.2	13.5			108	19	16	.2	<.10
16	1319	9.80	85	7.6	13.0			102				
16	1320	11.0							27	22		<.10
16	1322	13.2	85	6.9	13.0		9.7	92				
16	1323	16.5	85	6.7	12.5							
16	1324	20.0	85	6.7	12.5			82				
16	1325	21.5	85	6.6	12.5			79	25	21	10	.00
June 197												
13	1305	1.60	87	6.0	18.5	3.2	9.8	103		17	33	<.10
13	1307	3.30	87	5.6	18.0			103				
13	1308	6.60	87	5.4	18.0							
13	1309	9.80	87	5.3	17.5							
13	1310	13.1	87	5.3	17.5					18	175	<.10
13	1312	16.5	87	5.3	17.0							
13	1315	18.0								18		< .10
13	1317	20.0	87	5.2	15.5							
13	1320	21.3	87	5.3	15.5					20	194	.01
July 197												
11	1145	1.60	94	8.0	25.0	2.40	9.8	117		19	.4	<.10
11	1147	3.30	94	8.0	25.0							
11	1149	6.60	93	8.0	24.0							

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

					Lake site	L5Cont	inued				
	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Trans- par- ency (secchi disk)	Oxygen, dis- solved	Oxygen, dis- solved (per- cent satur-	Alka- linity field (mg/L as	Carbon dioxide dis- solved (mg/L	Nitro- gen, NO2+NO3 total (mg/L
Date		(ft)	(µS/cm)	units)	(°C)	(m)	(mg/L)	ation) (CACO3) a	s CO2)	as N)
											, -
July 197	77										
11	1150	9.80	93	8.0	23.5		9.4	109	20	. 4	.01
11	1153	13.0	92	7.7	22.5		8.4	96			
11	1155	15.0							20		.01
11 11	1157 1200	16.0 18.0	91	7.8	18.0		1.8	19	21		.01
11	1200	20.0	90	7.8	16.5		.0	0			.01
11	1205	21.0	90	7.8	15.5		.0	ŏ	23	.7	.03
August 1											
04	1245	1.60	96	7.4	24.5	3.00	8.9	105	19	1.5	<.10
04 04	1247 1249	3.30 6.60	96 96	7.4 7.3	24.0 24.0		9.0 8.8	106 104			
04	1250	9.80	96	7.1	23.5		8.6	100	19	2.9	<.10
04	1253	13.0	95	7.1	23.0		7.7	89			
04	1255	15.0							19		<.10
04	1258	16.0	94	6.9	21.0		2.5	28	20		- 10
04 04	1300 1302	18.0 20.0	91	7.2	19.0		.0	0	20		<.10
04	1305	21.0	91	7.3	18.0		.0	0	21	2.0	<.10
Septembe											
07	1210	1.60	86		24.0	3.7	8.2	96	20		<.10
07 07	1211 1212	3.30	86 88		24.0 23.5		8.2 8.2	96 95			
07	1215	6.60 9.80	88		23.5		8.0	93	22		<.10
07	1216	13.0	88		23.0		7.9	91			
07	1220	15.0							20		<.10
07	1221	16.0	87		22.0		4.6	52	21		<.10
07 07	1225 1226	18.0 20.0	86		21.0		2.1	23			·, 10
07	1230	21.0	86		21.0		1.0	11	22		<.10
October											_
04	1330	1.60	96		15.5	2.00	8.2	81	22		.02
04 04	1331 1335	3.30 6.60	96 96		15.5 15.5		8.1 8.1	80 80	24		.02
04	1336	9.80	96		15.5		7.9	78			
04	1340	11.5							23		.02
04	1341	13.2	98		15.5		7.9	78			
04 04	1342 1345	16.4 18.0	98 		15.5		7.6	75 	24		.03
04	1346	19.8	98		15.5		6.3	62			.03
04	1350	21.3	98		15.5		.0	0	23		.03
21	1010	1.64	85	7.4	12.0	1.70	7.9	73	21	1.6	. 05
21 21	1011 1015	3.30 6.60	85 85	7.4 7.3	12.0 12.0		7.9 7.9	73 73	22	2.1	.05
21	1015	9.80	85	7.3	12.0		7.8	73		2.1	.03
21	1020	11.5							21		.05
21	1021	13.0	85	7.3	12.0		7.8	72			
21	1025	16.4	85 85	7.2	12.0			70 70	22		.05
21 21	1026 1030	19.8 21.3	85 85	7.2 7.2	12.0 12.0			70 70	21	2.6	.02
Novembe		21.3	03	,,,	12.0		· • /	, 0		2.0	. 02
16	1330	1.60	80	7.3	9.5		4.9	43			
16	1331	3.30	80	7.3	9.5		4.9	43			
16	1332 1333	6.60 9.80	80 80	7.3 7.3	9.0 9.0			41 41			
16 16	1333	13.2	80 80	7.3	9.0			40			
16	1335	16.4	80	7.3	9.0			40			
16	1336	19.8	80	7.3	9.0			41			
April 1			00	7 4		2 22	3.0				
13 13	1300 1305	1.50 9.84	80 82	7.3 7.3	6.0 5.5						
	1303	3.04	02	7.1	5.0						

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

				<u>Lake site</u>	L5Cont	inued			
Date	Time	Sam- pling depth (ft)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)	Phos- phorus, ortho, total (mg/L as P)	Carbon, organic total (mg/L as C)
April 19	77								
04 04 04 04 04 04 04 04 26	1525 1526 1527 1528 1530 1531 1532 1533 1535 1415 1416	1.60 3.30 6.60 9.80 13.2 16.5 20.0 23.1 25.0 1.60 3.30	.080	.52	.70 .70 .84 .62	3.1 3.1 3.7 2.7	.010	<.010 <.010 <.010 <.010	3.0 5.4 5.3
26 26 26 26 26	1420 1421 1425 1426 1427	4.90 6.60 9.80 13.2 16.5	.080	. 50	.59	1.9	.030	<.010 <.010	
26 26 May 1977	1428 1430	20.0 21.5	. 120	.98	1.2	5.2	.100	<.010	
16 16 16 16 16	1310 1312 1315 1319 1320	1.60 3.30 6.60 9.80 11.0	.010	.41	.42	1.9	.020	<.010 <.010 <.010	
16 16 16 16	1322 1323 1324 1325	13.2 16.5 20.0 21.5	.070	.33	.40	1.8	.030	<.010	
June 197 13 13	1305 1307 1308	1.60 3.30 6.60	.040	.21	.25	1.1	.030	<.010 	
13 13 13	1309 1310 1312 1315	9.80 13.1 16.5 18.0	.060	. 24	.30	1.3	.030	<.010 <.010	
13 13 July 197	1317 1320 7	20.0	.060	. 29	.36	1.6	. 030	<.010	
11 11 11	1145 1147 1149	1.60 3.30 6.60	.040	. 57	.61 	2.7	.010	 	

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

				Lake site	L5Cont	inued			
Date	Time	Sam- pling depth (ft)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)	Phos- phorus, ortho, total (mg/L as P)	Carbon, organic total (mg/L as C)
July 19	77								
11	1150	9.80	.030	. 60	.64	2.8	.020		
11	1153	13.0							
11	1155	15.0	.070	. 56	.64	2.8	.030		
11	1157	16.0							
11	1200	18.0	. 130	.36	.50	2.2	.030		
11	1202	20.0							
11	1205	21.0	.180	2.1	2.3	10	.090		
August									
04	1245	1.60	1.60	.97	1.0	4.4	.040		5.9
04	1247	3.30							
04	1249	6.60							
04	1250	9.80	.030	. 95	.98	4.3	.040		8.1
04	1253	13.0 15.0	020		46	2.0	040		3.8
04 04	1255 1258	16.0	.020	. 44	.46	2.0	. 040		
04	1300	18.0	. 050	. 43	.48	2.1	.040	<.010	5.6
04	1302	20.0	.030	.45	.40		.040	1.010	3.0
04	1305	21.0	.010	1.1	1.1	4.9	.070		8.5
Septemb									•.•
07	1210	1.60	.020	. 34	.36	1.6	.020		
07	1211	3.30							
07	1212	6.60							
07	1215	9.80	.020	. 47	.49	2.2	.020		
07	1216	13.0							
07	1220	15.0	.020	. 39	.41	1.8	. 020		
07	1221	16.0							
07	1225	18.0	.040	. 38	.42	1.9	.030		
07 07	1226 1230	20.0 21.0	. 130	. 42	.55	2.4	.080		
October		21.0	, 130	.42	. 55	2,4	.000		
04	1330	1.60	.070	.37	.46	2.0	. 100		
04	1331	3.30	.070				. 100		
04	1335	6.60	.030	. 44	.49	2.2	.060		
04	1336	9.80	.050						
04	1340	11.5	.030	. 39	.44	1.9	.030		
04	1341	13.2							
04	1342	16.4							
04	1345	18.0	.030	. 39	.45	2.0	. 030		
04	1346	19.8							
04	1350	21.3	.020	. 39	. 44	1.9	. 100		
21	1010	1.64	.070	.57	.69	3.1	. 030		
21	1011	3. 3 0							

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

Date	Time	Sam- pling depth (ft)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)
October	1977						
21	1015	6.60	.050	.44	. 54	2.4	.030
21	1016	9.80					
21	1020	11.5	.070	.68	.80	3.5	.030
21	1021	13.0			•••		
21	1025	16.4	.070	.43	. 55	2.4	.030
21	1026	19.8					
21	1030	21.3	.040	.48	. 54	2.4	.030
Novembe	r 1977						
16	1330	1.60					
16	1331	3.30					
16	1332	6.60					
16	1333	9.80					
16	1334	13.2					
16	1335	16.4					
16	1336	19.8					
April 1	978						
13	1300	1.50					.008
13	1305	9.84					.010
13	1310	23.0					.012

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued Lake site L5--Continued

Phytoplankton analyses

() OTF TIME		4•77 525		26•77 200		26,77 415		16,77 310		13•77 305		11.73
DIVERSITY: DIVISION •CLASS		0.8 0.8		0.9		0.9		0.9		0.0		0.2
ORDER		1.5		1.6		1.6		0.9		0.9		0.3
···FAMILY ····GFNUS		2.0		1.6 1.6		1.6 1.6		1.6 1.6		1.6 1.6		0.3 0.6
ORGANISM	CELLS /ML	PER+ CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT
CHLOROPHYTA (GREEN ALGAE)												
.CHLOROPHYCFAECHLOROCOCCALES												
CHARACIACEAESCHROEDERIA		_		_		-		_	*	0	140	1
MICRACTINIACEAE				•				0		_		_
MICRACTINIUMOOCYSTACEAE		-	*	0		-						_
ANKISTRODESMUSCHODATELLA	*	0		0		-	*	0	*	0		-
DICTYOSPHAERIUM		-		-		-		-	*	0		-
NEPHRDCYTIUMSELENASTRUM		-		-		-		0		0		-
•••• TETRAEDRON ••• SCENEDESMACEAE	*	0		-		-	*	0		-		-
CRUCIGENIA		-		-		-		-		0		-
SCENEDESMUSTETRASTRUM	*	0		0		:		0		0		-
••TETRASPORALES •••PALMELLACEAE												
SPHAEROCYSTIS		-		-		-		-		0	700	3
••VOLVOCALES •••CHLAMYDOMONADACEAE												
••••CHLAMYDOMONAS ••ZYGNEMATALES	*	0		0		-		-		-		•
DESMIDIACEAE		_		_		_		_		0		
CHRYSOPHYTA		_								v		
.BACILLARIOPHYCEAECENTRALES												
COSCINODISCACEAE								•		0		_
····CYCLOTELLA ····MELOSIRA	*	25		-	"	33		0	*	0		-
RHIZOSOLENIACEAERHIZOSOLENIA		0				-		-		-		-
PENNALES		Ü										
ACHNANTHACEAECOCCONEIS		-	•	0		-		-		-		-
CYMBELLACEAEEPITHEMIA		-		_		_		_		-		-
FRAGILARIACEAE	*	25	#	33	#	33	,	3 3		0		
••••ASTERIONELLA ••••FRAGILARIA	"	-		-	"	-		-		0		-
SYNEDRAGOMPHDNEMATACEAE		•		0		•	*	0		-		-
GOMPHONEMA	*	25		-		-		-		-		-
NITZSCHIA	٠	0		-		-		0		-		-
TABELLARIACEAE		-		-		-		-	٠	0		-
.CHRYSOPHYCEAECHRYSOMONADALES												
OCHROMONADACEAE				•				_		_		_
DINOBRYDN OCHROMONAS		0		<u> </u>		-		-		0		-
CYANDPHYTA (BLUE-GREEN ALGAE)												
.CYANOPHYCEAECHROCCOCCALES												
CHROCCOCCAEAEANACYSTIS		-		-		-		-		0	1600	6
HORMOGONALES NOSTOCACEAE												
ANABAENA		-		-		-		-	,	# 3 3	280	1
APHANIZOMENONOSCILLATORIACEAE		-	•	0		-		¥ 33				-
OSCILLATORIACHROCCOCCALES	,	2 5	,	¥ 33	*	¥ 33	•	¥ 33	1	# 3 3		•
CHROCCOCCAEAE						_		_		# 33	24000#	. 00
••••GOMPHOSPHAERIA EUGLENOPHYTA (EUGLENOIDS)		-	•	* 33		-	-	_		. 33	24000	70
•CRYPTOPHYCEAE ••CRYPTOMONIDALES												
• • • CRYPTOMONODACEAE	_					_		_		0		_
CRYPTOMONAS .FUGLENOPHYCEAE	*	0	•	0		_		-	-	•	- -	
EUGLENALESEUGLENACEAE												
· · · · EUGLENA		-		-		-		-		0		-
PHACUSTRACHELOMONAS		-		-		-		0		0		-
PYRRHOPHYTA (FIRE ALGAE)												
.DINOPHYCEAEPERIDINIALES												
CERATIACEAECERATIUM		-		-		-		-		-		-
PERIDINIACEAE PERIDINIUM		-		0		-		-		-		_
* - Present in sample but not a domin	ant type			-								

^{* =} Present in sample but not a dominant type.
= A dominant type in sample.

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued $Lake\ site\ L5--Continued$

			Phytoj	lankton	analyses					
DATE TIME	AtiG	4,77 245		17•77 130	SEP	7•77 1210	0C1 1	4•77 330		21•77 010
DIVERSITY: DIVISION .CLASS .ORDERFAMILYGENUS		1.8 1.8 2.3 2.6 2.6		0.8 0.8 1.5 1.5		0.0 0.0 1.0 1.0 2.0		1.0 1.5 1.9 2.3 2.6		1.6 1.6 1.6 1.6
ORGANISM	CFLLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT
CHLOPOPHYTA (GREEN ALGAE) •CHLOROPHYCFAF										
CHLOROCOCCALES										
CHARACIACEAE SCHROFDFRIA		-		-		-		-		-
MICRACTINIACEAEMICRACTINIUM		-		-		-		-		-
OOCYSTACEAEANKISTRODESMUS	•	0		-		-		-		-
····CHODATELLA ····DICTYOSPHAERIUM		-		-		-		-		-
NEPHROCYTIUM SELENASTRUM		-		-		-		-		-
TETRAEDRONSCENEDESMACEAE		-		-		-		-		-
CRUCIGENIA SCENEDESMUS	*	0 0		-		-		-		-
••••TETRASTRUM ••TETRASPORALES		-		-		-		-		-
•••PALMELLACEAE ••••SPHAEROCYSTIS		17		-		_		_		_
VOLVOCALFSCHLAMYDOMONADACEAE										
CHLAMYDOMONASZYGNEMATALES		-		-		-		-		-
DESMIDIACEAE STAURASTRUM		0		_		_		_		_
CHRYSOPHYTA		v								
-BACILLARIOPHYCEAE										
CENTRALES COSCINODISCACEAE										
····CYCLOTELLA ····MELOSIRA		-		-		-		-	#	33
RHIZOSOLENIACEAERHIZOSOLENIA		-		-		-		-		-
••PENNALES •••ACHNANTHACEAE										
COCCONEISCYMBELLACEAE		-		-		-		-		-
EPITHEMIAFRAGILARIACEAE		0		-		-		-		-
····ASTERIONELLA ····FRAGILARIA		-		-		-	#	17		-
••••SYNEDRA •••GOMPHONEMATACEAE		-		-		-		-		-
GOMPHONEMÁNITZSCHIACEAE		-		-		-		-		-
••••NITZSCHIA •••TABELLARIACEAE		-		-		-		-		-
TABELLARIA .CHRYSOPHYCEAE		-		-		-	#	17		-
CHRYSOMONADALES OCHROMONADACEAE										
DINOBRYON OCHROMONAS		-		-		-	#	17		-
		-		-		-		-		-
CYANOPHYTA (BLUE-GREEN ALGAE) •CYANOPHYCEAE										
••CHROCCOCCALES •••CHROCCOCCAEAE										
ANACYSTISHORMOGONALES		-		-	#	25	*	17		-
ANABAENA		.=	#			25				-
••••APHANIZOMENON •••OSCILLATORIACEAE		17	#	25	*	25	#	17	#	33
OSCILLATORIACHROCCOCCALES	#	17		-		-		-		-
CHROCCOCCAEAEGOMPHOSPHAERIA	#	17	#	25	*	25	#	17		-
EUGLENOPHYTA (EUGLENOIDS) •CRYPTOPHYCEAE										
••CRYPTOMONIOALES •••CRYPTOMONODACEAE										
••••CRYPTOMONAS •FUGLENOPHYCEAE		-		-		-		-		-
EUGLENALES EUGLENACEAF										
EUGLENA PHACUS		-		-		-		-		-
TRACHELOMONAS	#	17	#	25		-		-	#	33
PYRRHOPHYTA (FIRE ALGAE) • PINOPHYCEAE										
PERIDINIALESCERATIACEAE										
CERATIUMPERIDINIACEAE	#	17		-		-		-		-
PERIDINIUM		-		-		-		-		-
<pre>* = Present in sample but not a domina # = A dominant type in sample.</pre>	int type.									

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

			<u>Lake</u>	site L6				
	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper-	Oxygen, dis- solved	Oxygen, dis- solved (per- cent satur-	
Date		(ft)	(µS/cm)	units)	(°C)	(mg/L)	ation)	
			(ps/ cm/	4117037		1-9/-/	a c t o ii j	
April 19		1 60	77	6 7	• 0	14.1	112	
04 04	1555 1556	1.60 3.30	77 77	6.7 6.7	6.0 6.0	14.1 14.1	113 113	
04	1557	6.60	77	6.7	6.0	14.0	112	
04	1558	9.80	77	6.6	6.0	13.8	110	
04	1559	13.2	78	6.6	6.0	13.8	110	
04	1600	16.5	78	6.6	6.0	13.7	110	
04	1601	18.1	77	6.6	6.0	13.7	110	
26	1445	1.60	77	8.9	12.0	12.7	118	
26	1446	3.30	75	8.8	11.5	12.4	114	
26	1447	6.60	76	8.6	11.5	12.3	113	
26	1448	9.80	78	8.2	11.0	12.0	109	
26	1449	13.2	77	7.3	10.0	11.3	100	
26	1450	16.5	77	7.2	9.5	10.4	90	
May 1977								
16	1335	1.60	87	8.1	15.0	11.0	108	
16	1336	3.30	86	8.3	13.5	11.3	108	
16	1337	6.60	86	8.0	13.0	10.5	99	
16	1338	9.80	86	7.0	13.0	10.0	94	
16	1339	13.2	86	6.9	13.0	9.8	92	
16	1340	16.5	86	6.8	12.5	9.4	88	
16	1341	20.0	86	6.8	12.5	9.3	87	
June 197	7							
13	1340	1.60	88	7.1	18.5	9.7	103	
13	1342	3.30	87	6.2	18.5	9.7	103	
13	1344	6.60	87	5.8	18.5	9.7	103	
13	1346	9.80	87	5.5	18.0	9.8	103	
13	1348	13.2	87	5.6	18.0	9.8	103	
13	1350	15.0	87	5.5	17.5	9.6	100	
			Lake	site L7				
			Spe-				Oxygen, dis-	
			cific				so I v ed	
		Sam-	con-	pН		Oxygen,	(per-	
		pling	duct-	(stand-	Temper-	dis-	cent	
	Time	depth	ance	ard	ature	solved	satur-	
Date		(ft)	(µS/cm)	units)	(°c)	(mg/L)	ation)	
	17.7							
April 19	1120	1 60	79	7.7	E 0	12 0	109	
ሰፍ		1.60			5.0	13.9		
06			70	77	E 0	120		
06	1121	3.30	79	7.7	5.0	13.8		
06 06	1122	6.60	80	7.7	5.0	13.8	108	
06 06 06	1122 1123	6.60 9.80	80 79	7.7	5.0 5.0	13.8 13.8	108 108	
06 06 06	1122 1123 1124	6.60 9.80 11.0	80 79 80	7.7 7.7 7.7	5.0 5.0 5.0	13.8 13.8 13.8	108 108 108	
06 06 06 27	1122 1123 1124 1315	6.60 9.80 11.0 1.60	80 79 80 81	7.7 7.7 7.7 8.7	5.0 5.0 5.0 11.5	13.8 13.8 13.8 12.6	108 108 108 116	
06 06 06 27 27	1122 1123 1124 1315 1316	6.60 9.80 11.0 1.60 3.30	80 79 80 81 81	7.7 7.7 7.7 8.7 8.6	5.0 5.0 5.0 11.5 11.5	13.8 13.8 13.8 12.6 12.6	108 108 108 116 116	
06 06 06 27 27	1122 1123 1124 1315 1316 1317	6.60 9.80 11.0 1.60 3.30 6.60	80 79 80 81 81 81	7.7 7.7 7.7 8.7 8.6 8.5	5.0 5.0 5.0 11.5 11.5	13.8 13.8 13.8 12.6 12.6	108 108 108 116 116	
06 06 06 27 27 27 27	1122 1123 1124 1315 1316 1317 1318	6.60 9.80 11.0 1.60 3.30	80 79 80 81 81	7.7 7.7 7.7 8.7 8.6	5.0 5.0 5.0 11.5 11.5	13.8 13.8 13.8 12.6 12.6	108 108 108 116 116	
06 06 06 27 27 27 47 May 1973	1122 1123 1124 1315 1316 1317 1318	6.60 9.80 11.0 1.60 3.30 6.60 9.80	80 79 80 81 81 81	7.7 7.7 7.7 8.7 8.6 8.5 8.4	5.0 5.0 11.5 11.5 11.5	13.8 13.8 12.6 12.6 12.4 12.1	108 108 108 116 116 114 110	
06 06 06 27 27 27 May 197;	1122 1123 1124 1315 1316 1317 1318	6.60 9.80 11.0 1.60 3.30 6.60 9.80	80 79 80 81 81 81 82	7.7 7.7 7.7 8.7 8.6 8.5 8.4	5.0 5.0 5.0 11.5 11.5 11.5 11.0	13.8 13.8 13.8 12.6 12.6 12.4 12.1	108 108 108 116 116 114 110	
06 06 06 27 27 27 May 197 16	1122 1123 1124 1315 1316 1317 1318 7	6.60 9.80 11.0 1.60 3.30 6.60 9.80	80 79 80 81 81 81 82 88	7.7 7.7 7.7 8.6 8.5 8.4 8.5	5.0 5.0 5.0 11.5 11.5 11.0 14.5	13.8 13.8 13.8 12.6 12.6 12.4 12.1	108 108 108 116 116 114 110	
06 06 06 27 27 27 47 48y 197; 16	1122 1123 1124 1315 1316 1317 1318 7 1355 1356 1357	6.60 9.80 11.0 1.60 3.30 6.60 9.80 1.60 3.30 6.60	80 79 80 81 81 81 82 88 87	7.7 7.7 7.7 8.7 8.6 8.5 8.4 8.5 8.6	5.0 5.0 5.0 11.5 11.5 11.0 14.5 14.0	13.8 13.8 13.8 12.6 12.6 12.4 12.1 11.4 11.5	108 108 108 116 116 114 110	
06 06 06 27 27 27 27 May 1977 16 16	1122 1123 1124 1315 1316 1317 1318 7 1355 1356 1357 1358	6.60 9.80 11.0 1.60 3.30 6.60 9.80	80 79 80 81 81 81 82 88	7.7 7.7 7.7 8.6 8.5 8.4 8.5	5.0 5.0 5.0 11.5 11.5 11.0 14.5	13.8 13.8 13.8 12.6 12.6 12.4 12.1 11.4 11.5	108 108 108 116 116 114 110	
06 06 06 27 27 27 May 197: 16 16 16 June 19;	1122 1123 1124 1315 1316 1317 1318 7 1355 1356 1357 1358	6.60 9.80 11.0 1.60 3.30 6.60 9.80 1.60 3.30 6.60 9.80	80 79 80 81 81 81 82 88 87 86	7.7 7.7 8.7 8.6 8.5 8.4 8.5 8.6 8.2 7.6	5.0 5.0 5.0 11.5 11.5 11.0 14.5 14.0 13.5	13.8 13.8 12.6 12.6 12.4 12.1 11.4 11.5	108 108 108 116 116 114 110 111 111 107	
06 06 06 06 27 27 27 27 May 197; 16 16 16 June 19;	1122 1123 1124 1315 1316 1317 1318 7 1355 1356 1357 1358 77	6.60 9.80 11.0 1.60 3.30 6.60 9.80 1.60 3.30 6.60 9.80	80 79 80 81 81 81 82 88 87 86	7.7 7.7 7.7 8.6 8.5 8.4 8.5 8.6 8.2 7.6	5.0 5.0 5.0 11.5 11.5 11.0 14.5 14.0 13.5 13.0	13.8 13.8 13.8 12.6 12.4 12.1 11.4 11.5 11.2 10.7	108 108 108 116 116 114 110 111 111 107 101	
06 06 06 27 27 27 27 48ay 197; 16 16 16 13	1122 1123 1124 1315 1316 1317 1318 7 1355 1356 1357 1358 77	6.60 9.80 11.0 1.60 3.30 6.60 9.80 1.60 3.30 6.60 9.80	80 79 80 81 81 81 82 88 87 86 86	7.7 7.7 7.7 8.7 8.6 8.5 8.4 8.5 8.6 8.2 7.6	5.0 5.0 11.5 11.5 11.5 11.0 14.5 13.5 13.0	13.8 13.8 13.8 12.6 12.6 12.4 12.1 11.4 11.5 11.2 10.7	108 108 108 116 116 114 110 111 111 107 101	
06 06 06 27 27 27 27 May 197 16 16 16 13 3	1122 1123 1124 1315 1316 1317 1318 7 1355 1356 1357 1358 77 1358	6.60 9.80 11.0 1.60 3.30 6.60 9.80 1.60 3.30 6.60 9.80	80 79 80 81 81 82 88 87 86 86 87	7.7 7.7 7.7 8.6 8.5 8.4 8.5 8.6 8.2 7.6 7.4 6.6 5.9	5.0 5.0 5.0 11.5 11.5 11.0 14.0 13.5 13.0 18.0 17.5	13.8 13.8 13.8 12.6 12.6 12.4 12.1 11.4 11.5 11.2 10.7 9.9 9.9	108 108 108 116 116 114 110 111 111 107 101	
06 06 06 27 27 27 27 4ay 197; 16 16 16 3une 19; 13	1122 1123 1124 1315 1316 1317 1318 7 1355 1356 1357 1358 77	6.60 9.80 11.0 1.60 3.30 6.60 9.80 1.60 3.30 6.60 9.80	80 79 80 81 81 81 82 88 87 86 86	7.7 7.7 7.7 8.7 8.6 8.5 8.4 8.5 8.6 8.2 7.6	5.0 5.0 11.5 11.5 11.5 11.0 14.5 13.5 13.0	13.8 13.8 13.8 12.6 12.6 12.4 12.1 11.4 11.5 11.2 10.7 9.9 9.9	108 108 108 116 116 114 110 111 111 107 101	

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

						Lake site	e L8					
	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Trans- par- ency (secchi disk)	Oxygen, dis- solved	Oxygen, dis- solved (per- cent satur-	Bicar- bonate FET-fld (mg/L as	Alka- linity field (mg/L as	Carbon dioxide dis- solved (mg/L	Nitro- gen, NO2+NO3 total (mg/L
Date		(ft)	(µS/cm)	units)	(°C)	(m)	(mg/L)	at ion)	HC03)	CACO3) a	s CO2)	as N)
April 19	17.7											
06	1140	1.60	76	7.7	5.0	1.30	13.9	109	19	16	.6	.17
06	1141	3.30	77	7.7	5.0		13.8	108				
06	1142	6.60	77	7.7	5.0		13.9	109				
06	1143	9.80	77	7.6	5.0		13.8	108				
06	1144	13.2	77	7.6	5.0		13.9	109				
06	1145	14.0							19	16		.17
06	1146	16.5	77	7.6	5.0		13.B	108				
06	1147	20.0	78	7.6	5.0		13.8	108				
06	1148	23.1	78	7.6	5.0		13.8	108				
06	1149	26.4	78	7.6	5.0		13.8	108				
06	1150	28.9	78	7.5	5.0		13.8	108	18	15	.9	.20
27	1340	1.60	81	8.7	12.0	1.50	12.6	117	20	16	.0	<.10
27	1342	3.30	81	8.7	11.5		12.5	115				
27	1345	4.90							20	16		<.10
27	1346	6.60	81	8.6	11.5		12.3	113				
27	1347	9.80	80	8.4	11.0		12.0	109				
27	1350	11.5							24	20		.01
27	1351	13.2	79	7.6	10.0		12.5	102				
27	1352	16.5	79	7.1	9.5		12.2	97				
27	1353	20.0	80	6.6	9.0		9.6					
27	1354	23.1	80	6.6	9.0		9.4	81				
27	1355	26.4	80	6.9	9.0		9.2		19	16	3.8	.08
May 1977												
04	1305	27.0										
16	1415	1.60	88	8.4	15.0	1.00		111	19	16	.1	<.10
16	1416	3.30	88	8.4	14.5		11.4	111				
16	1417	6.60	88	8.4	14.0		11.4	111				
16	1420	8.20							26	21		<.10
16	1421	9.80	87	8.5	13.5		11.4	109				
16	1425	13.2	87	8.0	13.0		11.0	104	19	16	.3	< .10
16	1426	16.5	87	7.2	13.0		10.4	98				
16	1427	20.0	87	7.2	13.0		10.3	97				
16	1428	23.1	87	6.9	12.5		9.8	92				
16	1430	26.4	87	6.6	12.0		8.0	74	24	20	9.6	.02
June 197												
13	1415	1.60	89	7.6	18.0	3.00	10.0	105		16	.8	.00
13	1417	3.30	89	7.5	18.0		10.0					
13	1418	6.60	89	6.0	18.0		10.1	106				
13	1420	9.80	88	5.9	16.5		9.5	97		17	41	.02
13	1421	13.2	88	5.6	16.0		8.7	88				
13	1423	16.5	88	5.7	16.0		8.4					

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

					Lake site	EUCOIN	. mueu				
	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Trans- par- ency (secchi disk)	Oxygen, dis- solved	Oxygen, dis- solved (per- cent satur-	Alka- linity field (mg/L as	Carbon dioxide dis- solved (mg/L	Nitro- gen, NO2+NO3 total (mg/L
Date		(ft)	(µS/cm)	units)	(°C)	(m)	(mg/L)	at ion)	CACO3) a	s CO2)	as N)
June 197	77										
13	1425	18.0							17		.04
13 13	1426 1430	20.0 21.3	88 89	5.7 5.7	15.0 14.0		4.7	46 4	18	70	. 01
15	1430	21.5	03	3.7	14.0		.,	-	10	70	.01
July 197 11	77 1225	1,60	96	8.0	25.0	2.40	9.8	117	19	. 4	<.10
11	1227	3.30	96	8.0	25.0		10.0	119			
11	1229	6.60	96	8.1	24.5		10.1	120			
11 11	1230 1233	9.80 13.0	95 93	8.0 7.8	24.5 22.0		9.6 8.4	11 <i>4</i> 95	19	. 4	<.10
11	1235	15.0							19		<.10
11	1237	16.0 20.0	91 90	7.9	18.5		1.7	18 0			
11 11	1239 1240	20.0 21.0	90	7.9	16.0		.0		20		<.10
11	1243	23.0	88	7.9	15.0		.0	0			
11 August 1	1245	26.0	105	8.0	14.0		.0	0	25	. 5	<.10
04	1320	1.60	98	7.6	25.0	3.00	8.9	106	19	.9	<.10
04	1322	3.30	97	7.5	24.0		9.1	107			
04 04	1324 1325	6.60 9.80	97 97	7.3 7.3	24.0 23.5		9.1 9.0	107 95	18	1.7	<.10
04	1327	13.0	97	7.3	23.0		8.2	94			
04	1330	16.0	95	7.3	21.5		5.6	63	20	1.9	<.10
04 04	1332 1335	20.0 23.0	91 94	7.4 7.4	18.5 16.0		.7 .0	7 0	23	1.8	<.10
04	1340	26.0	116	7.5	14.5		.0	0	28	1.7	<.10
Septembe		1 60	0.7		22.5	2.6	0.4	00	20		00
07 07	1245 1246	1.60 3.30	87 87		23.5 23.5	3.6	8.4 8.4	98 98	20		.00
07	1247	6.60	87		23.0		8.4	97			
07	1250	9.80	87 88		23.0 22.5		8.2 7.7	94 88	20		.00
07 07	1251 1255	13.0 16.0	86		21.0		4.4	49	21		.00
07	1256	20.0	90		19.5		.6	6			
07 07	1300 1301	21.0 23.0	99		17.0		.0	0	24		<. 10
07	1305	25.0	124		15.0		.0	ő	3 7		.00
October		1.60	0.7		16.0	2 10	7 1	72	25		03
04 04	1 40 5 1 40 6	1.60 3.30	97 97		16.0 16.0	2.10	7.1 7.2	73	25		.03
04	1410	6.60	97		16.0		7.2	73	23		.04
04 04	1411 1415	9.80 11.5	97		16.0		7.1	72	 2 5		
04	1416	13.2	99		15.0		7.2	71			
04	1417	16.4	99		15.5		7.2	71			
04 04	1420 1421	19.8 23.0	99 99		15.5 15.5		7.2 7.0	71 69	24		.03
04	1425	24.7	99		15.5		7.2	71			. 03
21	1040	1.60	90 90	7.2				70 7 0	22	2.7	.05
21 21	1041 1045	3.30 6.60	90	7.2 7.2	12.5			69	22	2.7	.06
21	1046	9.80	90	7.1	12.0		7.2	66			
21 21	1050 1051	13.1 13.2	90 90	7.1 7.0	12.0 12.0			66 65	22	3.4	.06
21	1051	20.0	90	7.0				64			
21	1055	21.3		7.0	12.0				23		.06
21 21	10 56 1100	23.0 24.7	90 90	7.0 7.1	12.0 12.0			63 61	22	3.4	.07
Novembe	r 1977										
16	1340	1.60	85 85	7.3 7.3	10.0 9.5		_	43 42			
16 16	1341 1342	3.30 6.60	85	7.3				42			
16	1343	9.80	84	7.3	9.0		4.8	41			
16 16	1344 1345	13.2 16.0	80 80	7.2 7.2				41 40			
16	1346	19.8	80	7.2				40			
16	1347	23.0	80	7.2				38			
April 1 13	978 1240	3.28	85	7.0	5.5	2.10	7.1				
13	1245	13.1	85	6.9	5.0		6.8				
13	1250	26.2	85	6.9	5.0		6.5				

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

					site L8		-				
Date	Time	Sam- pling depth (ft)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)	Phos- phorus, ortho, total (mg/L as P)	Carbon, organic total (mg/L as C)	Algal growth poten- tial, bottle test (mg/L)	
April 1	077										
06	1140	1.60	. 050	. 55	.77	3.4	.010	<.010	3.0	.7	
06	1141	3.30	.050				.010				
06	1142	6.60									
06	11 43	9.80									
06	1144	13.2									
06	1145	14.0	.080	. 32	.57	2.5	.010	<.010	2.8		
06	1146	16.5									
06	1147	20.0									
06	1148	23.1									
06	1149	26.4									
06	1150	28.9	.060	. 54	.80	3.5	.020	<.010	3.6		
27	1340	1.60	.050	. 37	.42	1.9	.030	<.010			
27	1342	3.30									
27	1345	4.90	.060	. 61	.67	3.0	.030	<.010			
27	1346	6.60									
27	1347	9.80									
27	1350	11.5	.050	. 46	.52	2.3	.020	<.010			
27	1351	13.2									
27	1352	16.5									
27	1353	20.0									
27	1354	23.1									
27	1355	26.4	.090	1.4	1.6	7.0	.170	<.010			
May 197	7										
04	1305	27.0									
16	1415	1.60	. 040	. 33	.37	1.6	.030	<.010			
16	1416	3.30									
16	1417	6.60									
16	1420	8.20	.040	.27	.31	1.4	.030	<.010			
16	1421	9.80									
16	1425	13.2	.050	. 30	.35	1.6	. 030	<.010			
16	1426	16.5									
16	1427	20.0									
16	1428	23.1									
16	1430	26.4	.070	. 44	.53	2.3	. 090	<.010			
June 19											
13	1415	1.60	.010	. 32	.34	1.5	.015	<.010			
13	1417	3.30									
13	1418	6.60		20	26	1.6	020	- 010			
13	1420	9.80	.050	. 29	.36	1.6	.030	<.010			
13	1421	13.2									

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

				Lake s	ite L8	Cont inued	Ĺ				
Date	Time	Sam- pling depth (ft) as	Nitro- gen, ammonia dis- solved (mg/L N) as	Nitro- gen, organic total (mg/L N) as	Nitro- gen, total (mg/L N) as	Nitro- gen, total (mg/L NO3) as	Phos- phorus, total (mg/L : P) as	Phos- phorus, ortho, total (mg/L P) as	Carbon, organic total (mg/L C) (mg	Algal growth poten- tial, bottle test	
June 197	,										
13	1425	18.0	.050	. 32	.41	1.8	.030	<.010			
13	1426	20.0									
13	1430	21.3	.050	. 45	.51	2.3	.030	<.010			
July 197											
11	1225	1.60	.040	. 49	.53	2.3	.020				
11	1227	3.30									
11	1229	6.60	**								
11	1230	9.80	.060	. 50	.56	2.5	.020				
11	1233	13.0	050	20	42	1 0	020				
11	1235 1237	15.0 16.0	.050	. 38	.43	1.9	.030				
11 11	1239	20.0									
11	1240	21.0	.070	1.3	1.4	6.2	.040				
11	1243	23.0									
11	1245	26.0	.300	2.0	2.3	10	.100				
August 1											
04	1320	1.60	.010	. 52	.53	2.3	.020		4.9	.7	
04	1322	3.30									
04	1324	6.60									
04	1325	9.80	.010	. 49	.50	2.2	. 020		3.9		
04	1327	13.0									
04	1330	16.0	.020	. 38	.40	1.8	.030		9.4		
04 04	1332 1335	20.0 23.0	.070	.60	.67	3.0	.070		8.9		
04	1340	26.0	.190	.80	.99	4.4	.130		6.7		
Septembe		20.0	.130	.00	.55	7.7	. 130		0.7		
07	1245	1.60	.050	. 30	.36	1.6	.020				
07	1246	3.30									
07	1247	6.60									
07	1250	9.80	.040	. 41	.46	2.0	.030				
07	1251	13.0									
07	1255	16.0	.030	. 38	.42	1.9	.030				
07	1256	20.0									
07	1300	21.0	.170	. 39	.56	2.5	.050				
07	1301	23.0									
07	1305	25.0	.890	.31	1.2	5.4	. 240				
October	1405	1 60	.130	27	E2	2 2	.100				
04 04	1405	1.60 3.30	. 130	.37	.53	2.3	. 100				
04	1410	6.60	. 130	.37	.54	2.4	.080				
04	1411	9.80	. 130	.57	.54		.000				
04	1415	11.5									
04	1416	13.2									

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

					L8Cont	mueu		
	Date	Time	Sam- pling depth (ft)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)
,	ctober	1077						
	04	1417	16.4					
	04	1420	19.8	.130	.38	. 54	2.4	.110
	04	1421	23.0	.130	.50	. 54	2,4	.110
	04	1425	24.7	.130	.41	. 57	2.5	.090
	21	1040	1.60	.140	.43	,62	2.7	.040
	21	1041	3.30					
	21	1045	6.60	.140	.48	. 68	3.0	.030
	21	1046	9.80					
	21	1050	13.1	.140	.55	. 75	3.3	.040
	21	1051	13.2					
	21	1052	20.0					
	21	1055	21.3	.140	.50	. 70	3.1	.030
	21	1056	23.0					
	21	1100	24.7	.140	.79	1.0	4.4	.030
	lovember							
	16	1340	1.60					
	16	1341	3.30					
	16	1342	6.60					
	16	1343	9.80					
	16	1344	13.2					
	16	1345	16.0					
	16	1346	19.8					
	16	1347	23.0					
	pril 19							
	13	1240	3.28					.016
	13	1245	13.1					.015
	13	1250	26.2					.013

Lake site L8--Continued

DATE TIME	APR 6+77 1140	APR 27,77 1200	APR 27,77 1340	MAY 16+77 1415	JUN 13+77 1415	JUL 1 . 17 1225
DIVEPSITY: DIVISION •CLASS ••ORDER ••FAMILY	0.9 0.9 1.6 1.6	0.9 0.9 0.9 1.6	0.9 0.9 1.6 1.6	0.9 0.9 0.9 1.6	1.5 1.5 1.5 1.5	0.5 0.5 1.1 1.7
· · · · GENUS	1.6	1.6	1.6	1.6	5.0	1.8
ORGANISM	CELLS PFR- /ML CENT	CELLS PER- /ML CENT				
CHLOROPHYTA (GREEN ALGAE) •CHLOROPHYCEAE						
CHLOROCOCCALESCHARACIACEAE						
SCHROEDFRIACOELASTRACEAE						* ()
COELASTRUMMICRACTINIACFAE		* 0		* 0	* 0	
MICRACTINIUMOOCYSTACEAE		* 0			* 0	
ANKISTRODESMUSCHODATELLADICTYOSPHAERIUM		* 0		* 0		 670 7
TETRAEDRONSCENEDESMACEAE		* 0		* 0		
CRUCIGENIASCENEDESMUS		* 0		 * 0		91 1 # 0
••TETRASPORALES •••PALMELLACEAE						
SPHAEROCYSTISVOLVOCALES				* 0		
CHLAMYOOMONADACEAECHLAMYDDMONAS		* 0				
ZYGNEMATALESDESMIDIACEAECLOSTERIUM				* 0	<u></u> -	
···STAURASTRUM					a 0	* 0
CHRYSOPHYTA .BACILLARIOPHYCEAE						
CENTRALESCOSCINODISCACEAE						
····CYCLOTELLA ····MELOSIRA	# 33	* 0	# 33	* 0		
STEPHANODISCUSPENNALES		* 0		* 0		
ACHNANTHACEAEACHNANTHES		* 0				
FRAGILARIACEAEASTERIDNELLA	# 33	* 33	# 33	# 33 * 0	# 25 # 25	
FRAGILARIASYNEDRAGOMPHONEMATACEAE				* 0		
GOMPHONEMANITZSCHIACEAE				• 0		
NITZSCHIATABELLARIACEAE		* 0				
TABELLARIA .CHRYSOPHYCEAE				* 0		
CHRYSOMONADALESOCHROMONADACEAE				* n		
DINOBRYON OCHROMONAS		* 0		* 0		
CYANOPHYTA (BLUE-GREEN ALGAE) •CYANOPHYCEAE						
CHROCCOCCALESCHROCCOCCAEAE						
AGMENELLUM ANACYSTIS						
HORMOGONALES NOSTOCACEAE						
ANABAENA APHANIZOMENON	== =	# 33	# 33	# 33	# 25 	1500# 16
OSCILLATORIACEAEARTHROSPIRA						 5600# 58
OSCILLATORIAPHORMIMIDIUM		# 33		# 33		* 0
CHROCCOCCALESCHROCCOCCAEAEGOMPHOSPHAERIA					* 0	1600# 17
EUGLENOPHYTA (EUGLENOIDS) .CRYPTOPHYCEAE						
CRYPTOMONIDALES CRYPTOCHPYSIDACEAE						
CHROOMONAS CRYPTOMONOOACEAE		* 0				
CRYPTOMONAS .FUGLENOPHYCEAE		* 0		* 0		
EUGLENALESEUGLENACEAE					# 25	۰ 0
TRACHFLOMONAS		* 0			# 25	- 1)
PYRRHOPHYTA (FIRE ALGAE) OINOPHYCEAE OPERIOINIALES						
CERATIUM						* 0
GLENODINIACEAF				* 0		
PERIDINIACEAEPERIDINIUM		* 0		a 0		
<pre>* = Present in sample but not a domin # = A dominant type in sample.</pre>	ant type					
\$ P - 10 - 1		70				

⁷⁰

Table 11.--Chemical, physical, and biological chara teristics of lake water--Continued $Lake\ site\ L8--Continued$

		Phytoplankton			
DATE TIME	AUG 4.7 1320		SEP 7.77 1245	OCT 4.77 1405	OCT 21.77
DIVERSITY: DIVISION	1.3	0.8	0.0	1.0	0.8
•CLASS	1.3	0.8 1.5	0.0	1.5	0 • ⊭
FAMILY	1.3 1.9	1.5	1.0 1.4	2.3 2.3	1.5 1.5
····GFNUS	2.6	2.0	2.3	2.3	2.0
ORGANISM	CELLS PER		CELLS PER-	CELLS PER- /ML CENT	CELLS PER-
CHLOROPHYTA (GRFEN ALGAE)					
.CHLOROPHYCFAE CHLOROCOCCALES					
CHARACIACEAE					
••••SCHROEDERIA •••COELASTRACEAE					
••••COELASTRUM •••MICRACTINIACEAE					
MICRACTINIUM			- - -		
•••OOCYSTACEAE •••ANKISTRONESMUS	* 0				
••••CHODATELLA ••••DICTYOSPHAERIUM					
· · · · SCENEDESMACEAE					
CRUCIGENIA					
••••SCENEDESMUS ••TETRASPORALES					-
•••PALMELLACEAE ••••SPHAEROCYSTIS	# 17	# 25			
VOLVOCALESCHLAMYDOMONADACEAE	- "				
CHLAMYDOMONAS					
••ZYGNEMATALES •••DESMIDIACEAE					
····CLOSTERIUM ····STAURASTRUM	* 0				
CHRYSOPHYTA BACILLARIOPHYCEAE					
••CENTRALES •••COSCINODISCACEAE					
····CYCLDTELLA ····MELOSIRA	* 0			# 20	 # 25
STEPHANODISCUS				" -	
••PENNALES •••ACHNANTHACEAE					
••••ACHNANTHES •••FRAGILARIACEAE					
ASTERIONELLA FRAGILARIA				# 20	# 25
SYNEDRA	== =		== =		# 25
GOMPHONEMATACEAEGOMPHONEMA					
NITZSCHIACEAE					
TABELLARIACEAE TABELLARIA					
.CHRYSOPHYCEAE					
CHRYSOMONADALESOCHROMONADACEAE					
DINOBRYONOCHROMONAS				# 20	
CYANOPHYTA (BLUE-GREEN ALGAE)					
.CYANOPHYCEAE					
CHROCCOCCALESCHROCCOCCAEAE					
····AGMENELLUM ····ANACYSTIS	* 0		# 20 # 20	# 20	
• HORMOGONALES • • NOSTOCACEAE					
ANABAENA	# 17	# 25			
••••APHANIZOMFNON •••OSCILLATORIACEAE	# 17	# 25	# 20	# 20	
····ARTHROSPIRA ····OSCILLATORIA	# 17 # 17		# 20		
PHORMIMIDIUMCHROCCOCCALES					
CHROCCOCCAEAE	* 0				
••••GOMPHOSPHAERIA EUGLENOPHYTA (EUGLENOIDS)	* 0	# 25	# 20		
•CRYPTOPHYCEAE ••CRYPTOMONIDALES					
CRYPTOCHRYSIDACEAE					
CHROOMONAS CRYPTOMONODACEAE	-	-			
CRYPTOMONAS .EUGLENOPHYCEAE					
EUGLENALESEUGLENACEAE					
TRACHELOMONAS	# 17				# 25
PYRRHOPHYTA (FIRE ALGAE)					
.DINOPHYCEAEPERIDINIALES					
•••CEPATIACFAE •••CERATIUM	* 0				
GL ENODINIACEAE		-			
GLENODINIUM PERIDINIACFAE	•	-	-		-
* = Present in sample but not a domi	nant type				-
# = A dominant type in sample.					

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

			Lake	site L9			
	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Oxygen, dis- solved	Oxygen, dis- solved (per- cent satur-
Date		(ft)	(µS/cm)	units)	(°C)	(mg/L)	ation)
			VE-911	<u> </u>	1 -7	\ 3 <i>t</i> = 1	
April 197							
06	1210	1.60	76	7.7	5.0	13.9	109
06	1211	3.30	77	7.7	5.0	13.9	109
06	1212	6.60	77	7.7	5.0	13.9	109
06	1213	9.80	77	7.7	5.0	13.9	109
06	1214	11.6	77	7.7	5.0	13.9	109
27	1400	1.60	82	8.8	12.0	12.6	117
27 27	1402 1404	3.30 6.60	83 83	8.8 8.8	12.0 12.0	12.6 12.5	117 116
27	1404	9.80	81	7.9	11.0	11.2	102
27	1408	11.6	81	8.1	10.5	11.5	102
27 May 1977	1400	11.0	01	0.1	10.5	11.5	103
16	1440	1.60	87	8.3	15.5	11.2	111
16	1441	3.30	87	8.3	15.0	11.2	110
16	1442	6.60	87	8.3	15.0	11.2	110
June 1977		0.00	σ,	0,0	10.0		110
13	1440	1.60	89	6.0	18.5	10.0	106
13	1442	3.30	89	5.9	18.5	10.0	106
13	1444	6.60	89	5.7	18.0	10.1	106
13	1446	9.80	89	5.7	18.0	10.0	105
			l aka	cita 110			
			Lake	site L10			
						0× y	jen,
			Spe-			0× y	dis-
		_	cific				dis- solved
		Sam-	cific con-	pH		0x ygen ,	dis- solved (per-
		pling	cific con- duct-	(stand-	Temper-	0xygen, dis-	dis- solved (per- cent
	Time		cific con-		ature	0x ygen ,	dis- solved (per-
Date	Time	pling	cific con- duct-	(stand-		0x ygen , dis- sol ved	dis- solved (per- cent
		pling depth	cific con- duct- ance	(stand- ard	ature	0xygen, dis- solved	dis- solved (per- cent satur-
April 197	'7	pling depth (ft)	cific con- duct- ance (US/cm)	(stand- ard units)	ature (°C)	0×ygen, dis- solved (mg/L)	dis- solved (per- cent satur- ation)
April 197 06	7 1230	pling depth (ft)	cific con- duct- ance (µS/cm)	(stand- ard units) 7.8	ature (°C) 5.0	Oxygen, dis- solved (mg/L)	dis- solved (per- cent satur- ation)
April 197 06 06	7 1230 1231	pling depth (ft) 1.60 3.30	cific con- duct- ance (µS/cm)	(stand- ard units) 7.8 7.8	ature (°C) 5.0 5.0	0xygen, dis- solved (mg/L) 14.0	dis- solved (per- cent satur- ation)
April 197 06 06 06	7 1230 1231 1232	pling depth (ft) 1.60 3.30 6.60	cific con- duct- ance (µS/cm)	(stand- ard units) 7.8 7.8 7.8	ature (°C) 5.0 5.0 5.0	0xygen, dis- solved (mg/L) 14.0 14.0	dis- solved (per- cent satur- ation)
April 197 06 06 06	77 1230 1231 1232 1233	pling depth (ft) 1.60 3.30 6.60 8.20	cific con- duct- ance (µS/cm)	(stand- ard units) 7.8 7.8 7.8 7.8	sture (°C) 5.0 5.0 5.0 5.0	0xygen, dis- solved (mg/L) 14.0 14.0 14.0	dis- solved (per- cent satur- ation) 109 109 109
April 197 06 06 06 06	1230 1231 1232 1233 1234	pling depth (ft) 1.60 3.30 6.60 8.20 9.80	cific con- duct- ance (µS/cm) 77 77 77 77 77	(stand- ard units) 7.8 7.8 7.8 7.8 7.8	ature (°C) 5.0 5.0 5.0 5.0 5.0	0xygen, dis- solved (mg/L) 14.0 14.0 14.0 14.0	dis- solved (per- cent satur- ation) 109 109 109 109
April 197 06 06 06 06 27	1230 1231 1232 1233 1234 1415	pling depth (ft) 1.60 3.30 6.60 8.20 9.80 1.60	cific con- duct- ance (μS/cm) 77 77 77 77 77 77 77 81	(stand- ard units) 7.8 7.8 7.8 7.8 7.8 8.9	sture (°C) 5.0 5.0 5.0 5.0 5.0 12.0	0xygen, dis- solved (mg/L) 14.0 14.0 14.0 14.0 12.8	dis- solved (per- cent satur- ation) 109 109 109 109 109
April 197 06 06 06 06 27	7 1230 1231 1232 1233 1234 1415 1416	pling depth (ft) 1.60 3.30 6.60 8.20 9.80 1.60 3.30	cific con- duct- ance (µS/cm) 77 77 77 77 77 77 81 82	(stand- ard units) 7.8 7.8 7.8 7.8 7.8 8.9 8.9	ature (°C) 5.0 5.0 5.0 5.0 5.0 12.0	0xygen, dis- solved (mg/L) 14.0 14.0 14.0 12.8	dis- solved (per- cent satur- ation) 109 109 109 109 119 119
April 197 06 06 06 06 27 27	7 1230 1231 1232 1233 1234 1415 1416 1417	pling depth (ft) 1.60 3.30 6.60 8.20 9.80 1.60 3.30 6.60	cific con- duct- ance (///S/cm) 77 77 77 77 77 81 82 81	(standard units) 7.8 7.8 7.8 7.8 7.8 7.8 9.9 8.9	5.0 5.0 5.0 5.0 5.0 12.0 12.0	0xygen, dis- solved (mg/L) 14.0 14.0 14.0 12.8 12.8 12.8	dis- solved (per- cent satur- ation) 109 109 109 109 119 119
April 197 06 06 06 06 27 27 27	7 1230 1231 1232 1233 1234 1415 1416	pling depth (ft) 1.60 3.30 6.60 8.20 9.80 1.60 3.30	cific con- duct- ance (µS/cm) 77 77 77 77 77 77 81 82	(stand- ard units) 7.8 7.8 7.8 7.8 7.8 8.9 8.9	ature (°C) 5.0 5.0 5.0 5.0 5.0 12.0	0xygen, dis- solved (mg/L) 14.0 14.0 14.0 12.8	dis- solved (per- cent satur- ation) 109 109 109 109 119 119
April 197 06 06 06 06 27 27 27 May 1977	1230 1231 1232 1233 1234 1415 1416 1417 1418	1.60 3.30 6.60 8.20 9.80 1.60 3.30 6.60 8.20	cific conduct - ance (µS/cm) 77 77 77 77 77 81 82 81 81	(standard units) 7.8 7.8 7.8 7.8 7.8 8.9 8.9 8.9 8.9 8.8	5.0 5.0 5.0 5.0 5.0 12.0 12.0 12.0	0xygen, dis- solved (mg/L) 14.0 14.0 14.0 12.8 12.8 12.8	dis- solved (per- cent satur- ation) 109 109 109 109 119 119 119
April 197 06 06 06 06 27 27 27 May 1977	1230 1231 1232 1233 1234 1415 1416 1417 1418	1.60 3.30 6.60 8.20 9.80 1.60 8.20	cific con- duct- ance (µS/cm) 77 77 77 77 81 82 81 81 81	(stand- ard units) 7.8 7.8 7.8 7.8 7.8 9.9 8.9 8.9	ature (°C) 5.0 5.0 5.0 5.0 12.0 12.0 12.0 12.0	0xygen, dis- solved (mg/L) 14.0 14.0 14.0 12.8 12.8 12.7 11.6	dis- solved (per- cent satur- ation) 109 109 109 109 119 119 118
April 197 06 06 06 06 27 27 27 47 May 1977 17	1230 1231 1232 1233 1234 1415 1416 1417 1418	1.60 3.30 6.60 8.20 9.80 1.60 3.30 6.60 8.20	cific con- duct- ance (\(\mu S/cm\)) 77 77 77 77 81 82 81 81 84	(stand- ard units) 7.8 7.8 7.8 7.8 7.8 9.9 8.9 8.9 8.9 8.9 8.8	sture (°C) 5.0 5.0 5.0 5.0 12.0 12.0 12.0 16.5	0xygen, dis- solved (mg/L) 14.0 14.0 14.0 12.8 12.8 12.7 11.6	dis- solved (per- cent satur- ation) 109 109 109 119 119 118 118
April 197 06 06 06 06 27 27 27 27 17 17	1230 1231 1232 1233 1234 1415 1416 1417 1418 1035 1036 1038	pling depth (ft) 1.60 3.30 6.60 8.20 9.80 1.60 3.30 6.60 8.20	cific conduct - ance (µS/cm) 77 77 77 77 77 81 82 81 81 84 84	(stand- ard units) 7.8 7.8 7.8 7.8 7.8 8.9 8.9 8.9 8.9 8.9 8.8 8.3 8.4 8.3	ature (°C) 5.0 5.0 5.0 12.0 12.0 12.0 16.5 16.0 15.0	0xygen, dis- solved (mg/L) 14.0 14.0 14.0 12.8 12.8 12.7 11.6 11.6 11.8	dis- solved (per- cent satur- ation) 109 109 109 109 119 119 118 118 118
April 197 06 06 06 06 27 27 27 27 17 17 17	1230 1231 1232 1233 1234 1415 1416 1417 1418 1035 1036 1038 1039	1.60 3.30 6.60 8.20 9.80 1.60 3.30 6.60 8.20	cific con- duct- ance (\(\mu S/cm\)) 77 77 77 77 81 82 81 81 84	(stand- ard units) 7.8 7.8 7.8 7.8 7.8 9.9 8.9 8.9 8.9 8.9 8.8	sture (°C) 5.0 5.0 5.0 5.0 12.0 12.0 12.0 16.5	0xygen, dis- solved (mg/L) 14.0 14.0 14.0 12.8 12.8 12.7 11.6	dis- solved (per- cent satur- ation) 109 109 109 119 119 118 118
April 197 06 06 06 06 27 27 27 27 17 17 17 June 1977	1230 1231 1232 1233 1234 1415 1416 1417 1418 1035 1036 1038 1039	pling depth (ft) 1.60 3.30 6.60 8.20 9.80 1.60 3.30 6.60 8.20	cific conduct - ance (µS/cm) 77 77 77 77 77 81 82 81 81 84 84	(stand- ard units) 7.8 7.8 7.8 7.8 7.8 8.9 8.9 8.9 8.9 8.9 8.8 8.3 8.4 8.3	ature (°C) 5.0 5.0 5.0 12.0 12.0 12.0 16.5 16.0 15.0	0xygen, dis- solved (mg/L) 14.0 14.0 14.0 12.8 12.8 12.7 11.6 11.6 11.8	dis- solved (per- cent satur- ation) 109 109 109 119 119 118 118 117 116
April 197 06 06 06 06 27 27 27 27 17 17 17	1230 1231 1232 1233 1234 1415 1416 1417 1418 1035 1036 1039	pling depth (ft) 1.60 3.30 6.60 8.20 9.80 1.60 3.30 6.60 8.20 1.60 3.30 6.60 8.20	cific conduct duct ance (µS/cm) 77 77 77 77 81 82 81 81 82 81 81 84	(stand- ard units) 7.8 7.8 7.8 7.8 7.8 8.9 8.9 8.9 8.9 8.9 8.3 8.4 8.3 8.4	ature (°C) 5.0 5.0 5.0 5.0 12.0 12.0 12.0 15.0 16.5 16.0 14.5	0×ygen, dis- solved (mg/L) 14.0 14.0 14.0 12.8 12.8 12.7 11.6 11.8	dis- solved (per- cent satur- ation) 109 109 109 119 119 118 118 117 116 113
April 197 06 06 06 06 27 27 27 27 17 17 17 June 1977	1230 1231 1232 1233 1234 1415 1416 1417 1418 1035 1038 1039	1.60 3.30 6.60 8.20 9.80 1.60 3.30 6.60 8.20 1.60 8.20	cific conduct-ance (µS/cm) 77 77 77 77 81 82 81 81 84 84 82 82 87	(stand-ard units) 7.8 7.8 7.8 7.8 7.8 8.9 8.9 8.9 8.9 8.8 8.3 8.4 6.9	ature (°C) 5.0 5.0 5.0 5.0 12.0 12.0 12.0 12.0 15.0 14.5	0xygen, dis- solved (mg/L) 14.0 14.0 14.0 12.8 12.8 12.7 11.6 11.8 11.6	dis- solved (per- cent satur- ation) 109 109 109 109 119 119 118 118 118 117 116 113
April 197 06 06 06 06 06 27 27 27 27 17 17 17 17 17 14 14	1230 1231 1232 1233 1234 1415 1416 1417 1418 1035 1038 1039	pling depth (ft) 1.60 3.30 6.60 8.20 9.80 1.60 3.30 6.60 8.20 1.60 3.30 6.60 8.20	cific conduct - ance (µS/cm) 77 77 77 77 77 81 82 81 81 84 84 84 82 82 87	(stand- ard units) 7.8 7.8 7.8 7.8 7.8 8.9 8.9 8.9 8.9 8.9 8.3 8.4 6.9 6.8	ature (°C) 5.0 5.0 5.0 12.0 12.0 12.0 16.5 16.0 14.5 19.5	0xygen, dis- solved (mg/L) 14.0 14.0 14.0 12.8 12.8 12.7 11.6 11.8 11.6	dis- solved (per- cent satur- ation) 109 109 109 109 119 119 118 118 117 116 113

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

								Ovygon				
	Time	Sam- pling de pth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Trans- par- ency (secchi disk)	0xygen, dis- solved	Oxygen, Oxygen dis- solved (per- cent satur-	Bicar- bonate FET-fld (mg/L as	Alka- linity field (mg/L as	Carbon dioxide dis- solved (mg/L	Nitro- gen, NO2+NO3 total (mg/L
Date		(ft)	(US/cm)	units)	(°C)		(mg/L)	at ion)	HC03)		CO2)	as N)
			Marit									
lpril 19		1 60	26			1 20		100	10	1.5	-	20
06	1245	1.60	76	7.8	5.0	1.30	14.0	109	18	15	.5	.20
06	1247	3.30	76	7.8	5.0		14.0	109				
06	1248	6.60	76	7.8	5.0		14.0	109				
06	1249	9.80	76	7.8	5.0		13.9	109				
06	1250	13.2	76 76	7.7 7.7	5.0		13.9	109				
06	1252	16.5			5.0		13.9	109	10			
06	1255	17.2	76	7.7			13.9	100	19	16		.21
06	1257 1259	20.0 23.1	76 76	7.7	5.0 5.0		13.9	109 109				
06	1301	26.4	76 76	7.7	5.0		13.9	109				
	1301	30.0	76 76	7.7	5.0		13.9	109				
06	1302	33.0	76 78	7.7	5.0		13.9	109				
06	1304	34.7	76 76	7.7	5.0		13.9	109		16	.6	.20
27	1425	1.60	82	8.9	12.0	1.50	12.8	119	20	16	.0	<.10
27	1423	3.30	82	8.9	12.0	1.50	12.9	120				\.10
27	1428	6.60	82	8.9	11.5		12.8					
27	1430	9.80	82	8.9	11.5		12.8		19	16	.0	<.10
27	1432	13.2	81	8.8	11.0		12.7	118				1,10
27	1435	16.5	79	7.7	10.0		11.6			16	.6	.03
27	1436	20.0	79	7.1	9.5		11.3					
27	1437	23.1	79	6.6	8.5		10.8					
27	1438	26.4	80	6.4	8.5		9.5					
27	1439	30.0	80	6.4	8.5		9.2					
27	1440	33.0	82	6.4	8.0		7.1			18	14	.06
lay 197		55.0	-		0.0			•		~~	- '	
17	1100	1.60	86	8.5	16.0	1.70	12.1	122				<.10
17	1101	3.30	87	8.8	16.0		12.2					
17	1105	6.60	85	8.8	14.5		12.9			16	.0	<.10
17	1106	9.80	84	8.6	13.5		12.4					
17	1110	11.0							25	21		<.10
17	1111	13.2	84	7.2	13.0		11.2	106				
17	1112	16.5	84	6,8	13.0		10.4					
17	1113	20.0	84	6.8	12.5		10.1					
17	1114	23.1	84	6.7	12.5		9.8					
17	1115	26.4	83	6.6	12.5		9.4	89				
17	1119	30.0	85	6.5	12.5		9.0					
17	1120	33.0	86	6.5	12.5		12.5			19	12	.01
June 19	77											
14	1200	1.60	87	7.2	19.5	3.8	10.4	112		19	2.3	<.10
14	1201	3.30	87	7.2	19.0		10.4	111				
14	1202	6.60	87	7.2	18.5		10.4	110				

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

				<u>L</u>	ake site	L11Cont	inued				
	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Trans- par- ency (secchi disk)	Oxygen, dis- solved	Oxygen, dis- solved (per- cent satur-	Alka- linity field (mg/L as	Carbon dioxide dis- solved (mg/L	Nitro- gen, NO2+NO3 total (mg/L
Date		(ft)	(µS/cm)	units)	(°C)	(m)	(mg/L)	at ion)	CACO3) a:	s CO2)	as N)
		-									
June 19			0.0					•••			
14	1203	9.80	86	7.2	17.5		10.2	106			. 10
14	1205	13.1	86	7.2	17.0		10.0	103	19	2.3	<.10
14	1206	16.5	86	7.1	16.0		8.0	81			
14	1210	19.7	85 96	7.2	15.0		5.0	49	19	2.3	<.10
14 14	1211 1212	23.1 26.4	85 85	7.1 7.1	14.0 13.5		2.0	19 0			
14	1213	30.0	90	7.0	12.5			Ö			
14	1215	31.2	92	6.9	12.5		.0 .1	1	22	5.4	<.10
July 19		31.2	32	0.3	12.5		.1	•	22	3.4	₹, 10
11	1300	1.60	97	7.9	25.0	2.40	9.7	115	20	.5	<.10
11	1302	3.30	97	7.9	25.0		9.8	117			
11	1304	6.60	97	7.9	25.0		9.8	117			
11	1305	9.80	97	7.9	25.0		9.8	117	19	.5	. 01
11	1307	13.0	94	7.8	22.5		8.8	101			
11	1309	16.0	92	7.8	19.0		3.8	40			
11	1310	18.0							20		<.10
11	1312	20.0	91	7.8	16.5		.4	4			
11	1314	23.0	89	7.9	14.5		1.0	10			
11	1315	26.0	92	7.9	13.5		.0	Ō	23	.6	<.10
11	1318	30.0	101	7.9	13.0		.0	0			
11	1320	31.0	101	7.9	13.0		.0	0	26	.6	.03
August	1977										
05	1045	1.60	95	7.6	25.0	2.40	9.3	111	20	1.0	<.10
05	1046	3.30	95	7.4	25.0		9.3	111			
05	1047	6.60	95	7.2	24.5		9.3	110			
05	1048	9.80	94	7.0	24.0		9.0	106			
05	1049	13.0	94	6.9	23.5		8.4	98			
05,	1050	16.0	93	6.9	22.0		6.8	77	20	4.9	<.10
05	1052	20.0	88	6.9	19.0		.6	6			
05	1055	21.0							19		<.10
05	1057	23.0	87	7.0	16.5		.0	0			
05	1059	26.0	100	7.0	14.5		.0	0			
05	1100	28.0	104	7.0	14 6				23		<.10
05	1102	30.0	104	7.0	14.5		.0	0		5.2	
05	1105	31.0	108	7.0	13.0		.0	0	27	5.2	.01
	er 1977	1 60	07		24 0	2.7	0 5	100	20		- 10
07	1310	1.60	87 84		24.0 23.5		8.5	100 99	20		<.10
07 07	1311 1312	3.30	84 85		23.5		8.5 8.4	99			
07	1312	6.60 9.80	85 85		23.0		8.2	94	20		<.10
07	1315	13.0	85 87		22.0		6.8	77	20		<.10
٠,	1320	16.0	86		21.0		4.2	47	20		<.10

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Trans- par- ency (secchi disk)	Oxygen, dis- solved	0xygen, dis- solved (per- cent satur-	Alka- linity field (mg/L as	Carbon dioxide dis- solved (mg/L	Nitro- gen, NO2+NO3 total (mg/L
Date		(ft)	(µS/cm)	units)	(°C)	(m)	(mg/L)	at ion)	CACO3) a	s CO2)	as N)
Septembe							_	_			
07	1321	20.0	90		19.5		.6	6			
07	1325	23.0	. 89		17.0		.2	2	22		<.10
07	1326	26.0	112		15.0		.3	3			
07	1330	3 0.0	124		14.0		. 4	4	32		<.10
October					16.5						22
05	1145	1.60	111		16.5	1.90	7.6	78	23		. 03
05	1146	3.30	113		16.0		7.6	77			
05	1147	6.60	113		16.0		7.2	73			
05	1150	9.80	115		15.5		6.8	67	25		.03
05	1151	13.2	115		15.5		6.6	65			
05	1155	16.4	115		15.5		6.6	65	24		.04
05	1156	19.8	115		15.5		6.6	65			
05	1200	23.0	116		15.5		6.6	65	25		.03
05	1201	26.0	116		15.5		6.9	68			
05	1205	30.0	116		15.5		7.0	69			.04
21	1105	1.60	90	7.1	12.0	1.80	7.1	65	22	3.4	.07
21	1106	3.30	90	7.1	12.0		7.1	65			
21	1107	6.60	90	7.1	12.0		7.0	64			
21	1110	9.80	90	7.1	12.0		7.0	64	22	3.4	.06
21	1111	13.2	90	7.0	12.0		7.0	64			
21	1115	16.4	90	7.1	12.0		7.0	64	23	3.5	.06
21	1116	20.0	90	7.0	12.0		6.9	63			
21	1117	23.0	90	7.0	12.0		6.9	63	22	4.3	
21	1120	26.2	90	7.0	12.0		6.8	62	22	4.3	.07
21	1121	30.0	90	7.0	11.5		6.8	61			
21	1125	31.2	90	7.0	11.5		6.8	61	22	4.3	.08
November	-	1 60	~-								
16	1350	1.60	85	7.2	9.5		4.9	44			
16	1351	3.30	85	7.2	9.5		4.9	44			
16	1352	6.60	85	7.2	9.5		4.8	43			
16	1353	9.80	85	7.2	9.5		4.8	43			
16	1354	13.2	85	7.2	9.5		4.8	43			
16	1355	16.0	85	7.2	9.5		4.8	43			
16	1356	20.0	85	7.2	9.5		4.8	43			
16	1357	23.0	85	7.2	9.5		4.7	41			
16	1358	26.0	85	7.2	9.5		4.7	41			
16	1359	30.0	85	7.2	9.5		4.6	40			

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

			<u>L</u>	ake site	L11Cont	inued			
Date	Time	Sam- pling depth (ft)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as ND3)	Phos- phorus, total (mg/L as P)	Phos- phorus, ortho, total (mg/L as P)	Carbon, organic total (mg/L as C)
April 19	77								
06	1245	1.60	.090	. 31	.60	2.7	.010	<.010	5.2
06	1247	3.30							
06	1248	6,60							
06	1249	9.80							
06	1250	13.2							
06	1252	16.5							
06	1255	17.2	.080	.32	.61	2.7	.010	<.010	3.3
06	1257	20.0							
06	1259	23.1							
06	1301	26.4							
06	1302	30.0							
06	1304	33.0							
06	1305	34.7	.080	. 42	.70	3.1	.010	<.010	3.0
27	1425	1.60	.040	. 51	.55	2.4	.020	<.010	
27	1427	3.30							
27	1428	6.60							
27	1430	9.80	.080	. 50	.47	2.1	.030	<.010	
27	1432	13.2							
27	1435	16.5	.040	. 41	. 48	2.1	.020	<.010	
27	1436	20.0							
27	1437	23.1							
27	1438	26.4							
27	1439	30.0							
27	1440	33.0	.220	. 58	.86	3.8	.040	<.010	
May 1977									
17	1100	1.60	.030	. 35	.38	1.7	.020	<.010	
17	1101	3.30							
17	1105	6.60	.050	. 40	.45	2.0	.030	<.010	
17	1106	9.80							
17	1110	11.0	.060	. 37	.43	1.9	. 030	<.010	
17	1111	13.2							
17	1112	16.5							
17	1113	20.0							
17	1114	23.1							
17	1115	26.4							
17	1119	30.0							
17	1120	33.0	.090	. 44	.54	2.4	. 040	<.010	
June 197				20	^-		000	. ^1^	
14	1200	1.60	.040	. 35	.39	1.7	.020	<.010	
14	1201	3.30							
14	1202	6.60							

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

			L	ake site	L11Cont	inued			
Date	Time	Sam- pling depth (ft)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)	Phos- phorus, ortho, total (mg/L as P)	Carbon, organic total (mg/L as C)
June 197	7								
14	1203	9.80							
14	1205	13.1	.080	. 28	.36	1.6	.020	<.010	
14	1206	16.5							
14	1210	19.7	.090	. 20	.29	1.3	.030	<.010	
14	1211	23.1							
14	1212	26.4							
14	1213	30.0							
14	1215	31.2	.110	. 39	.50	2.2	. 060	<.010	
July 197	1200	1 60	000	• • • • • • • • • • • • • • • • • • • •	4.6	2.0	000		
11	1300	1.60	.030	. 43	.46	2.0	.020		
11 11	1302 1304	3.30 6.60							
11	1305	9.80	.050	. 45	.51	2.3	.020		
11	1303	13.0	.030		.51	2.5	.020		
11	1309	16.0							
11	1310	18.0	.110	. 46	.57	2.5	.020		
11	1312	20.0							
11	1314	23.0							
11	1315	26.0	.140	. 58	.72	3.2	.060		
11	1318	30.0							
11	1320	31.0	.310	.89	1.2	5.4	.130		
August 1									
05	1045	1.60	.020	. 49	.51	2.3	.020		5.6
05	1046	3.30							
05	1047 1048	6.60 9.80						•-	
05 05	1048	13.0							
05	1050	16.0	.030	.38	.41	1.8	.030		5.3
05	1052	20.0					.000		
05	1055	21.0	. 040	. 48	.52	2.3	.030		4.1
05	1057	23.0							
05	1059	26.0							
05	1100	28.0	.040	1.1	1.1	4.9	.050		5.1
05	1102	30.0							
05	1105	31.0	. 240	.76	1.0	4.5	. 100		10
Septembe									
07	1310	1.60	.010	. 26	.27	1.2	.020		
07	1311	3.30							
07	1312	6.60 9.80	030	.27	.30	1.3	.020		
07 07	1315 1316	13.0	.030	.2/	.30	1.3	.020		

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

				L11Cont			
Date	Time	Sam- pling depth (ft)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)
Septembe	ar 1077						
D7	1321	20.0					
07	1325	23.0	.080	.49	. 57	3.5	.050
07	1326	26.0					
07	1330	30.0	.650	.35	1.0	4.4	.180
October							
05	1145	1.60	.140	.39	. 56	2.5	.110
05	1146	3.30			•••		
05	1147	6.60					
05	1150	9.80	.140	.38	. 55	2.4	.150
05	1151	13.2					
05	1155	16.4	.140	.40	. 58	2.6	.030
05	1156	19.8					
05	1200	23.0	.140	.37	. 54	2.4	.030
05	1201	26.0					
05	1205	30.0	.130	.36	. 53	2.3	.070
21	1105	1.60	.140	.61	.82	3.6	.040
21	1106	3.30					
21	1107	6.60					
21	1110	9.80	.170	.53	. 76	3.4	.040
21	1111	13.2					
21	1115	16.4	.160	.40	. 62	2.7	.040
21	1116	20.0					
21	1117	23.0	150	40		2.1	
21	1120	26.2	.150	.48	.70	3.1	.030
21	1121	30.0	160			2.7	040
21	1125	31.2	.160	.60	. 84	3.7	.040
Novembe 16	1350	1.60					
16	1350	3.30					
16	1351	6.60					
16	1353	9.80					
16	1354	13.2					
16	1355	16.0					
16	1356	20.0					
16	1357	23.0					
16	1358	26.0					
16	1359	30.0					

041F TIME	APR	6•77 1245		27 , 77		27•77 425		17.77		14.77
DIVEPSITY: DIVISION CLASS ORDER FAMILY GENUS		0.9 0.9 1.6 1.6		0.9 0.9 0.9 1.6 1.6		0.9 0.9 0.9 1.6 1.6		1.5 1.5 1.5 2.0 2.0		0.8 0.8 1.5 2.0 2.0
ORGANISM	CELL5 /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT
CHLOROPHYTA (GREEN ALGAE)										
.CHLOROPHYCEAECHLOROCOCCALES										
CHARACIACEAESCHROEDERIA		_		_		_		_		_
MICRACTINIACEAF										
MICRACTINIUMOOCYSTACEAE		-		0		-		-		•
ANKISTRODESMUSCHODATELLA		-	*	0		-		- 25		-
CLOSTERIOPSIS		-		0		-		-		-
DICTYOSPHAERIUMNFPHROCYTIUM		-		-		-	*	0 0		-
••••TETRAEDRON •••SCENEDESMACEAE		-	*	0		-		0		-
• • • • SCENEDESMUS		-		0		-		0		-
TETRASTRUMTETRASPORALES		-	*	0		-		-		-
PALMELLACEAE										
SPHAEROCYSTISVOLVOCALES		-		-		-		-		-
CHLAMYDOMONADACEAECHLAMYDOMONAS		_		0		_		_		_
ZYGNEMATALES				ŭ						
DESMIDIACEAECLOSTERIUM		-		-		-		-		-
STAURASTRUM		-		~		-		-		-
CHRYSOPHYTA										
.BACILLARIOPHYCEAECENTRALES										
COSCINODISCACEAE		33		0			٠	0		
MELOSIRA	"	-		0		-		0		-
••PENNALES •••FRAGILARIACEAE										
· · · · ASTERIONELLA	#		*		#		#	25	#	
FRAGILARIA SYNEDRA		-		-		-	*	0		-
NITZSCHIACEAENITZSCHIA		_		0		_		0		_
.CHRYSOPHYCEAE				v				U		
CHRYSOMONADALESOCHROMONADACEAE										
DINOBRYONSYNURACEAE	~ ~	-		-		-		-		-
SYNURA		-		-		-		-		-
CYANOPHYTA (BLUE-GREEN ALGAE)										
.CYANOPHYCEAECHROCCOCCALES										
CHROCCOCCAEAE										
AGMENELLUM ANACYSTIS		-		0		-		-	#	25
HORMOGONALESNOSTOCACEAE										
ANABAENA		-		-		-		-	#	25
APHANIZOMENONOSCILLATORIACEAE		-	#	33	*	33	*	25		-
····ARTHROSPIRA ····LYNGBYA		-		_		-		-		-
OSCILLATORIA	#	33	#	33	*	33	#	25	#	25
CHROCCOCCALESCHROCCOCCAEAE										
GOMPHOSPHAERIA		-		-		-		-	*	0
EUGLENOPHYTA (EUGLENOIDS)										
.CRYPTOPHYCEAECRYPTOMONIDALES										
CRYPTOCHRYSIDACEAECHROOMONAS		_		0		_		_		_
CRYPTOMONODACEAE		_				_		_		_
CRYPTOMONAS .FUGLENOPHYCEAE		-	*	0		-		-		-
EUGLENALES										
EUGLENACEAETRACHELOMONAS		-		-		-		-		0
PYPRHOPHYTA (FIRE ALGAE)										
.DINOPHYCEAE										
PERIDINIALESCEPATIACEAE										
CERATIUM GLENODINIACEAF		-		-		-		-		-
GL ENODINIUM		-		-		-	#	0		-
PERIDINIACEAE,PERIDINIUM		_		0		_		_		-
<pre>* = Present in sample but not a dominar # = A dominant type in sample.</pre>	it type			•						

^{# =} A dominant type in sample.

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued Lake site L11--Continued

			Phytop	lankton	anarvses					
DATE TIME		11.77		5•77 245		17•77 150		7•77 310		21•77 1105
DIVERSITY: DIVISION CLASS ORDER FAMILY OFFICE OFF		0.3 0.3 1.1 1.5		1.3 1.3 1.3 1.9 2.6		1.4 1.4 1.9 1.9		0.7 0.7 1.4 1.4 2.3		0.9 1.6 1.6 1.6
	CELLS	PER-	CELLS	PER-	CELLS	PER-	CELLS	PER-	CELLS	PER-
ORGANISM	/ML	CENT	/ML	CENT	/ML	CENT	/ML	CENT	/ML	CENT
CHLOROPHYTA (GREEN ALGAE) .CHLOROPHYCEAECHLOROCCALESCHARACIACEAESCHROEDERIA	200	1					<u></u>			
MICRACTINIACEAEMICRACTINIUM		-		-		-		-		-
OOCYSTACEAEANKISTRODESMUS		-	*	0		-		-		-
CHODATELLACLOSTERIOPSIS		-		-		-		-		-
DICTYOSPHAERIUM		-		-		-		-		-
••••NEPHROCYTIUM ••••TETRAEDRON		-		-		-		-		-
SCENEDESMACEAE		-		-		_		=	••	-
••••SCENEDESMUS ••••TETPASTRUM		-		-		-		-		-
•• TETRASPORALES		-		-		-		-		-
•••PALMELLACEAE •••SPHAERDCYSTIS ••VOLVOCALES ••CHLAMYDOMONADACEAE	480	2	#	17		-		-		-
CHLAMYDOMONASZYGNEMATALES		-		-		-		-		-
DESMIDIACEAECLOSTERIUM		_	*	0		_		_		_
STAURASTRUM	*	0	*	ő		-		-		-
CHRYSOPHYTA *BACILLARIOPHYCEAE *.CENTRALES										
COSCINODISCACEAE										
····CYCLOTELLA ····MELOSIRA		-		-		-		-	#	33
• • PENNALES										
FRAGILARIACEAEASTERIONELLA		0		-		-		-		-
FRAGILARIA		-	*	0	#	20		-		-
····SYNEDRA ···NITZSCHIACEAE		-		-		-		-		-
NITZSCHIA .CHRYSOPHYCEAE		-		-		-		-		-
••CHRYSOMONADALES •••CHROMONADACEAE										
····DINOBRYDN ···SYNURACEAE		-		0		-		-	*	33
SYNURA	*	0		-		-		-		-
CYANOPHYTA (BLUE-GREEN ALGAE) .CYANOPHYCEAECHROCCOCCALESCHRDCCOCCAEAE										
ANACYSTISHORMOGONALESNOSTOCACEAE	300 930	3		0 -		-	#			-
ANABAENA APHANIZOMENON	3000 1000	9 3		17 17		20 20		-		-
OSCILLATORIACEAEARTHROSPIRA		-	#	17		-		-		-
LYNGBYA OSCILLATORIA	3500 17000#	11 53		17		-	#	20		-
CHROCCOCCALESCHROCCOCCAEAEGOMPHOSPHAERIA	5000#	16	*	0	#	20		20		-
EUGLENOPHYTA (EUGLENOIDS) CRYPTOPHYCEAE CRYPTOMONIDALES						-				
CRYPTOCHRYSIDACEAECHROOMONAS	300	1		-		-		-		~
CRYPTOMONODACEAECRYPTOMONAS .EUGLENOPHYCEAEEUGLENALESFUGLENACEAE		-		-		-		-		-
TRACHELOMONAS	#	0	#	17	*	20	#	20	#	33
PYRRHOPHYTA (FIRE ALGAE) .DINOPHYCEAEPERIDINIALESCEPATIACEAE										
••••CERATIUM •••GLENODINIACEAE	*	0	*	0		-		-		~
• • • • GLENODINIUM		-		-		-		-		-
•••PEPIDINIACEAE •••PERIDINIUM		-		-		-		-		_
* = Present in sample but not a domin	nant type.									

^{* =} Present in sample but not a dominant type.
= Adominant type in sample.

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

			Lake	site L12			
	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	0×ygen, dis- solved	Oxygen, dis- solved (per- cent satur-
Date		(ft)	(µS/cm)	units)	(°C)	(mg/L)	ation)
April 06 06	133D	1.60 3.30 6.60	76 76 76	7.9 7.8 7.8	5.0 5.0 5.0	14.0 13.9 13.9	109 109 109
06 06 06 06	1333 1334 1335 1336 1337	9.80 13.2 16.5 20.0	76 76 76 76	7.7 7.7 7.7 7.7	5.0 5.0 5.0 5.0	13.9 13.9 13.9 13.9	109 109 109 109
06 06 06 27	1338 1339 1340 1500	23.1 26.4 30.0 31.4 1.60	76 76 76 76 84	7.7 7.7 7.6 7.6 8.8	5.D 5.0 5.0 5.0 12.5	13.9 13.9 13.9 13.8 12.5	109 109 109 108 117
27 27 27 27 27	1502 1504 1506 1508	3.30 6.60 9.80 13.2	84 83 82 80	8.9 8.9 8.7 8.2	11.5 11.5 11.0 10.5	13.0 12.8 12.2 11.8	119 117 110 105
27 27 27 27	1512 1514 1516	16.5 20.0 23.1 26.4 28.0	81 81 81 81	7.8 7.2 6.8 6.8 6.8	10.0 9.5 9.0 8.5 8.5	11.5 11.2 10.6 10.3 10.0	102 97 91 87 85
May 19 17 17 17	1135 1136 1137	1.60 3.30 6.60	87 87 86	8.4 8.5 8.7	16.0 15.5 14.0	12.0 12.1 12.6	121 120 121
17 17 17 17	1139 1140 1141	9.80 13.2 16.5 20.0 23.1	84 84 84 84	8.5 7.9 8.4 6.9 6.8	14.0 13.5 13.0 12.5 12.5	12.3 11.8 11.0 10.4 10.2	115 112 104 97 95
17 17 June 1 14	1143 1144 977	26.4 28.0	84 84 90	6.7 6.6 7.3	12.5 12.5 20.5	9.6 9.1 10.2	90 85 112
14 14 14 14	1242 1244 1246	3.30 6.60 9.80 13.2 15.0	89 89 89 88 88	7.3 7.4 7.3 7.2 7.2	19.0 18.0 17.5 16.5 16.5	10.4 10.5 10.6 9.2 8.5	111 111 110 94 87

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

			Lake	site L13	!		
	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper-	Oxygen, dis- solved	Oxygen, dis- solved (per- cent satur-
Date		(ft)	(µS/cm)	units)	(°C)		ation)
Date		(11)	(µ3/cm)	units)	()	(mg/ r)	actony
April 19							
06	1425	1.60	74	8.0	5.0	14.1	110
06	1426	3.30	74	8.0	5.0	14.1	110
06	1427	6.60	74	8.0	5.0	14.1	110
06	1428	9.80	74	8.0	5.0	14.0	109
06	1429	13.2	75	8.0	5.D	14.0	109
06	1430	16.5	75	8.0	5.0	14.0	109
06	1431	20.0	75	8.0	5.0	14.0	109
06	1432	23.1	75	8.0	5.0	14.0	109
06	1433	26.4	75	8.0	5.0	14.0	109
06	1434	30.0	75	8.0	5.0	14.0	109
06	1435	33.0	75	8.0	5.0	14.0	109
			75				
06	1436	36.3		8.0	5.0	14.0	
27	1510	1.60	83	8.9	12.0	12.7	118
27	1512	3.30	83	8.9	11.5	12.8	117
27	1514	6.60	81	8.5	10.5	12.2	109
27	1516	9.80	80	8.4	10.0	11.9	105
27	1518	13.2	80	7.3	9.5	11.4	
27	1520	16.5	80	7.0	9.0	11.2	97
27	1522	20.0	80	6.9	9.0	11.1	96
27	1524	23.1	80	6.7	9.0	10.8	93
27	1526	26.4	81	6.6	8.5	10.2	
27	1528	30.0	79	6,3	8.5	10.2	
27	1530	33.0	81	6.8	8.5	9.2	
May 197		33.0	0.	0.0	0,5	,,,	
17	1205	1.60	88	8.6	15.5	12.2	121
17	1206	3.30	88	8.7	15.0	12.4	
17	1207	6.60	88	8.6	14.5	12.2	
17	1208	9.80	86	8.5	13.5	12.3	
17	1209	13.2	85	6.9	13.0	10.4	
17	1210	16.5	85	6.8	12.5	10.2	
17	1211	20.0	85	6.7	12.5	9.8	
17	1212	23.1	85	6.7	12.5	9.6	
17	1213	26.4	85	6.6	12.5	8.9	
17	1214	30.0	86	6.5	12.5	8.2	77
17	1215	33.0	86	6.5	12.5	8.1	76
June 19							
14	1255	1.60	90	7.2	20.0	10.1	110
14	1257	3.30	89	7.3	18.5	10.3	
14	1259	6.60	89	7.3	18.0	10.4	
14	1300	9.80	88	7.3	16.5	9.8	
			88		16.0		
14	1302	13.2		7.2			
14	1304	16.5	88	7.2	16.0		
14	1306	20.0	88	7.1	16.0		
14	1308	23.1	88	7.1	14.0		
14	1310	26.4	88	7.1	13.0		
14 14	1312	30.0	90	7.1	12.5		
	1314	33.0	94	7.2	12.5	.0	

Table 11.--Chemical, and biological characteristics of lake water--continued

	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Trans- par- ency (secchi disk)	Oxygen, dis- solved	0xygen, dis- solved (per- cent satur-	Bicar- bonate FET-fld (mg/L as	Alka- linity field (mg/L as	Carbon dioxide dis- solved (mg/L	Nitro- gen, NO2+NO3 total (mg/L
Date		(ft)	(µS/cm)	units)	(°C)	(m)	(mg/L)	at ion)	HC03)	CACO3) as	s CO2)	as N)
April 19	77											
06	1455	1.60	76	8.0	5.0	1.20	14.0	109	20	16	.3	.23
06	1456	3.30	76	8.0	5.0		14.0	109				
06	1457	6.60	76	8.0	5.0		14.0	109				
06	1458	9.80	76	8.0	5.0		13.9	109				
06	1459	13.2	76	8.0	5.0		13.9	109				
06	1501	16.5	76	8.0	5.0		13.9	109				
06	1505	20.0	76	8.0	5.0		13.9	109	19	16	.3	.20
06	1506	23.1	76	8.0	5.0		13.9	109				
06	1507	26.4	76	8.0	5.0		13.9	109				
06	1508	30.0	76	8.0	5.0		13.9	109				
06	1509	33.0	76	8.0	5.0		13.8	108				
06	1510	36.3	76	8.0	5.0		13.8	108				
06	1515	38.0			3.0		15.0		19	16		.21
06	1516	39.0	76	8.1	5.0		13.7	108				.21
27	1540	1.60	86	8.9	11.5	1.60	12.8		21	17	.0	<.10
27	1541	3.30	85	8.9	11.5	1.00	12.0	117			.0	
27	1541	6.60	85	8.8	11.0		12.9	118 115	19		.0	.01
27	1545	9.80	84	8.7	10.0		12.3	109	19	16		
27	1546	13.2	83	8.1	9.5		11.7	109				
27	1550	15.2		0.1	9.5			_	21			.02
27	1551	16.5	83	7.0	9.0		11.3	97		17		.02
27	1552	20.0	83	6.8	9.0		11.3	96				
27	1552	23.1	83	6.7	8.5		10.6	90				
27	1554	26.4	82	6.6			10.8					
27	1555	30.0	83		8.0			86				
	1556	33.0	83	6.6 6.5	8.0		10.3	87				
27					8.0		9.6	81				
27 27	1557	36.3	85 86	6.3	8.0		7.9	66	20	16	16	07
27 lay 1977	1600	38.0	86	6.3	8.0		7.2	61	20	16	16	.07
	1325	40.0										
04 17	1225	1.60	90	8.5	16.6	1 60	12 1	122	23	10		- 10
					16.5	1.60	12.1	123		19	.1	<.10
17	1226	3.30	90	8.5	16.0		12.2	123	27			. 10
17	1230	6.60	90	8.7	15.0		12.6	123	27	22	.0	<.10
17	1231	9.80	87	7.4	13.0		11.3	107				
17	1235	11.0					10.0		24	20		<.10
17	1236	13.2	87	7.0	13.0		10.8	102				
17	1237	16.5	87	6.8	12.5		10.2	95				
17	1238	20.0	87	6.7	12.5		9.8	92				
17	1239	23.1	87	6.6	12.5		9.2	86				
17	1240	26.4	87	6.6	12.5		9.0	84				

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

							-Continue					
	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Trans- par- ency (secchi disk)	Oxygen, dis- solved	Oxygen, dis- solved (per- cent satur-	Bicar- bonate FET-fld (mg/L as	Alka- linity field (mg/L as	Carbon dioxide dis- solved (mg/L	Nitro- gen, NO2+NO3 total (mg/L
Date		(ft)	(µS/cm)	units)	(°c)	(m)	(mg/L)	ation)	HC03)	CACO3) a	s CO2)	as N)
1ay 1977												
17	1241	30.0	87	6.5	12.5		8.6	80				
17	1242	33.0	87	6.5	12.0		8.1	75				
17	1243	36.3	87	6.4	12.0		7.7	71				
17	1245	38.0	87	6.4	12.0		7.2	66	24	20	15	.02
June 1977			-						-			
14	1315	1.60	91	7.2	20.0	3.7	10.2	111		18	2.2	<.10
14	1316	3,30	90	7.2	18.5		10.5	111				
14	1318	6.60	89	7.2	17.5		10.5	109				
14	1320	9.80	88	7.3	17.0		10.4	107		18	1.7	<.10
14	1322	13,2	88	7.2	16.5		9.8	100				
14	1323	16.5	88	7.2	16.0		8.6	87				
14	1325	19.7	88	7.2	15.5		6.4	64		18	2.2	<.10
14	1327	23.1	88	7.2	14.0		2.2	21				
14	1329	26.4	88	7.2	13.0		.0	0				
14	1330	30.0	90	7.2	12.5		.0	0				
14	1332	33.0	95	7.2	12.0		.0	0				
14	1333	36.3	103	7.2	11.5		.0	0				
14	1335	37.7	106	7.2	11.5		.0	0		25	3.1	<.10
July 1977	7											
11	1340	1.60	97	7.9	25.0	2.90	9.8	117				<.10
11	1342	3.30	97	7.9	25.0		10.0	119				
11	1344	6.60	97	7.9	24.5		10.0	118				
11	1345	9.80	95	7.9	23.0		9.7	111		19	.5	<.10
11	1347	13.0	93	7.7	21,0		7.0	78				
11	1349	16.0	91	7.8	19.0		3.6	38				
11	1350	20.0	91	7.9	16.5		.6	6		20	.5	<.10
11	1352	23.0	90	7.9	15.0		.2					
11	1353	26.0	95	8.0	13.5		.0					
11	1354	30.0	102	8.0	12.5		.0					
11	1355	33.0	107	7.9	12.0		.0			•	.9	<.10
11	1357	36.0	115	7.9	12.0		٥.					
11	1 400	38.0	115	7.9	12.0		.0	0		35	.9	<.10
August 19				_		_	_					_
05	1115	1.60	96	7.7	25.0						.8	
05	1116	3.30	96	7.7	25.0		9.2					
05	1118	6.60	96	7.4	24.5		9.2					
05	1119	9.80	96	7.1	24.0		9.0					
05	1120	13.0	95	7.1	23.0		8.0				2.9	
		16.0	94	7.1	21.5		5.6	63				
05	1122											
05 05	1125 1127	20.0	90 85	7.3 7.3	19.0 15.0		.5	5		20	1.9	

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

				Ī	ake site	L14Cont	inued					
	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Trans- par- ency (secchi disk)	Oxygen, dis- solved	Oxygen, dis- solved (per- cent satur-	Alka- linity field (mg/L as	Carbon dioxide dis- solved (mg/L	Nitro- gen, NO2+NO3 total (mg/L	
Date		(ft)	(µS/cm)	units)	(°C)	(m)	(mg/L)	at ion)	CACO3) a	s CO2)	as N)	
August 19		30 O	112	7 2	12.0		0	0				
05 05	1129 1130	30.0 33.0	113 118	7.3 7.3	13.0 12.0		.0 .0	0	41	4.0	<.10	
05	1133	36.0	123	7.3	12.0		.0	0		4.0		
05	1135	38.0	131	7.3	11.5		.0	ő	46	4.5	<.10	
September												
07	1340	1.60	87		24.0	3.6	8.6	101	20		<.10	
07	1341	3.30	88		23.5		8.6	100				
07	1342	6.60	88		23.5		8.5	99				
07	1345	9.80	88		23.0		8.5	98	20		.10	
07	1346	13.0	88 87		22.5		8.2	94 46	20		< 10	
Ð7 07	1350 1351	16.0 20.0	88		20.5 19.0		4.2	46 2	20		<.10	
07	1352	23.0	88		16.5		.2	2				
07	1353	26.0	114		14.0		.2	2				
07	1355	30.0	129		13.0		.2	2	28		<.10	
07	1356	33.0	132		12.5		.2	2				
07	1400	36.0	137		12.5		.2	2	51		<.10	
October :		1 60	111		16.5	2 00	0 0	0.2	24		0.E	
05 05	1225 1226	1.60 3.30	111 113		16.5 16.D	2.00	8.0 7.8	82 79	24		.05	
05	1230	6.60	111		16.0		7.5	76	24		.04	
05	1231	9.80	113		16.0		6.9	70				
05	1232	13.2	115		15.5		6.9	68				
05	1235	16.4	115		15.5		7.0	68	25		.03	
05	1236	19.8	115		15.5		7.0	68				
05	1237	23.0	115		15.5		6.9	68				
05	1238	26.0	116		15.5		6.5	64				
05 05	1239 1240	30.0 33.0	122 148		15.0 13.0		4.3	4 2 0	30		.02	
05	1241	36.0	185		12.0		.0	0			.02	
05	1245	37,9	186		12.0		.0	ō	71		.01	
21	1130	1.60	90	7.1	12.5	1.80	6.9	64	23	3.5	.05	
21	1131	3.30	90	7.1	12.5		6.9	64				
21	1132	6.60	90	7.0	12.0		6.8	62				
21	1135	9.80	90	7.0	12.0		6.8	62	21	4.1	.01	
21 21	1136 1137	13.2 16.0	90 90	7.0 7.0	12.0 12.0		6.8 6.7	62 61				
21	1140	19.8	90	7.0	12.0		6.7	61	23	4.5	.06	
21	1141	23.0	90	7.1	12.0		6.7	61				
21	1142	26.0	90	7.0	12.0		6.7	61				
21	1145	29.9	90	7.0	12.0		6.7	61	23	4.5	.06	
21	1146	33.0	90	7.0	12.0		6.7	61				
21 21	1147 1150	36.0 38.0	90 90	7.0 7.0	12.0 12.0		6.7 6.7	61 61	23	4.5	05	
November		JO.U	30	7.0	12.0		0.7	01	23	4.5	. 05	
16	1400	1.60	85	7.2	9.5		4.8	42				
16	1401	3.30	85	7.2	9.5		4.8	42				
16	1402	6.60	85	7.2	9.5		4.7	41				
16	1403	9.80	85	7.2	9.5		4.7	41				
16	1404	13.2	85 05	7.2	9.5		4.6	40				
16	1405	16.4	85 85	7.2	9.5		4.6	40				
16 16	1406 1407	19.8 23.0	85 85	7.2 7.2	9.5 9.5		4.6 4.6	40 40				
16	1407	26.0	85	7.2	9.5		4.5	39				
16	1409	30.0	85	7.2	9.5		4.5	39				
16	1410	33.0	85	7.2	9.5		4.5	39				
16	1411	36.0	85	7.1	9.0		4.4	38				
16	1412	39.0	85	7.1	9.0		4.4	38				
April 19		2.20	or			2.00						
13 13	1320 1325	3.28	85 85	6.9	6.0	2.00	6.0					
	1343	19.7	85	6.7	4.5		6.0					

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

Date	Time	Sam- pling depth (ft)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)	Phos- phorus, ortho, total (mg/L as P)	Carbon, organic total (mg/L as C)	Algal growth poten- tial, bottle test (mg/L)	
April 197	7										
D6	1455	1.60	.060	. 44	.73	3.2	.020	.010	5.5	1.9	
06	1456	3.30									
06	1457	6.60									
06	1458	9.80									
06	1459	13.2									
06	1501	16.5									
06	1505	20.0	.060	. 34	.60	2.7	.010	<.010	3.9		
06	1506	23.1									
06	1507	26.4									
06	1508	30.0									
06	1509	33.0									
06	1510	36.3									
06	1515	38.0	.070	. 43	.71	3.1	.010	<.010	5.1		
06	1516	39.0									
27	1540	1.60	. 040	. 56	.60	2.7	.020	<.010			
27	1541	3.30									
27	1545	6,60	. 050	. 38	.44	1.9	. 030	<.010			
27	1546	9.80									
27	1547	13.2				2.4	020	. 010			
27 27	1550	15.0	.050	. 48	.55	2.4	.020	<.010			
27	1551 1552	16.5 20.0									
27	1553	23.1									
27	1554	26.4									
27	1555	30.0									
27	1556	33.0									
27	1557	36,3									
27	1600	38.0	. 190	. 53	.79	3.5	.040	<.010			
May 1977					• • •						
04	1325	40.0									
17	1225	1.60	.020	. 26	.28	1.2	.010	<.010			
17	1226	3.30									
17	1230	6.60	.060	. 36	.42	1.9	.020	<.010			
17	1231	9.80									
17	1235	11.0	.050	.52	.57	2.5	.020	<.010			
17	1236	13.2									
17	1237	16.5									
17	1238	20.0									
17	1239	23.1									
17	1240	26.4									

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

							-			
		Sam- pling	Nitro- gen, ammonia dis- solved	Nitro- gen, organic total	Nitro- gen, total	Nitro- gen, total	Phos- phorus, total	Phos- phorus, ortho, total	Carbon, organic total	Algal growth poten- tial, bottle
Date	Time	depth (ft)	(mg/L as N)	(mg/L as N)	(mg/L as N)	(mg/L as NO3)	(mg/L as P)	(mg/L as P)	(mg/L as C)	test (mg/L)
May 19 17		30.0								
17		33.0								
17		36.3								
17		38.0	.210	. 35	.58	2.6	.040	<.010		
June		30.0			,,,,					
14		1.60	.030	. 27	.30	1.3	.010	<.010		
14		3.30								
14		6.60								
14	. 1320	9.80	.100	. 57	.67	3.0	.030	<.010		
14		13.2								
14		16.5								
14		19.7	.090	. 32	.41	1.8	.020	<.010		
14		23.1								
14		26.4								
14		30.0								
14		33.0								
14		36.3								
14		37.7	. 300	. 34	.64	2.8	.090	.020		
July		1	0.40		0.0					
11		1.60	.040	. 32	.36	1.6	.020			
11		3.30								
11		6,60				2				
11		9.80	.050	. 52	.57	2.5	.020			
11		13.0		••						
11 11		16.0	.090	. 29	.38	1.7	.020			
11		20.0 23.0	.090	. 29	.38	1.7	.020			
11		26.0								
11		30.0								
11		33.0	.840	. 76	1.6	7.1	.390			
11		36.0	.040	.70	1.0	7.1	.550			
11		38.0	.040	. 38	.42	1.9	.020			
	t 1977	50.5			. 12	,	.020			
05		1,60	.020	. 62	.65	2.9	.020		6.2	2.6
05		3.30							•••	
05		6,60								
05		9.80								
05		13.0	.020	. 50	.52	2.3	.040		4.5	
05		16.0								
05	. 1125	20.0	.020	. 39	.41	1.8	.030		4.5	
05	. 1127	23.0								
05	. 1128	26.0								

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

			<u>L</u> .	ake site	L14Cont	inued		
	Date	Time	Sam- pling depth (ft)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)
Δ	ugust 19	77						
	05	1129	30.0					
	05	1130	33.0	1.40	1.1	2.5	11	.480
	05	1133	36.0					
	05	1135	38.0	1.80	.90	2.7	12	.650
	eptember	1977						
	07	1340	1.60	.010	.34	. 35	1.6	.020
	07	1341	3.30					
	07	1342	6.60	060	21	47	2 1	0.40
	07	1345 1346	9.80 13.0	.060	.31	. 47	2.1	.040
	07	1350	16.0	.020	.34	. 36	1.6	.030
	07	1351	20.0	.020				
	07	1352	23.0					
	07	1353	26.0					
	07	1355	30.0	.510	.37	. 88	3.9	.170
	07	1356	33.0					
_	07	1400	36.0	2.10	.50	2.6	12	.640
	ctober 1		1 60	1.50		67	2.0	050
	05	1225	1.60	.150	.47	. 67	3.0	.050
	05	1226 1230	3.30	150	40	67	2 0	030
	05 05	1231	6.60 9.80	.150	.48	. 67	3.0	.030
	05	1232	13.2					
	05	1235	16.4	.140	.47	. 64	2.8	.040
	05	1236	19.8					
	05	1237	23.0					
	05	1238	26.0					
	05	1239	30.0					1.20
	05	1240	33.0	.500	.60	1.1	5.0	.130
	05 05	1241 1245	36.0	3.30	1.0	4.3	19	1.10
	21	1130	37.9 1.60	.200	.48	.73	3.2	.030
	21	1131	3.30					.030
	21	1132	6.60					
	21	1135	9.80	.090	.32	. 42	1.9	.030
	21	1136	13.2					
	21	1137	16.0					
	21	1140	19.8	.210	.72	. 99	4.4	.040
	21	1141	23.0					
	21	1142	26.0 29.9	.210	.49	. 76	3.4	.030
	21	1145 1146	29.9 33.0	.210	.49	./6	3.4	.030
	21	1147	36.0					
	21	1150	38.0	.200	.70	.95	4.2	.030
· ·	November							
	16	1 400	1.60					••
	16	1401	3.30					
	16	1402	6.60					
	16	1403	9.80					
	16 16	1404 1405	13.2 16.4					
	16	1405	19.8					
	16	1407	23.0					
	16	1408	26.0					
	16	1409	30.0					
	16	1410	33.0					
	16	1411	36.0					
	16	1412	39.0					
•	April 197 13	1320	3.28					.020
	13	1325	19.7					.020
	13	1330	39.4					.015

Lake site L14--Continued

	1	6•77 1455		27•77 200		27•77 1540		17•77 225		11•77 1340
DIVERSITY: DIVISION •CLASS		0.9		0.9		0.9		1.6		0.3
• ORDER		1.6		1.6		0.9		1.6		1.3
FAMILY		1.6		1.6		1.6		1.6		1.7
••••GENUS		1.6		1.6		1.6		1.6		2.2
ORGANISM	CELLS /ML	PER-	CELLS /ML	PER~	CELLS /ML	PER-	CELLS /ML	PFR- CENT	CELLS /ML	PER- CENT
CHLOROPHYTA (GREEN ALGAE)	, , ,			•1.2	,					
•CHLOROPHYCEAF ••CHLOROCOCCALES										
COELASTRACEAE		-		_		_		_		-
MICRACTINIACEAEMICRACTINIUM		_		33		0		_		_
OOCYSTACEAEANKISTRODESMUS		_		-		_		_		_
CHODATELLA		-		-		0	#	33		-
DICTYOSPHAERIUM		-		-		-		-	450	3
···NEPHROCYTIUM ···SCENEDESMACEAE		-		-		-	•	O		-
CRUCIGENIA SCENEDESMUS		-		-		0		0	240	2
· · · · TETRASTRUM		_		0		-		-		-
ZYGNEMATALES										
DESMIDIACEAE CLOSTERIUM		_		_		-	٠	0		-
COSMARIUM		-	*	0		-		-		-
STAURASTRUM		-		-		-		-	•	0
CHRYSOPHYTA .BACILLARIOPHYCEAE										
CENTRALES										
COSCINODISCACEAE CYCLOTELLA	4	33	ø	0		0		0		_
MELOSIRA		-	. #	33		-		0		-
••••STEPHANODISCUS ••PENNALES		-	*	0		-	•	0		-
FRAGILARIACEAE										
····ASTERIONELLA ····SYNEDRA	*	33	*	33 0	#	33	*	33		-
NITZSCHIACEAE							_			
···NITZSCHIA ···TABELLARIACEAE		-		-		-	*	0		-
TABELLARIA		-	ø	0		-		-		-
.CHRYSOPHYCEAE CHRYSOMONADALES										
OCHROMDNADACEAE DINOBRYON		_		_		_		_		_
				_						
CYANOPHYTA (BLUE-GREEN ALGAE) CYANOPHYCEAE										
CHROCCOCCALES										
CHROCCOCCAEAE AGMENELLUM		_		_		-		_	510	3
ANACYSTIS		-		-		-		-		-
HORMOGONALES NOSTOCACEAE										
ANABAENA		-		-		-	•	0	1400	9
APHANIZOMENON OSCILLATORIACEAE		-	۰	0	#	33		0		-
ARTHROSPIRA		-		-		-		-	160	1
OSCILLATORIA PHORMIMIDIUM	*	33		-	#	33	#	33	5400# 1300	3 5
CHPOCCOCCALES									2000	
CHROCCOCCAEAE GOMPHOSPHAERIA		_		-		-		_	5800#	37
EUGLENOPHYTA (EUGLENOIDS)										
CRYPTOPHYCEAE										
.CRYPTOMONIDALES										
CRYPTOCHRYSIDACEAE ,CHROOMONAS		-	4	0		_		_		_
CRYPTOMONODACEAE						0				
CRYPTOMONAS .EUGLENOPHYCEAE		-		-	*	U		-		-
.EUGLENALES										
EUGLENACEAE EUGLENA		_		_		-		0		_
TRACHELOMONAS		-	•	0		0		0	110	1
PYRRHOPHYTA (FIRE ALGAE)										
DINOPHYCEAF PERIOINIALES										
CERATIACEAE										
CERATIUM		-		-		-		-	*	0
CLENODINITACEAE		_		_		_		0		_
GLENODINIACEAF										
		_		0		_		_	_	_

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued Lake site L14--Continued

		۲۱	iytop i ank i	on ana i	A262			
DOTE TIME	AUG 11	5•77 115		17,77 206		7•77 340		5•77 225
DIVERSITY: DIVISION	(0.0		0.9		0.0		1.6
·CLASS	(0.0		0.9		0.0		1.6 1.6
ORDERFAMILY		0.9 0.9		1.6 1.6		1.5		1.6
GENUS	1	1.6		1.6		2.0		1.6
ORGANISM	CELLS /ML	PER- CFNT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT
CHLOROPHYTA (GREEN ALGAE)								
.CHLOROPHYCEAE CHLOROCOCCALES								
COELASTRACEAE								_
COELASTRUM MICRACTINIACEAE	7	0		-		-		-
MICRACTINIUM OOCYSTACEAE		-		-		-		-
ANKISTRODESMUS	#	0		-		-		-
CHODATELLA DICTYOSPHAERIUM		-		-		-		-
NEPHROCYTIUM		-		-		-		-
SCENEDESMACEAE		_		_		-		_
SCENEDESMUS		-		-		-		-
TETRASTRUM ZYGNEMATALES		-		-		•		-
DESMIDIACEAE						_		_
CLOSTERIUM COSMARIUM	*	0		-		-		-
STAURASTRUM	٠	0		-		-		-
CHRYSOPHYTA								
.RACILLARIOPHYCEAE CENTRALES								
COSCINODISCACEAE								
CYCLOTELLA MELOSIRA		0		-		-		-
STEPHANODISCUS		-		-		-		-
PENNALES Fragilariaceae								
ASTERIONELLA		-		-		-		-
SYNEDRA NITZSCHIACEAE		-		-		-		-
NITZSCHIA		-		-		-		-
TABELLARIACEAE TABELLARIA		_		-		-		-
CHRYSOPHYCEAE								
CHRYSOMONADALES OCHROMONADACEAE								
DINOBRYON		0		-		-	#	33
CYANOPHYTA (BLUE-GREEN ALGAE)								
.CYANOPHYCEAE CHROCCOCCALES								
CHROCCOCCAEAE								
AGMENELLUM ANACYSTIS	#	33 0		-	*	25		-
HORMOGONALES	**	٠						
NOSTOCACEAE ANABAENA	*	0	4	33		_		-
APHANIZOMENON		Ö		-	#	25		-
OSCILLATORIACEAE ARTHROSPIRA		_		-		-		-
OSCILLATORIA	#	33		-		25		-
PHORMIMIDIUM CHROCCOCCALES		-		-		-		-
CHROCCOCCAEAE	ш	22		33		25		33
GOMPHOSPHAERIA	#	33	#		•	25	#	33
EUGLENOPHYTA (EUGLENOIDS)								
CRYPTOPHYCEAECRYPTOMONIDALES								
CRYPTOCHRYSIDACEAE CHROOMONAS		_		-		_		_
CRYPTOMONODACEAE		_						
CRYPTOMONAS .EUGLENOPHYCEAE		-		-		-		-
EUGLENALES								
EUGLENACEAE EUGLENA		_		-		-		-
TRACHELOMONAS	4	0	#	33		-	#	33
PYRRHOPHYTA (FIRE ALGAE)								
.DINOPHYCEAE								
PERIDINIALESCERATIACEAE								
CERATIUM		-		-		-		-
GLENODINIACEAE GLENODINIUM		_		-		-		_
PERIDINIACEAE								_
••••PERIDINIUM		-		-		-		-

^{* =} Present in sample but not a dominant type.
= A dominant type in sample.

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

			<u>Lake</u>	site L15			
							Oxygen,
			Spe-				dis-
		Sam-	cific con-	рH		0xygen,	solved (per-
		pling	duct-	(stand-	Temper-	dis-	cent
	Time	depth	ance	ard	ature	solved	satur-
Date		(ft)	(µS/cm)	units)	(°C)		ation)
			(J= 07 57			X37 × 7	
April 19							
06	1515	1.60	76	8.1	5.0	14.0	112
06	1516	3.30	76	8.1	5.0	13.9	111
06	1517	6.60	76	8.1	5.0	13.9	111
06	1518	9.80	76	8.1	5.0	13.8	111
06	1519	13.2	76	8.1	5.0	13.8	111
06	1520	16.5	76	8.1	5.0	13.8	111
06	1521	20.0	76	8.1	5.0	13.7	111
06	1522	23.1	76	8.1	5.0	13.7	111
06	1523	26.4	76	8.1	5.0	13.7	111
06	1524	30.0	76	8.0	5.0	13.7	111
06	1525	31.4	76	8.0	5.0	13.6	111
27	1615	1.60	86	8.9	12.0	12.8	119
27	1616	3.30	86	8.9	11.5	12.8	117
27	1617	6.60	83	8.5	10.5	12.0	107
27	1618	9.80	83	8.4	10.0	11.9	105
27	1619	13.2	84	8.6	10.0	12.0	106
27	1620	16.5	83	7.6	9.5	11.4	99
27	1621	20.0	83	7.0	9.0	11.0	95
27	1622	23.1	83	6.8	8.5	10.4	88
27	1623	26.4	83	6.8	8.5	10.1	86
27	1624	30.0	84	6.9	8.5	9.2	80
May 1977		30.0		0.3	0.5	3.2	30
17	1250	1.60	91	8.4	17.5	11.8	123
17	1251	3.30	91	8.5	16.0	12.1	122
17	1252	6.60	90	8.7	15.0	12.4	122
17	1253	9.80	88	8.2	13.5	12.0	114
17	1254	13.2	88	7.2	13.0	11.1	105
17	1255	16.5	88	7.0	13.0	10.6	100
17	1256	20.0	88	6.8	13.0	10.1	95
17	1257	23.1	87	6.7	12.5	9.4	87
17	1258	26.4	87	6.8	12.5	9.3	87
June 197							
14	1350	1.60	91	7.4	19.5	10.3	110
14	1352	3.30	91	7.4	18.5	10.4	110
14	1354	6.60	90	7.4	17.5	10.4	107
14	1356	9.80	90	7.4	17.0	10.3	106
14	1358	13.2	90	7.3	16.5	9.5	97
14	1400	16.5	95	7.3	16.0	8.6	87
14	1402	20.0	95	7.3	15.5	6.8	67
14	1404	23.1	95	7.3	14.5	2.5	24
14	1406	26.4	95	7.3	13.5	.0	0
14	1408	28.1	97	7.2	13.0	.0	0
17	1 400	20.1	31	, . 4	13.0		U

Table 11.~-Chemical, physical, and biological characteristics of lake water--Continued

						<u>Lake site</u>	<u> L16</u>					
	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper-	Trans- par- ency (secchi disk)	Oxygen, dis- solved	Oxygen, dis- solved (per- cent satur-	Bicar- bonate FET-fld (mg/L as	Alka- linity field (mg/L as	Carbon dioxide dis- solved (mg/L	Nitro- gen, NO2+NO3 total (mg/L
Oate		(ft)	(µS/cm)	units)	(°C)	(m)	(mg/L)	at ion)	HC 0 3)	CACO3) as	s CO2)	as N)
April 19	1530	1.60	76	٥ ۸	5.0	1.30	14.1	110	20	16	.3	.21
06		3.30	76 76	8.0 8.0			14.1	109		10		
06	1531 1535	6.60	76 76	8.0	5.0 5.0			109	19	16	.3	.21
06			76 76				14.0 14.0		19	10		
06	1536	9.80		8.0	5.0			109	19	16		.21
06	1540	11.0	77	8.1			13.9	109				
06	1541	13.0			5.0	1 60						- 10
27	1630	1.60	85	8.9	11.5	1.60	12.9	118	21	17	.0	
27	1633	3.30	84	8.9	11.0		12.8	116				
27	1635	6.60	84	8.9	10.5		12.6	113	19	16	.0	
27	1637	9.80	83	8.8	10.5		12.5	112				
27	1640	11.6	82	8.7	10.0		12.2	108	20	16	.0	.01
May 1977												
04	1350	10.0										
17	1310	1.60	91	8.5	17.5	2.00		123	24	20	.1	< .10
17	1311	3.30	91	8.5	17.5		11.9	124				
17	1315	4.90							23	19		<.10
17	1316	6.60	89	8.6	15.0		12.2	120				
17	1320	9.80	88	7.8	14.0		11.2	108	24	20	.6	< .10
June 197												
14	1405	1.60	92	7.3	19.0	4.0	10.2	109		19	1.8	<.10
14	1407	3.30	92	7.3	18.5		10.1	107				
14	1410	8.20	91	7.3	18.0		10.0	105		18	1.7	<.10
14	1412	9.80	91	7.2	17.0		8.4	87				
14	1415	11.5	90	7.2	17.0		8.7	90		18	2.2	
July 197		11.5	30	,	17.10		0.,	30				0
11	1420	1.60	97	7.9	24.0	2.70	10.6	125		19	.5	<.10
11	1422	3.30	97	7.9	24.0	2.70	10.8					
11	1424	6.60	96	7.9	23.5		10.0					
11	1425	7.40		7.9	23.3		10.0			19		
11	1425	9.80	94	7.7	23.0		8.8					
11	1427	13.0	94	7.7	23.0		6.2			19	.7	
August 1		13.0	94	/./	21.3		0.2	70		19	.,	.01
05	1125							5				
		1.60	96	7.5	24.5		9.2			20	1.2	
05	1200											
05	1203	3.30	96	7.5	24.5		9.2			10	2.0	
05	1205	6.60	96	7.1	24.0		9.2			19	2.9	
05	1207	9.80	96	7.0	24.0		9.0					
05	1210	11.0	95	6.9	23.0		6.8	78		19	4.6	<.10
Septembe		1.65										
07	1405	1.60	90		24.0					20		
07	1406	3.30	90		24.0		8.6					
07	1410	4.10								20		
07	1415	6.60	90		24.0		8.6	101		20		.00
October												
05	1255	1.60	109		16.5	2.00	7.8	79		24		.03

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

	Time	Sam- pling depth	Spe- cific con- duct- ance	pH (stand- ard	Temper- ature	Trans- par- ency (secchi disk)	Oxygen, dis- solved	Oxygen, dis- solved (per- cent satur-	Alka- linity field (mg/L as	Carbon dioxide dis- solved (mg/L	Nitro- gen, NO2+NO3 total (mg/L
Date		(ft)	(µS/cm)	units)	(°C)	(m)	(mg/L)	at ion)	CACO3) a	s CO2)	as N)
October	1977										
05	1256	3.30	111		16.0		7.9	80			
05	1300	5.70							24		.03
05	1301	6.60	111		16.0		7.8	79			
05	1305	9.80	111		16.0		7.7	78	24		.03
21	1155	1.60	92	7.1	12.5	1.80	7.1	66	23	3.5	.06
21	1156	3.30	92	7.1	12.5		7.1	66			
21	1200	5.70							24		.06
21	1201	6.60	92	7.1	12.5		7.1	66			
21	1205	9.80	92	7.1	12.0		7.0	64	23	3.5	.06
November	1977										
16	1410	1.60	85	7.2	9.5		4.7	40			
16	1411	3.30	85	7.2	9.5		4.6	39			
16	1412	6.60	85	7.2	9.5		4.6	39			
16	1413	9.80	85	7.2	9.5		4.6	39			
16	1414	13.2	85	7.2	9.5		4.9	44			

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

			Ī	ake site	16Cont	inued			
Date	Time	Sam- pling depth (ft)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)	Phos- phorus, ortho, total (mg/L as P)	Carbon, organic total (mg/L as C)
April 19	77								
06	1530	1,60	.060	.34	.61	2.7	.020	<.010	4.1
06	1531	3.30							
06	1535	6.60	.080	. 32	.61	2.7	.020	<.010	
06	1536	9.80							
06	1540	11.0	.070	. 43	.71	3.1	.020	<.010	4.0
06	1541	13.0							
27	1630	1.60	.040	. 48	.52	2.3	.020	<.010	
27	1633	3.30							
27	1635	6,60	.040	. 39	.44	1.9	.020	<.010	
27 27	1637 1640	9.80	.050	. 49	.55	2.4	.020	<.010	
May 1977	1040	11.6	.030	. 43	.33	2.4	.020	1.010	
04	1350	10.0							
17	1310	1.60	.010	.36	.37	1.6	.010	<.010	
17	1311	3.30							
17	1315	4.90	.010	. 39	.40	1.8	.010	<.010	
17	1316	6.60							
17	1320	9.80	.050	. 44	.49	2.2	.030	<.010	
June 197									
14	1405	1.60	.030	. 27	.30	1.3	.010	<.010	
14	1407	3.30							
14	1410	8.20	.080	. 28	.36	1.6	. 030	<.010	
14	1412	9.80		20	20	1 7	040	010	
14 July 197	1415	11.5	.080	. 30	.38	1.7	.040	<.010	
11	1420	1.60	.030	. 58	.61	2.7	.020		
11	1422	3.30	.050		.01	2.7			
11	1424	6,60							
11	1425	7.40	.060	. 42	.48	2.1	.020		
11	1427	9.80							
11	1430	13.0	.070	. 64	.72	3.2	.030		
August 1									
05	1200	1.60	.010	. 41	.42	1.9	.020		1.8
05	1203	3.30	01.0			2	020		6 7
05	1205	6.60	.010	. 56	.57	2.5	.020		6.7
05 05	1207 1210	9.80 11.0	.020	. 58	.60	2.7	.020		9.6
Septembe		11.0	. 02.0		.00	2.7	. 02.0		3.0
07	1405	1.60	.010	. 36	.37	1.6	. 020		
07	1406	3,30							
07	1410	4.10	.020	. 31	.34	1.5	.020		
07	1415	6.60	.030	. 32	.36	1.6	.040		
October 0	1977								
05	1255	1.60	.120	. 32	.47	2.1	.060		
05	1256	3.30							

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued

Date	Time	Sam- pling depth (ft)	Nitro- gen, ammonia dis- solved (mg/L as N)	Nitro- gen, organic total (mg/L as N)	Nitro- gen, total (mg/L as N)	Nitro- gen, total (mg/L as NO3)	Phos- phorus, total (mg/L as P)
October	1977						
05	1300	5.70	.140	.47	. 64	2.8	.080
05	1301	6.60					
05	1305	9.80	.130	.46	. 62	2.7	.120
21	1155	1.60	.200	.57	.83	3.7	.030
21	1156	3.30					
21	1200	5.70	.180	.64	.88	3.9	.030
21	1201	6.60					
21	1205	9.80	.200	.51	.77	3.4	.040
Novembe					-		
16	1410	1.60					
16	1411	3.30					
16	1412	6.60					
16	1413	9.80					
16	1414	13.2					

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued Lake site L16--Continued

DATE TIME.		6•77 530		27•77 200		27•77 630		17•77 310		14•77 405		11•77 420
DIVERSITY: DIVISION	-	0.9		0.9		0.0		1.0		1.0		0.7
•CL ASS		0.9		0.9		0.0		1.0		1.0 1.5		0.7
ORDER FAMILY		1.6 1.6		1.6 1.6		0.0 0.0		1.0		1.5		1.6 2.0
GENUS		1.6		1.6		0.0		1.0		5.0		2.0
	CELLS	PER-	CELLS	PER-	CELLS	PER-	CELLS	PER-	CELLS	PER-	CELLS	PER-
ORGANISM	/ML	CENT	/ML	CENT	/ML	CENT	/ML	CENT	/ML	CENT	/ML	CENT
CHLOROPHYTA (GREEN ALGAE) •CHLOROPHYCEAE												
• • CHLOROCOCCALES												
•••CHARACIACEAE •••SCHROEDERIA		-		-		-		-		0	*	0
COELASTRACEAECOELASTRUM		_		_		_		_		-		-
MICRACTINIACEAE			4	0			*	0		_		_
MICRACTINIUMOOCYSTACEAE		-		U		-						-
CHODATELLADICTYOSPHAERIUM		-		-		-	*	0		-	1300	13
••••TETRAEDRON	*	0		-		-	4	ō		-		-
•••SCENEDESMACEAE •••SCENEDESMUS		-	*	0		_	*	0		-		-
TETRASTRUM		-	#	0		-		-		-		-
••TETRASPORALES •••PALMELLACEAE												
••••SPHAEROCYSTIS ••ZYGNEMATALES		-		-		-		-		-		-
DESMIDIACEAE												•
····COSMARIUM ····STAURASTRUM		-		-		-		-		0		0 0
CHRYSOPHYTA												
.BACILLARIOPHYCEAE												
CENTRALESCOSCINDDISCACEAE												
CYCLOTELLA		33	*	0		-	*	0		-		-
····MELOSIRA ··PENNALES	_	v	-	•				·				
FRAGILARIACEAEASTERIONELLA	4	33	*	33	#	100	*	50	#	25		-
FRAGILARIA	*	0		-		-		-	#	25		•
SYNEDRANITZSCHIACEAE		-		U		-	-	U		_		_
NITZSCHIA TABELLARIACEAE	*	0		-		-		-		-		-
TABELLARIA		-		0		-		0		-		-
• CHRYSOPHYCEAE • • CHRYSOMONADALES												
OCHROMONADACEAE		_		_		_		_		_		_
DINOBRYON		-		-		-		_				
CYANOPHYTA (BLUE~GREEN ALGAE) CYANOPHYCEAE CHROCOCCALES												
CHROCCOCCALESCHROCCOCCAEAE												
••••ANACYSTIS ••HORMOGONALES		-	#	33		-		-	#	25		-
NOSTOCACEAE									_	26	1200	12
····ANABAENA ····APHANIZOMENON	*	0		33		-	*	0	"	<u>-</u> S2	1200	12
•••OSCILLATORIACEAE •••OSCILLATORIA	*	33		-		_	#	50		_	3700#	36
CHROCCOCCALES	Ţ	. 33						30			3,000	-
CHROCCOCCAEAEGOMPHOSPHAERIA		-		-		-		-	•	0	3800#	37
EUGLENOPHYTA (EUGLENOIDS)												
.CRYPTOPHYCEAE												
CRYPTOMONIDALESCRYPTOCHRYSIDACEAE												
CHROOMONAS .EUGLENOPHYCEAE		-	*	0		-		-		-		-
EUGLENALES												
EUGLENACEAETRACHELOMONAS		_		-		-		0	9	0	68	1
PYRRHOPHYTA (FIRE ALGAE)												
•DINOPHYCEAE												
CERATIACEAE												
CERATIUMGLENODINIACEAE		-		-		-		-		0	130	1
GLENODINIUM		-		-		-	*	0		-		-
* = Present in sample but not a domin	ant type.											

^{* =} Present in sample but not a dominant type.
= A dominant type in sample.

Table 11.--Chemical, physical, and biological characteristics of lake water--Continued Lake site L16--Continued

	onytoplankton analyses												
DATE TIME		5•77 200		17,77 215		7•77 405		5,77 255		21•77 155			
DIVERSITY: DIVISION .CLASS .ORDERFAMILYGENUS		0.0 0.0 0.9 0.9		0.8 0.8 1.5 1.5		0.0 0.0 1.0 1.0		1.6 1.6 1.6 1.6		0.9 0.9 1.6 1.6			
ORGANISM	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT	CELLS /ML	PER- CENT			
CHLOROPHYTA (GREEN ALGAE) •CHLOROPHYCEAE ••CHLOROCOCCALES													
CHARACIACEAESCHROEDERIA		-		-		-		-		-			
COELASTRACEAE		0		-		-		-		-			
MICRACTINIACEAEMICRACTINIUM		-		-		_		-		-			
CHODATELLA		_		_		_		_		_			
DICTYOSPHAERIUMTETRAEDRON		0		-		-		-		-			
SCENEDESMACEAE		_						_		_			
SCENEDESMUS TETRASTRUM TETRASPORALES		Ξ		-		-		Ξ		=			
PALMELLACEAESPHAEROCYSTISZYGNEMATALES		0		-		-		-		-			
DESMIDIACEAE COSMARIUM		0		_		-		_		_			
· · · · STAURASTRUM	*	Ö		-		-		-		-			
CHRYSOPHYTA .RACILLARIOPHYCEAECENTRALES													
COSCINODISCACEAECYCLOTELLA		_		-		_		_		_			
MELOSIRA PENNALES		-		-		-		-	#	33			
FRAGILARIACEAE ASTERIONELLA		0		_		_		_		_			
FRAGILARIA		-		-		-		-		-			
SYNEDRA NITZSCHIACEAE	-	_		_		-		-		-			
NITZSCHIA TABELLARIACEAE		•		-		-		•		-			
TABELLARIA .CHRYSOPHYCEAE		-		-		-		-	#	33			
OCHRYSOMONADALESOCHROMONADACEAEDINOBRYON		-		-		-	*	33		_			
CYANOPHYTA (BLUE-GREEN ALGAE)													
.CHROCCOCCALES													
CHROCCOCCAEAEANACYSTISHORMOGDNALES	*	33		-	#	25		-		-			
ANABAENA	#	33	*	25	*	25		_		_			
APHANIZOMENONOSCILLATORIACEAE		-	#		#			-		-			
OSCILLATORIA CHROCCOCCALES		-		-		-		-		-			
CHROCCOCCAEAEGOMPHOSPHAERIA	#	33	*	25	#	25		-		-			
EUGLENOPHYTA (EUGLENOIOS) •CRYPTOPHYCEAE													
CRYPTOMONIDALESCRYPTOCHRYSIDACEAECHROOMONAS .FUGLENOPHYCEAE		-		-		-		-		-			
EUGLENALESEUGLENACEAETRACHELOMONAS	•	0	#	25		_	a	33	#	33			
PYRRHOPHYTA (FIRE ALGAE) .DINOPHYCEAE													
CEPATIACEAE								22					
CERATIUMGLENODINIACEAE		-		-		-	#	33		-			
* = Present in sample but not a domination	ent type	-		-		-		-		-			
# = A dominant type in sample.	o cypc.												

⁹⁷